3 1761 06238777 4

MANUAL
OF
OTOLOGY

Presented to
The Library
of the
University of Toronto
by

DR. E.S. RYERSON

Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

http://www.archive.org/details/manualofotology00bacouoft

MEar
B

A MANUAL

OF

OTOLOGY.

BY

GORHAM BACON A.B., M.D.,

PROFESSOR OF OTOLOGY IN THE COLLEGE OF PHYSICIANS AND SURGEONS, COLUMBIA UNIVERSITY, NEW YORK; AURAL SURGEON, NEW YORK EYE AND EAR INFIRMARY.

WITH AN INTRODUCTORY CHAPTER.

BY

CLARENCE JOHN BLAKE, M.D.,

PROFESSOR OF OTOLOGY IN HARVARD UNIVERSITY.

FOURTH EDITION, REVISED AND ENLARGED.

WITH 134 ILLUSTRATIONS AND 11 PLATES.

497368
20.9.49

LEA BROTHERS & CO.,

NEW YORK AND PHILADELPHIA.

1906.

Entered according to Act of Congress, in the year 1906, by
LEA BROTHERS & CO.,
In the Office of the Librarian of Congress. All rights reserved.

WESTCOTT & THOMSON,
ELECTROTYPERS, PHILADA.

PRESS OF
WILLIAM J. DORNAN. PHILADA

PREFACE TO THE FOURTH EDITION.

From the demand for a new edition the author is gratified to infer that the work has attained its primary purpose as a student's text-book, and that perhaps it has been of service to not a few of his professional brethren in general practice.

The revision has been thorough and interstitial, resulting in an increase of about forty pages, the replacement of many illustrations, and the addition of several new plates.

Among the new topics considered may be mentioned osteomyelitis, primary jugular bulb thrombosis and suppurative inflammation of the labyrinth. The paragraphs in the third edition relating to leucocytosis, lumbar puncture and the treatment of facial paralysis have been rewritten and additions made thereto.

In view of the importance of the examination of pus, especially in suppurative middle ear disease and its complications, an appendix has been added in which the method of preparing smears from pus, the making of cultivations and the physiological inoculation experiment have been considered.

The author wishes again to express indebtedness to his colleague, Dr. Clarence John Blake, of Boston, for the introductory chapter, and especially to his assistant Dr. George Sloan Dixon for his valuable aid in the work of revision, and preparation of the new illustrations.

G. B.

47 West Fifty-fourth Street,
New York.

CONTENTS.

PAGE

INTRODUCTION 17

CHAPTER I.

Anatomy and Physiology of the Ear 30

CHAPTER II.

Methods of Examination of the Ear 67

CHAPTER III.

Diseases of the Auricle 100

CHAPTER IV.

Diseases of the External Auditory Canal 119

CHAPTER V.

Diseases and Injuries of the Drumhead and Middle Ear 148

CHAPTER VI.

Otitis Media Purulenta Acuta, or Acute Purulent Otitis Media 164

CHAPTER VII.

Adenoid Growths. Enlarged Tonsils. Diseases of the Nasal Passages 185

CHAPTER VIII.

PAGE

Otitis Media Catarrhalis Chronica, or Chronic Catarrhal Otitis Media 219

CHAPTER IX.

Otitis Media Purulenta Chronica, or Chronic Purulent Otitis Media, or Chronic Suppurative Inflammation of the Middle Ear 257

CHAPTER X.

Granulations and Polypi—Caries and Necrosis of the Temporal Bone 289

CHAPTER XI.

Diseases of the Mastoid Process 310

CHAPTER XII.

Intracranial Complications 370

CHAPTER XIII.

Diseases of the Sound-perceiving Apparatus 436

CHAPTER XIV.

Deaf-mutism 458

APPENDIX 469

INDEX 473

MANUAL OF OTOLOGY.

INTRODUCTION.

The growth of medical education during the past thirty years, shown in no way more markedly than by the establishment of special lines of research, has resulted in the practical advance of the so-called specialties in medicine, and, as a necessary result, an extension of medical school work, to enable it to include for the student such information in the special branches of his profession as he may be able to apply in the course of a general practice.

The extent of this instruction is limited both by the time which may be allotted to it in deference to the demands of other departments and by the ease or difficulty attending the particular form of clinical instruction.

The inaccessibility of all but the outermost parts of the organ of hearing to other than individual inspection, and the difficulty, therefore, of demonstration of diseases of the deeper portions of the ear, except to small classes or sections of a class, greatly limit the amount of instruction in clinical otology which may be given in the usual medical school course. The supplementing of undergraduate teaching by graduate instruction in the special branches of medicine has done something to com-

pensate for this defect, but has opened the door to an evil in the establishment of short courses in otology, as in other departments, which can, in a limited time, afford only a little knowledge and enable the superficial student to assume that he has acquired an ability in special practice which could really be obtained only through long and serious study.

The difficulty here presented, which can be met only by a considerable extension of the medical school curriculum and by the establishment of graduate departments of medical schools and graduate schools giving only long courses to small classes or to individual students, and including abundance of clinical material with adequate opportunities for continual observation of cases, is in a measure temporarily provided for by the publication of elaborate text-books, carefully compiled and richly illustrated. The demand for such text-books has resulted in their numerous production, and at the same time has awakened in both the student and the practitioner the desire for more compact compilations which afford in a small and conveniently accessible compass the information most desirable from a practical point of view. In a field comprising, as does otology, so much of interest and importance, and so many minute definitions, the selection and condensation of the subject-matter are a task of no little difficulty, and one which is to be accomplished only by the presentation of some of the subjects in a merely suggestive form, in order that space may be accorded for the necessary elaboration of others.

The book here presented has this aim in view, and, in addition to a text covering the more important points in regard to which the student and practitioner require

immediate information, it is so compiled as to be a text-book of moderate dimensions and a book of ready reference.

So intricate is the structure of the temporal bone and so important the study of the relationships of its component parts, not only to the understanding of the physiology of the organ of hearing, but also as a preparation for intelligent surgical treatment of its diseases, that even the most careful and painstaking reading must be supplemented or accompanied by a study of sections of the bone itself in order to gain an accurate knowledge of its anatomy.

If, in addition to this, the student himself makes sections of the temporal bone, it will not only render his knowledge more exact, but the manual process of section-making will help to fix in his memory dimensions and points of relationship which as a surgeon it will be necessary for him to bear constantly in mind.

In that important portion of clinical work in otology which includes hearing-test, such knowledge of acoustics as is necessary to an understanding of the process of sound-transmission within the ear itself is of importance to the student and a necessity to the special practitioner, and to this may well be added such further knowledge of the general subject as may form a basis for original research.

This is to be sought either in collateral reading or in lectures on acoustics; and a course of instruction in otology to be complete should include not only individual study of the minute anatomy of the temporal bone, but a demonstrative course in experimental acoustics as well, in order to lay a proper foundation for the intelligent clinical study of diseases of the ear.

In the objective study of his patients the student cannot impress upon himself too forcibly the importance of accurate and repeated observation; the field for inspection is so small and the variations in appearance in disease so manifold that differential diagnosis is often most difficult, and inspection must be supplemented by careful revision of the history of the individual case and of its cognate symptoms.

No amount of reading can compensate for the lack of personal instruction and individual observation, but to the student and the practitioner, to whom this book is commended for the condensed information which it contains, it will be found also to bring that stimulation to further study which is one of the purposes of all teaching.

The training of hand and eye essential to manipulation and observation in so minute a field as that presented by the human ear is enough in itself to constitute this branch of surgical work a special one, and the general practitioner, covering as he does a wider field, cannot expect to do more than to draw general conclusions and to determine the limit beyond which he cannot with safety go, and in which he must call for assistance upon his specially educated colleague.

That this sort of definition of his possibilities by the general practitioner is very necessary is shown by the constantly recurring cases in aural clinics and private practice of patients who have been treated too much or unskilfully for a simple aural disease, or in whom a serious aural trouble has been neglected for want of proper recognition.

To this state of things the inadequate instruction in otology in the undergraduate departments of many of

our medical schools, and the short courses of instruction to graduates, which apparently promise a speedy equipment, have much contributed; while, on the other hand, the eagerness of the special practitioner to establish himself remuneratively tends to put the question of his more serious and scientific education into the background.

The remedy, inevitable in the course of time with the advance of medical education, may be furthered by limiting the instruction of the undergraduate student to such points as it may be necessary for him to know in order for him to meet successfully the average aural cases coming to him in general practice, and by making his instruction largely clinical; while of the special student there should be required at least a year of additional study and practical experience in some one of the special aural hospitals or hospital aural departments.

Short courses to graduates, except within the limits above defined, should as a whole be discouraged; while for the undergraduate, or graduate intending to enter special practice or desiring something more than the superficial knowledge which has within it an element of danger, long courses properly graded, and including a maximum of clinical training, should be required.

A knowledge of the appearance of the normal external auditory canal and drumhead is an essential prerequisite to the appreciation of the value of the objective symptoms of disease of these and deeper-seated parts, and this knowledge the general practitioner who has been instructed in methods of examination can, in time, acquire by his own observation; and in the cases of patients applying for relief on account of symptoms occurring in one ear only, it is well to make it a rule to examine both ears, the normal ear not only affording a

standard of comparison in the individual case, but refreshing the memory of the observer with regard to the normal appearance. The differences of contour of the external canal and the variations in the surface of the drumhead within normal limits, though slight, are so many that the superposition of many mental pictures is necessary to the attainment of the composite picture which shall represent the normal average.

In the diseases of children especially, a familiarity with the methods of examination of the ear, based upon previous instruction and self-acquired in the course of a general practice, will often enable the physician to determine for himself that the ear is not a factor in the causation of perplexing symptoms or that its condition is one in which special attention is demanded, and he may be able to hold himself free from the blame emphasized by the late Edward H. Clark, and approvingly quoted by von Troeltsch, to the effect that "the physician who fails to examine the ears of a child in the course of one of the acute exanthemata may justly be denominated an unscrupulous practitioner."

Very nearly 25 per cent. of the patients attending our aural clinics are children under fourteen years of age; of this number, 50 per cent. are the subjects of suppurative inflammation of the middle ear, and of this number 10 per cent. or more owe their origin to the exanthemata of childhood; while of the cases of deaf-mutism throughout the United States, 27 per cent. have been the result of suppurative middle-ear disease in early childhood, and it is probable that the more thorough investigation and the systematic examination of the ears of children in our deaf-mute schools will show a still larger percentage of these cases.

Of the diseases of the ear which come under the consideration of the general practitioner, and which do not menace life, the first in the list is the so-called chronic catarrhal inflammation of the middle ear, which makes in the temperate zone about 30 per cent. of all the material ultimately referred to the aural specialist, and includes a series of changes in the soft tissues of the sound-transmitting apparatus which are still the subject of interested study and investigation; and one of the many fields of research open to the student of the future in otology is the further acquired differentiation of those alterations in structure, the result of disease, which interfere with the transmission of sound, and the etiology of which is often to be traced to a remote source. As an example of this, may be taken the peculiar and often puzzling cases of variable hearing occurring in young women who ultimately become very deaf to all sounds aërially conveyed. The great variations, occurring as they did in girls soon after puberty, or in young women who were the subjects of considerable nervous and emotional strain, led, without other ascertainable cause, to the conclusion that these variations were purely of neural origin, and, being merely symptomatic, were to be dealt with solely through attention to the general condition. A special study of these cases, however, through the aural speculum, showed, under conditions of the greatest hearing impairment, in many of them, a localized congestion of the tympanic mucous membrane, especially of the niche of the oval window around the stapes. This peculiar blush was found most commonly, periodically, in connection with disturbances of the menstrual function, or with other evidences of pelvic disorder, and, with the assistance of the gynecologist, it was found,

in the great majority of cases, that there was either malposition of the uterus, erosion of its neck, or atresia of its cervical canal, and that the remedying of these defects had a very material and beneficial influence upon the localized circulatory disturbance, and accompanying tissue-change, in so important a portion of the middle ear as is the inner end of the sound-transmitting chain of bones.

Another example is shown in the cases of chronic non-suppurative middle-ear disease in which the deleterious changes in the lining membrane of the tympanic cavity are furthered by general nervous overstrain, overtire, and the coincident train of disturbances of nutrition. In these cases the local treatment of the ear is futile, or of but little and temporary benefit, without careful attention is given to general hygiene and to a consideration of the condition of environment of the patient, climatic or otherwise. The dependence upon local treatment alone in such cases is sometimes mischievous in its results, and at least fails of the benefit which might accrue from a broader view of the situation, and emphasizes the importance of a consultation between the aurist and the physician, who as the family adviser has cognizance of conditions which may have an important bearing upon the etiology and progress of the case in question.

The first tendency in specialism is to lay stress upon the special character of the investigation to be followed and to emphasize its individuality, and with the broadening of the study, its relationship to and dependence upon general scientific research becomes more and more evident. Specialized specialism ceases to exist in proportion as it separates itself, and grows and broadens

only as it keeps up its relationship with the main body; and here the military simile may be employed for the specialist, who, as a skirmisher exploring unknown country, cuts himself off from the main body which is his support, and to which he owes his allegiance, and who, seeking to exist to himself alone, fails of that usefulness in the line of duty which is the invariable price of any existence whatever.

One of the most striking examples of elucidative work in otology is that which has had to do with the definition and explanation of the curious complex of symptoms of which vertigo is the most marked and incapacitating manifestation.

With the recognition of the ear as a peripheral organ of equilibration, and with increased knowledge of its reflex relationship to the sympathetic nervous system, the better definition of its blood-supply, and the differentiation of its vasomotor tracts, there is an explanation of much that has been vague and mystifying in those cases of vertigo of sudden onset, formerly referred either to abnormalities in the circulation in the brain or to disturbances of the digestive apparatus, as their amenable cause; and the advances made in the study of the physiology of the sound-transmitting apparatus, and in the application of surgical rules to the treatment of diseases of the middle ear, has brought to our knowledge the possible advantage of establishing still another differentiation, that of *aural* as distinguished from *auditory* vertigo.

An effusion or hemorrhage occurring in the labyrinth may affect the hearing or the balancing power, or both, of the individual, to a greater or less degree, according to its location and extent, and the symptoms, so

far as equilibration is concerned, may gradually disappear, to be reawakened at a later period by general conditions which induce a suspense of vasomotor inhibition, and thereby a secondary irritation of the terminal nerve apparatus. It is further established that a fixation of the sound-transmitting apparatus of the middle ear, by doing away with or decreasing the function of one of the outlets from the labyrinth for pressure exerted by dilatation of the intralabyrinthine vessels, favors disturbance of the nervous end apparatus by a lesser degree of circulatory pressure than would be possible were the elasticity of the membranes of the round and oval windows and the mobility of the ossicular chain unimpaired.

Aural records show an increasing observation of cases of suppurative disease of the middle ear in which pressure upon the stapes, as the result of a new growth, or of epidermal exfoliation, causes a series of symptoms very closely resembling those of the so-called Ménière's complex, and in which vertigo plays a prominent and disturbing part. These cases are almost invariably relieved by the removal of the morbid growths or accumulations, and the consequent release of the labyrinth from extrinsic mechanical pressure, and constitute the class of what may, therefore, be called the mechanical or aural vertigoes. For purposes of distinction, the first series of cases just cited may be termed auditory or labyrinthine vertigo, and the second class may be considered as lying between these two extremes. Beyond this grosser differentiation it is possible to go into minute definitions as to immediate or reflex cause and effect, and to follow the subject into the domain of the neurologist, before it shall in any degree have received

the attention in investigation which its importance demands; while the question of intralabyrinthine lesions as an accompaniment of diseases of the kidneys, of the digestive apparatus, of the pelvic organs in women, and of the constitutional diseases, of which syphilis stands at the head of the list in the number of cases of labyrinth implication, and of diseases of the internal ear in children, presents problems toward the solution of which the general practitioner and the otologist may well take counsel together.

There is one other subject which should command the attention both of the aurist and the general practitioner, and that is what may be called the "fatigue of deafness," a common and often unestimated factor in the causation and persistence of neurasthenia, and one imposing upon sometimes feeble shoulders a burden which it becomes us to lighten if we cannot lift.

It may be taken as an accepted fact, among other evidences of the bounty of nature, that those of us who have normal ears start in life with double the amount of hearing absolutely necessary for the ordinary purposes of social communion, and it is, therefore, possible to lose one-half of one's hearing power before becoming conscious of the loss. Under these conditions, where the daily ordinary use of the hearing is regarded as the standard of perfection, it is easily understandable that a chronic progressive disease impairing the hearing may become fairly established before attention is turned to the necessity of interference, with a view to prevention and repair; and it is, therefore, not unusual to find in the majority of the chronic progressive diseases, of which the slow impairment of hearing is the important symptom, that they have by the time they come into the

hands of the aurist long passed the possibility of prevention, and have established a condition which can be repaired in only a moderate degree. The apparent hopelessness of many of these cases, the impossibility of effecting by the treatment more than a slight improvement, should be no deterrent to the honest and strenuous effort to ameliorate the condition of cases of this class; and it is in many of them that the general practitioner, the family physician, and the specialist can best work hand in hand. The deleteriously causative and influencing factors in the general condition in these cases are so many, and the progress toward improvement so slow, that the local treatment may often advisedly be limited to that which the patient can himself effect under the observation of his physician, with occasional examination and suggestion from the specialist; and there is probably no class of cases, although coming under the care of the aurist, which is so likely to become the subject of what may be termed overtreatment, either surgical or medical, as the chronic non-suppurative diseases of the middle ear. It is especially in connection with these cases of slowly progressive impairment of hearing that the fatigue of deafness manifests itself. Few intelligent observers outside of the very deaf themselves, or of those who have to deal much with persons so afflicted, can appreciate the profound exhaustion resulting from the effort to compensate for a deficiency in this particular line of communication with the outer world in the person whose perception of that mode of motion to which we give the name of sound was once made without appreciable effort, and who has, under conditions of impaired hearing, first of all to make an effort to hear, and, in default of hearing, an effort to

appreciate vocal utterances by watching the motion of the lips of the speaker, and finally, in default, either through lack of perceptive power or through ability only to catch consonant sounds which are formed in front of the mouth, to solve the puzzle of the spoken sentence by filling in the missing consonant sounds, those imperfectly heard, or inadequately seen, from the appreciated context of the sentence; so that where once understanding came without effort, three distinct and appreciable brain efforts are required: the first, to hear; the second, to see; and the third, to understand. In our estimate of the burdens of physical life, the consideration of which belongs especially to the ministry of the medical profession, we should endeavor to estimate more justly as scientific observers, more kindly, perhaps, as fellow-creatures, the nervous strain, the restrictive limit to useful expenditure of energy, and the demand for compensatory consideration for the pitiably deaf.

CHAPTER I.

ANATOMY AND PHYSIOLOGY OF THE EAR.

In order to study the anatomy and physiology of the ear in a satisfactory manner, the organ of hearing is divided into (I.) the sound-conducting and (II.) the sound-perceiving apparatus.

I. The sound-conducting apparatus includes
   a. The external ear, consisting of
      1. The auricle (or pinna) and
      2. The external meatus (or external auditory canal).
   b. The middle ear, composed of
      1. The tympanum (or tympanic cavity), with the membrana tympani and the ossicles,
      2. The Eustachian tube, and
      3. The mastoid process.

II. The sound-perceiving apparatus (or internal ear) comprises
      1. The vestibule,
      2. The semicircular canals, and
      3. The cochlea.

The latter receives the peripheral expansions of the auditory nerve, the excitation of which produces in the brain-centre sensations which we call sound.

(*a*) THE EXTERNAL EAR.

The external ear consists of

1. The auricle (or pinna) and
2. The external meatus (or external auditory canal).

1. The Auricle.

The auricle, or pinna, pyriform in shape, is attached to the side of the head by means of ligaments and

FIG. 1.



Auricle. (POLITZER.)

A. Helix. B. Antihelix. C. Tragus. D. Antitragus. E. Lobule. F. Concha. G. Orifice of the external meatus.

muscles. It usually forms an acute angle with the lateral portion of the head, although the angle varies in different individuals. The auricle is composed of yellow or reticulated cartilage, except the lower portion, called the lobule, which is formed mostly of connective tissue, in the meshes of which are globules of fat. The lobule

contains but few bloodvessels and nerves. The cartilage of the auricle, about $\frac{1}{10}$ inch in thickness, is covered by the perichondrium, and outside of the perichondrium is the integument. The latter is more firmly attached to the anterior than to the posterior surface of the cartilage. The anterior or concave portion of the auricle presents a number of ridges and depressions.

The helix (see Fig. 1, A) begins just above the external meatus at a point in the concha called the crista helicis, and extends upward, backward, and downward, and terminates just above the upper portion of the lobule.

The antihelix (B) is another elevation running in a parallel direction with the posterior portion of the helix. It starts by means of two crura above the crista helicis, and, passing backward and downward, terminates in the antitragus, D. Partly closing the external opening of the ear, and opposite the antitragus, is the tragus, C. Between the tragus and the antitragus is a depression called the incisura intertragica. The fossa of the antihelix, or fossa scaphoidea (H), lies between the two crura of the antihelix. There are a few sebaceous glands in the lobule, but they are much more numerous and of larger size in the concha.

There are two sets of muscles connected with the auricle. The first set consists of the

1. Attollens aurem, which takes its origin from the epicranial aponeurosis, and is inserted into the upper convex surface of the auricle. Its action is to support and pull the auricle upward.

2. Attrahens aurem, which also arises from the epicranial aponeurosis, is attached to the crista helicis. It draws the auricle forward and upward.

3. Retrahens aurem, which has its origin from the mastoid process, is inserted into the posterior convex surface of the concha. This muscle pulls the auricle backward, but the action of all these three muscles is

FIG. 2.



The muscles of the pinna. (GRAY.)

very limited. Some persons, however, have the power of moving their ears quite perceptibly.

The second set of muscles, five in number, are muscular fibres lying on the concave and convex surfaces of the auricle (see Fig. 2), and are called the

1. Tragicus.
2. Antitragicus.
3. Helicis major.
4. Helicis minor.

} On the concave surface.

5. Transversus auriculæ.
6. Obliquus auris.

} On the convex surface.

These muscles are rudimentary in man, and have very little action.

There are also various ligaments, some holding the auricle to the side of the head and others connecting the different portions of the cartilage together.

2. The External Meatus, or the External Auditory Canal.

The external meatus, about one inch and a quarter in length, is a continuation inward of the auricle, and consists of a cartilaginous and a bony portion. The cartilaginous portion is not cartilaginous throughout its whole extent, the cartilage being wanting in the upper and posterior portions of the canal, where a membranous layer attached to the lining membrane of the canal takes its place (Fig. 3). The fissures of Santorini, generally two in number, are situated in the lower and anterior wall of the cartilaginous meatus. These fissures, transverse to the long axis of the canal, are openings in the cartilage which are filled in with fibrous tissue, and are of importance to the aural surgeon, because in examinations of the ear they allow a certain straightening of the meatus, and, further, an inflammation of the external auditory canal may extend to the parotid gland through these fissures, and, *vice versa*, in case of parotiditis leading to the formation of an abscess, the pus may force its way through these openings into the external meatus and give rise to an otorrhœa, the source

of which can only accurately be determined by means of the head-mirror and speculum.

FIG. 3.



A front view of the organ of hearing. (Right side.) (GRAY.)

The Osseous Meatus.

The osseous meatus, forming about two-thirds of the entire external auditory canal, consists of a bony ring in the infant (the annulus tympanicus) (Fig. 5, C), which develops later in adult life into a bony canal about three-quarters of an inch in length. In the infant the tympanic ring is attached to the squamous portion of the temporal bone by its two extremities, between which is a space which, later, becomes the Rivinian segment, and connected with this ring is a membranous canal. Occasionally, in the fully developed temporal

bone fissures are observed in the upper and posterior walls of the bony meatus. As prolongations of connective tissue extend into these fissures, it is easy to understand how inflammatory material may pass through these openings and cause caries of the walls of the canal. The cartilaginous canal contains soft hairs and sebaceous and ceruminal glands. A few of the latter are found in the bony portion, especially in its upper wall. The

FIG. 4.



Transverse section through the cartilaginous external auditory canal at birth.

lining membrane of the external meatus is simply a continuation inward of the integument of the auricle, and is not a mucous membrane, as formerly was supposed. The dermal layer in the bony portion is very thin and closely attached to the periosteum, a fact which accounts for the excruciating pain sometimes

observed in acute inflammation of the lining of the external meatus.

The blood-supply for the auricle and external meatus comes from the temporal and internal maxillary arteries.

The anterior or concave surface of the auricle and the outer portion of the external meatus derive their supply from several branches of the superficial temporal artery. The posterior surface is supplied by a branch of the posterior auricular artery (a branch of the external

FIG. 5.



Temporal bone at birth.

A, squama; B, zygoma; C, annulus tympanicus; D, promontory with foramen rotundum; E, foramen ovale.

carotid). The inner portion of the external meatus is supplied by the deep auricular artery (a branch of the internal maxillary artery). The smaller branches of these arteries reach the deeper portions of the external meatus by passing through

1. The fissures of Santorini.

2. Through the fibrous connective tissue which joins the cartilaginous canal to the bony canal, and

3. Through the intermediate membranous layer which covers the margins of the cartilaginous groove.

They form a network of capillary vessels on the perichondrium and about the hair-sacs and sebaceous glands. These vessels are quite numerous about the upper border of the membrana tympani and extend along the posterior portion of the handle of the malleus to its inferior extremity.

The veins supplying the two surfaces of the auricle join the external jugular and mastoid veins, although in some cases there are veins that empty into the internal maxillary and temporal veins. Very little is known about the lymphatics of the ear.

The nerves distributed to the auricle and external meatus are derived

1. From the deep posterior auricular nerve (a branch of the facial), which supplies the posterior surface of the auricle.

2. From the cervical plexus through the auricularis magnus.

3. From the auriculo-temporal (a branch of the trigeminus), which supplies the skin of the auricle and external meatus.

4. From Arnold's nerve (a branch of the pneumogastric), which supplies the lining membrane of the external meatus.

In children, one of the difficulties in the way of a successful examination of the external meatus is the small size of the cartilaginous portion. The size and shape of the external meatus in the adult are subject to great variations. In some persons the cartilaginous canal is very wide, while in others it is extremely narrow. The diameter of the canal, which just inside of the

external orifice is about $\frac{2}{10}$ to $\frac{3}{10}$ inch at first, becomes wider in its passage inward, and then narrows again at the point of junction with the bony canal. From this point toward the membrana tympani the bony canal becomes wider. The narrowest part of the bony canal is called the *isthmus*. It is here that foreign bodies are apt to become wedged, while if they pass beyond this constriction it is sometimes impossible to remove them without displacing the auricle forward, an operation to be described later. The distance from the anterior margin of the membrana tympani to the isthmus is about $\frac{6}{20}$ to $\frac{7}{20}$ inch, while the distance from the posterior margin of the drumhead, along the posterior wall to the isthmus is but $\frac{1}{20}$ to $\frac{2}{20}$ inch. This fact should be borne carefully in mind in the case of impaction of a foreign body, which becomes lodged at this point, for serious injury may be done to the drumhead by means of instruments passed along the posterior wall. The cartilaginous canal, in passing inward, turns backward and upward, while the bony canal extends forward and downward. In examining the ear, after introducing the speculum, the auricle should be pulled upward and backward, in order to straighten the external meatus and thus afford a better view of the drumhead.

(*b*) THE MIDDLE EAR.

The middle ear, as previously mentioned, consists of

1. The tympanum, or tympanic cavity, with the membrana tympani and ossicles,
2. The Eustachian tube, and
3. The mastoid process.

1. The Tympanum, or Tympanic Cavity.

The tympanum, or tympanic cavity, is very irregular in shape. Its greatest diameters are those extending antero-posteriorly and supero-inferiorly. The membrana tympani, or drumhead, forms the greater part of the outer wall of the tympanum, and lies at the inner extremity of the bony auditory canal, being placed almost in a horizontal position in the infant at birth, and gradually becoming more perpendicular in the adult. The membrana tympani receives the sound-waves from the external air and carries them by means of the ossicular chain to the labyrinth. The drumhead in the adult is placed obliquely with the long axis of the bony canal, so that the inferior and the anterior walls of the auditory canal are longer than the superior and posterior walls, an important point to be remembered in the extraction of foreign bodies and in other instrumental manipulations. The membrana tympani at its margin is attached to the groove in the bony canal, called the *sulcus tympanicus*, which belongs to the tympanic ring. At the Rivinian segment, previously mentioned, the drumhead is attached partly to the margo tympanicus and partly to the lining membrane of the bony auditory canal. The normal drumhead is of a pearl or grayish color, somewhat oval in shape and drawn inward, the greatest point of concavity being at the *umbo*, at which point the inferior extremity of the malleus is attached by being enclosed in the layers of the drumhead. (See Fig. A, Plate V.) There is a partial convexity or bulging out of the membrana in its anterior and inferior segments, so that the drumhead, while concave as a whole, is convex from the centre to the periphery. The membrana, which resembles gold-beaters'-skin in appearance, consists of three layers:

1. An external or dermal,
2. A middle or fibrous (lamina propria), and
3. An inner or mucous layer.

The long handle of the malleus lies between these layers and extends downward to the umbo. The short process is recognized as a short, teat-like projection, and is a most important landmark in examinations of the drumhead. Situated just in front of and

Fig. 6.



Outer surface of the left tympanic membrane of an adult. (Politzer.)

V. Segment of the tympanic membrane lying in front of the handle of the malleus. H. Posterior segment of the tympanic membrane. S, S′. Prussak's striæ, passing from the short process of the malleus to the spina tymp. post. et minor. MS. Shrapnell's membrane.

behind this process are two folds, which are very important, for it will be demonstrated in another chapter that when the drumhead is drawn inward these folds stand out very prominently. There are two striæ, first described by Prussak (Fig. 6, S, S′), which lie just inside of the two folds, and which can often be made out in life as two tendinous bands. These striæ, together with the grooveless fissure, bound a triangular-shaped portion of the membrane called the *membrana*

flaccida, or *Shrapnell's membrane* (Fig. 6, MS). This membrane consists of very delicate and loosely arranged fibres of connective tissue, crossing each other irregularly, and covered by a thin dermic layer on the outer and a mucous layer on the inner surface, and is very much thinner than the other portions of the membrana tympani The inner lining of the membrana tympani is a continuation of the mucous membrane of the tympanum. The outer or dermal layer is a continuation inward of the skin covering the external meatus, and contains several strata of pavement epithelium. The middle layer, or lamina propria, consists of two sets of fibrous tissue. The outer fibres are attached to the tip of the malleus and radiate toward the circumference, while the inner fibres are circular in their arrangement. As previously stated, a network of vessels descends behind the long handle of the malleus, and at the centre of the drumhead there is a communication between these arteries and veins with a vascular zone which is situated at the periphery of the membrana, and through this latter zone with the vessels of the middle ear.

The superior wall or roof of the tympanum consists of a thin plate of bone, upon which the middle lobe of the brain rests. In the infant there is a suture (petroso-squamosal suture) in the tympanic roof, through which connective-tissue processes with bloodvessels pass from the dura into the middle ear, a fact which explains why an acute inflammation of the middle ear in an infant is apt to occasion more or less meningeal irritation. In the adult this suture is generally closed, but in some temporal bones the delicate superior wall is defective, there being several openings, an anomaly which is designated by Hyrtl as spontaneous dehiscence of the tegmen

tympani, and is in all probability due to an arrest of development. The inferior wall is much narrower than the superior, and close to it is the *fossa jugularis.* The proximity of this fossa to the middle ear should be remembered, because phlebitis and thrombosis of the jugular vein are complications liable to follow an inflammation of the tympanic cavity.

FIG. 7.

Section of foot-plate of stapes—foramen ovale, showing ligamentous attachment.

The inner wall separates the middle ear from the labyrinth or inner ear, and presents several points of interest in

1. The promontory, which corresponds to the first turn of the cochlea;

2. The fenestra ovalis (see Fig. 7), which contains

the foot-plate of the stapes, and which lies a little above and behind the promontory; below the fenestra ovalis is

3. The fenestra rotunda, closed by a membrane (membrana tympani secundaria). Behind these foramina are seen (Fig. 8).

4. The pyramid, a cone-shaped projection containing the stapedius muscle, and

5. The aqueductus Fallopii, enclosing the portio dura,

FIG. 8.



View of inner wall of tympanum. (Enlarged.) (GRAY.)

or facial nerve, a slight ridge of bone lying along the upper and posterior part of the inner wall. The facial nerve emerges at the stylomastoid foramen.

The upper portion of the tympanic cavity, which contains portions of the malleus and incus, is called the *attic*. The attic opens into the mastoid antrum posteriorly, while the *antrum* is the passage of communication between the mastoid cells and the middle ear. The

short process of the incus lies in the inferior angle of the antrum.

In the anterior wall of the middle ear the bony Eustachian canal is situated. The carotid artery also is in close proximity to the anterior wall, so that in cases of caries or necrosis fatal hemorrhage may occur. The *tensor tympani muscle* lies in a bony canal just above and parallel with the Eustachian tube, and, passing around the processus cochleariformis, is attached to the upper and posterior part of the handle of the malleus.

FIG. 9.



The small bones of the ear, seen from the outside. (Enlarged.) (GRAY.)

The Ossicles.

The ossicles consist of the malleus, incus, and stapes. The malleus has

1. A head which articulates with the incus.
2. A short process (processus brevis), and
3. A long process or handle, or manubrium, and (Fig. 9)
4. A processus gracilis.

The incus, besides

1. The body, which resembles the crown of a molar tooth, has

2. A long process (processus longus), which articulates with the head of the stapes as well as

3. A short process (processus brevis).

The latter is attached by a fan-shaped ligament to the lower angle of the mastoid antrum.

As its name implies, the stapes is shaped like a stirrup. Besides the head, already mentioned, which articulates with the incus, there are two

1. Crura, and

2. A foot-plate, the latter being attached to the foramen ovale by means of the annular ligament.

The *chorda tympani nerve* (see Fig. 75) enters the tympanum through a small canal at the posterior border of the membrana tympani, and, passing between the long process of the incus and the handle of the malleus, emerges at a small opening (canal of Huguier) close to the anterior border of the drumhead and at the inner extremity of the Glaserian fissure. It then passes down between the two pterygoid muscles, and meeting the lingual nerve accompanies it to the submaxillary gland.

The articular portions of the ossicles are connected together by capsular ligaments. In addition to these there are numerous ligaments between the ossicles and the tympanic cavity. The external ligament of the malleus should be mentioned, as it forms the upper boundary of *Prussak's space*. The other boundaries of this cavity, as shown in Fig. 10, are, internally, the neck of the malleus; below, the superior surface of the short process of the malleus, and, externally, the membrana flaccida, or Shrapnell's membrane. The conjoined bones, viz., the malleus and incus, serve to divide the attic into two compartments. The *outer compartment* or *external*

attic, is situated between the articulating bones and the outer bony wall, being bounded above by the superior ligament of the malleus and the superior malleo-incudal fold. This *external attic* is further subdivided into a *superior* and *inferior cavity* (see Fig. 10), the latter being known as Prussak's space, and being bounded superiorly by the external ligament of the malleus and the system of cavities described by Politzer; internally, by the neck

FIG. 10.



Prussak's space, section through the tympanic membrane, malleus, upper and outer tympanic wall of a decalcified preparation. (POLITZER.)

LS. Superior ligament of the malleus. LE. External ligament of the malleus. S. Shrapnell's membrane, or the membrana flaccida. O. Prussak's space. R. System of cavities between the body of the malleus and the incus and the external tympanic wall. T. Tendon of the tensor tympani muscle.

of the malleus; inferiorly, by the short process of the malleus; and externally, by the membrana flaccida. Prussak's space communicates with the upper part of the attic, and also with the posterior portion of the tympanic cavity by a small opening.

A good understanding of the subdivisions of the attic is most important to the aurist, for it is here that the inflammation of the middle ear is occasionally most

marked, leading frequently to caries of the walls and of the ossicles.

The origin of the tensor tympani muscle is

1. From the cartilaginous portion of the Eustachian tube.

2. From the osseous wall of the pyramid close to the carotid canal, and

3. From the bony canal through which it passes.

The tendon curves around a bony process, the processus cochleariformis (see Fig. 8), and is inserted into the upper and posterior portion of the long process of the malleus.

The stapedius muscle arises from the bony pyramid through which it passes, and the tendon is inserted into the neck of the stapes.

Experiments made by Politzer upon dogs proved

1. That the tensor tympani is supplied by the motor portion of the fifth nerve, and

2. That the central fibres of the stapedius muscle are under the control of the facial nerve.

The lining membrane of the middle ear is quite thin, and its epithelium is of the ciliated squamous variety in the upper part of its cavity, while in the lower portion it is ciliated cylindrical. The connective tissue which contains the bloodvessels and nerves serves as a periosteum, and the mucous membrane of the tympanum is a continuation inward of the mucous membrane of the pharynx and Eustachian tube. There are various folds of mucous membrane between the walls and the ossicles. In the newborn infant the lining membrane of the tympanum contains a great many bloodvessels and has a swollen appearance. During fœtal life the tympanum is filled with a gelatinous mass, which degenerates rapidly after birth into a thick fluid, owing to the

entrance of air into the middle ear. This fluid becomes absorbed.

The arteries supplying the tympanum are

1. The tympanic, from the internal maxillary.
2. Several branches from the middle meningeal.
3. A few branches from the internal carotid.
4. The ascending pharyngeal (a branch of the external carotid), and
5. The stylomastoid artery.

The veins supplying the tympanum communicate with those of the external meatus as well as with the plexus in the carotid canal and the plexus of the lower jaw, and further join the veins of the dura mater by passing through the petrososquamosal suture. Anatomical investigations undertaken by Politzer prove that "vascular connections are kept up between the middle ear and the labyrinth through the osseous wall separating them." This point is of importance, because it is easy to understand how involvement of the labyrinth may occur simultaneously with acute inflammation of the middle ear.

The nervous supply is derived from

1. The sympathetic.
2. Sensory fibres of the trifacial, and
3. The glosso-pharyngeal.

Jacobson's nerve, the tympanic branch of the glosso-pharyngeal, enters the tympanic cavity through a small canal in the inferior wall of the jugular fossa, and joins, at the promontory, the small superficial petrosal nerve.

The tympanic plexus lying on the inner wall of the middle ear is composed of

1. Branches of Jacobson's nerve.
2. The small superficial petrosal nerve, and
3. Branches from the carotid plexus.

This plexus distributes branches to the fenestra ovalis, the fenestra rotunda, and the lining membrane of the middle ear and Eustachian tube.

Fig. 11.



Eustachian tube and tympanic cavity. (Politzer.)

A. Membrana tympani. B. Head of the malleus. C. Lower end of the handle of the malleus. D. Body of the incus. E. Short process of the incus. F. Tensor tympani muscle. G. Pharyngeal opening of the tube. H. Isthmus of the tube. I. Tympanic opening of the tube. (Right ear.)

2. The Eustachian Tube.

This tube is named after Eustachius, who described it more minutely than other writers, although Vesalius

had previously discovered it. It is the passage connecting the middle ear with the pharynx, and through it the external air enters the tympanum. It is about one and one-half inches in length. The position of the tube is oblique, extending from the tympanum forward, downward, and inward, while its axis forms an angle of 135° with the horizontal axis of the meatus. It is wider at either extremity than at the junction of the cartilaginous and osseous portions, and is composed of a bony and a cartilaginous canal (Fig. 11).

The *osseous canal*, a continuation inward of the anterior portion of the tympanic cavity, is bounded externally by the tympanic bone, superiorly by the canal of the tensor tympani, inferiorly and to the inner side by the canal for the carotid artery. The average diameter of the tube is about $\frac{1}{12}$ inch, although the greatest diameters vary from $\frac{2}{12}$ to $\frac{3}{12}$ inch, while its smallest normal calibre is large enough ordinarily to admit a small probe.

The *cartilaginous canal*, about two-thirds the whole length of the tube, is attached at its upper end by connective tissue to the irregular margin of the bony canal, and has a funnel-shaped opening into the nasopharynx. The canal is not composed throughout of cartilage, being partly cartilaginous and partly membranous. The posterior and upper walls are cartilaginous. The membranous portion forms the lesser half of the circumference of the tube, while the cartilaginous portion forms the boundary of one side and the superior wall. The calibre of the tube is slightly sigmoid and cleft-like in shape. The cartilaginous portion is stiff enough to prevent complete closure of the lumen of the canal, although the opposite surfaces of the lining membrane touch one another except at a point very near the union of the cartilaginous and bony portions and at the pharyngeal

orifice. The cartilage is composed superficially of a hyaline, but in the deeper part of a fibrous, fundamental substance. Zuckerkandl and Urbantschitsch describe numerous irregular fissure-clefts in the cartilage, and occasionally a division of the cartilage into several separate portions. The opening of the tube in the naso-pharynx lies on the external wall of the latter, just behind the posterior extremity of the inferior turbinated bone, being rounded in the child, but oval-shaped in the adult. The Eustachian tube in children is much wider and shorter, its direction is more horizontal, and it contains less cartilage than in the adult; the opening also into the tympanum is larger, while the pharyngeal orifice is much smaller, and it is easier to inflate the middle ear in children than in adults, while the secretion passes more readily from the middle ear to the pharynx. The mucous membrane lining the Eustachian tube is of a light-yellow color, and projects somewhat at the pharyngeal opening, but is firmly adherent to the deeper tissues. It is of the ciliated cylindrical variety, the cilia moving in a direction from the tympanum toward the pharynx, thus aiding the flow of mucus or of fluid from the middle ear or tube toward the pharynx. The mucous membrane in the cartilaginous portion is more fully developed than in the bony portion, and contains many acinose glands. In the bony portion there are but few of these glands, and the membrane is very thin, smooth, and closely adherent.

The tensor palati and levator palati are the muscles that open the tube during the act of swallowing, and thus favor the passage of air toward the tympanic cavity. The tensor palati has its origin from (1) the under surface of the sphenoid, from (2) the extremity of the cartilaginous hook, and also from (3) the mem-

branous wall of the cartilaginous portion of the tube. Passing downward around the hamulus pterygoideus, it spreads out in the fibrous prolongation of the hard palate. The tendon being quite firmly attached to the hamulus pterygoideus, it acts more fully on the Eustachian tube than on the soft palate when the muscle contracts. Its action on the tube is to separate the outer or membranous wall, together with the cartilaginous hook, from the inner wall.

The levator palati arises from the surface of the petrous bone adjoining the carotid canal. In its course parallel to the Eustachian tube it is joined by some connective-tissue fibres to the membranous portion which forms the base of the tube, and also to the cartilaginous plate, and is inserted below the pharyngeal opening of the tube into the soft palate. The action of this muscle seems to increase the transverse diameter of the cartilaginous canal by pulling the lower wall of the tube upward and backward.

The arterial supply is derived from (1) the pharyngeal, a branch of the external carotid artery; (2) the middle meningeal, from the internal maxillary artery, and (3) some small branches from the internal carotid artery. The veins are numerous, and form a plexus which communicates with the cavernous sinus and with the plexus near the temporomaxillary articulation, and empty into the facial or internal jugular veins.

The tensor palati muscle derives its nervous supply from the otic ganglion and from the internal pterygoid nerve, which is a branch of the inferior maxillary nerve. The levator palati is supplied by a branch of the pneumogastric and by the facial through the Vidian and petrosal nerves.

3. The Mastoid Process.

The mastoid process under normal conditions communicates with the posterior part of the middle ear by means of the antrum. In the light of anatomical investigations we know that no two mastoid processes are exactly alike. According to Zuckerkandl, who made an examination of one hundred mastoid processes in the fresh state and of one hundred and fifty macerated ones, there were absolutely no air-cavities in 20 per cent. of the specimens. Of the entire number, 38.6 per cent. were perfectly pneumatic mastoid processes without diploëtic spaces. He found also that in some persons only the upper half of the process was pneumatic, while the lower half was diploëtic. It will thus be seen that there are three different kinds of mastoid processes: (*a*) the pneumatic, (*b*) the diploëtic, and (*c*) a mixed variety or a combination of the two. In the pneumatic variety (see Plate I.) very frequently numerous cellular spaces are found extending into the temporal bone in different directions—viz., around the lateral sinus, down to the apex, backward to the occipital suture and upward toward the root of the zygoma—a point to be remembered in operations on the mastoid, for it is most important in such cases to remove thoroughly all diseased tissue found in these cells. In an infant at birth there is usually found but one cell, the *antrum*, and the mastoid process is but a small tuberosity and undeveloped. From birth the mastoid process gradually develops by extending downward, until at five years of age it is very similar to that of an adult, as regards the position of the antrum and arrangement of the cells, except that it is smaller and the bony substance is not as firm and dense. In a pneumatic mastoid process, beside the *antrum* already mentioned, there are

PLATE I.



Section showing Middle Ear and Mastoid Cells, Pneumatic Variety.

1. Membrana tympani.
2. Aquæductus Fallopii.
3. Mastoid antrum.
4. Tensor tympani muscle.
5. Malleus and incus.
6. Mastoid cells.
7. Promontory.
8. Foramen rotundum.
9. Foot-plate of stapes in foramen ovale.
10. Semicircular canal.
11. Canal of carotid artery.
12. Bony wall of Eustachian tube.

usually a number of cells that communicate with one another as well as with the antrum. These contain air, and are lined with a continuation of the mucous membrane of the tympanic cavity, and the epithelium is of the non-ciliated variety. Those cells that are not pneumatic usually contain a fatty substance. The upper wall of the antrum is but a thin lamina of bone, which separates it from the dura mater. The lateral sinus is in close proximity to the mastoid cells, and rests against the inner wall of the process. The facial nerve may become involved in inflammation of the mastoid cells, or may be injured during operations, as its course is downward through the cells after passing along the upper portion of the tympanic cavity. A point which serves as an anatomical landmark in operations on the mastoid is the *spina supra meatum*. This is a prominence situated just a little above and behind the external meatus at the junction of the mastoid process with the posterior wall of the external auditory canal. Occasionally there are found cells in the occipital bone that communicate with the mastoid cells. Pus frequently makes its way down to the lowest extremity or tip of the process, and, as the inner wall is very thin, the bone is occasionally perforated, and an abscess may form underneath the muscles and deep fascia in the neck. The lateral sinus occasionally runs out of its usual course, so that great care should always be exercised in operations on the mastoid.

As we have seen that the internal structure of the mastoid process is subject to great variations in different individuals, so the outer surface of the process may have a very thick or a thin wall, or may be rounded or flattened. It is believed very generally that in a normal mastoid process in which the external wall is thick

the bone will be found thin at the mastoid tip and digastric fossa; and, further, that when the mastoid protuberance is small and with a well-defined digastric fossa the lateral sinus is apt to enroach upon the mastoid antrum. As the two mastoid processes in an individual are usually symmetrical, valuable information can often be obtained by comparing the normal with the one that is inflamed.

THE INTERNAL EAR.

The inner ear, or sound-perceiving apparatus, consists of a bony and a membranous labyrinth, together with the central origin and trunk of the auditory nerve.

The Labyrinth.

There is a bony labyrinth contained within the petrous portion of the temporal bone, and a membranous labyrinth within the bony one. The *bony labyrinth* is divided into (1) the vestibule, (2), the semicircular canals, and (3) the cochlea.

1. **The Vestibule** lies on the inner side of the tympanum in front of the semicircular canals and behind the cochlea, and is the central cavity by which the different parts of the bony labyrinth are in communication with one another. It is somewhat oval or elliptical in shape. On its outer wall is situated the *fenestra ovalis*, which is closed by the foot-plate of the stapes (Plate I.), and the latter is held in position by the annular ligament. There is a small depression, called the *fovea hemispherica*, at the anterior part of the inner wall, which contains several minute perforations (macula cribrosa) for the passage of the filaments of the vestibular nerve. The *crista vestibuli* is a ridge lying behind

this depression. At the posterior part of the inner wall is found the opening of the *aqueductus vestibuli* (Fig. 12), through which a small vein passes, and it is claimed by some authorities that it contains a tubular prolongation of the lining membrane of the vestibule, and that it terminates in a cul-de-sac between the layers of the dura mater.

FIG. 12.



The osseous labyrinth laid open. (Enlarged.) (GRAY.)

There is an opening in the vestibule which communicates with the scala vestibuli of the cochlea, called the apertura scalæ vestibuli cochleæ.

2. The Semicircular Canals, three in number, open into the posterior part of the vestibule by five orifices. They are of unequal length, flattened from side to side, and describe almost a circle in each case. The dilated end of each canal is called the *ampulla.* One of the

canals is vertical in direction, and is called (1) *the superior semicircular canal;* another, directed backward, is also vertical in direction, (2) *the posterior semicircular canal;* the third, the shortest of the three, runs in a direction outward and backward, (3) *the external or horizontal semicircular canal.* Each end that is dilated, or has an ampulla, as well as the smaller end of the external semicircular canal, has a separate opening into the vestibule. The smaller ends of the posterior and superior canals, which join before entering the vestibule, have a common opening.

FIG. 13.



The cochlea laid open. (Enlarged.) (GRAY.)

3. The Cochlea. The cochlea, resembling on a cross-section the ordinary garden-snail shell, consists of a tube which takes two and a half turns about its central axis or modiolus (see Figs. 12 and 13), In a macerated preparation it will be seen that there is an opening or passage between the cochlea and vestibule, and also a communication between the cochlea and tympanum through the *fenestra rotunda.* The cochlea, conical in shape, is placed almost horizontally in front of the vestibule, with its base directed toward the internal

PLATE II.



After Photomicrograph of a Section through Cochlea of Rabbit. Collection of Dr. J. E. Weeks. × 26 diam.

1. Scala vestibuli.
2. Scala media.
3. Scala tympani.
4. Membrane of Reissner.
5. Organ of Corti.
6. Cochlear nerve lying within the modiolus.
7. Ganglia spirale.

auditory meatus and its apex or cupola toward the tympanic cavity. There are several minute openings in the base for the passage of the cochlear branch of the auditory nerve. The tube or canal of the cochlea is partially subdivided by the *bony lamina spiralis*, which commences at about the fenestra rotunda and ends at the cupola in the *hamulus*. The *membranous lamina spiralis* is inserted into this bony lamina spiralis, so that the canal is thereby subdivided into two passages, the superior called the *scala vestibuli* and the inferior the *scala tympani*. A second membrane cuts off a portion of the scala vestibuli and forms the *scala media* or *ductus cochleæ* (see Plate II.), but the latter properly belongs to the membranous labyrinth. There is a communication at the apex of the cochlea between the scalæ tympani and vestibuli through the *helicotrema of Breschetius*. There is quite a large canal passing through the centre of the modiolus named the *canalis centralis modioli*, extending from the base to the cupola. It transmits a small nerve and artery. There is an orifice, the *aqueductus cochleæ*, which leads to a canal containing a small vein, and which opens on the basilar surface of the temporal bone. A swelling of the cochlear nerve, called the *ganglion spirale* (see Plate II.), lies in a small canal, the *canalis spiralis modioli*. This canal winds around the modiolus and lies at a point where the bony lamina is attached to the modiolus. Ganglion-cells are found in the ganglion spirale, from which nerves pass through the bony lamina to the organ of Corti. The lamina spiralis terminates in a curved extremity, the *sulcus spiralis internus*, the upper lip of which is called the *labium vestibulare*, the lower lip the *labium tympanicum*. Between the labium vestibulare and the origin of the membrane of Reissner is the *crista*

spiralis. A delicate membrane, the *membrana basilaris* (see Plate III.), completing the scala tympani, extends from the labium tympanicum to the *ligamentum spirale,* which appears as an expansion of the membrana basilaris attached to the outer bony wall of the cochlea. The *stria vascularis* is a thin layer of cells resting on that part of the ligamentum spirale between the attachment of the membrane of Reissner and the *prominentia spiralis.* Between the prominentia spiralis and the attachment of the membrana basilaris is the *sulcus spiralis externus,* and this point of attachment of the membrana basilaris is termed the *crista basilaris.* The *membrane of Reissner* is stretched across from the vestibular surface of the periosteum of the bony lamina to the ligamentum spirale, and forms the superior boundary of the scala media, the floor being the membrana basilaris. A very thin membrane, described by Corti and called the *membrane of Corti,* or the *membrana tectoria,* extends across from the attachment of Reissner's membrane and the labium vestibulare to the outer row of hair-cells. The *organ of Corti* consists of numerous rods or cells resting on the membrana

---

DESCRIPTION OF PLATE III.

After photomicrograph of a section of the cochlea of a rabbit, showing relations of the organ of Corti. 160 diameters. (Collection of Dr. J. E. Weeks.)

S. V. Scala vestibuli. *S. M.* Scala media. *S. T.* Scala tympani. *S. S. E.* Sulcus spiralis externus. *S. S. I.* Sulcus spiralis internus, with its epithelium.

1. Membrane of Reissner. *2.* Crista spiralis. *3.* Membrana tectoria, or membrane of Corti. *4.* Labium vestibulare. *5.* Inner hair-cells. *6.* Labium tympanicum. *7.* Zona arcuata, showing its floor-cells and two nerve-fibres passing to outer hair-cells. *8.* Inner rod of Corti. *9.* Outer rod of Corti. *10.* Lamina reticularis or membrane of Kölliker, covering the outer hair-cells. *11.* Supporting cells of Hensen. *12.* Cells of Claudius. *13.* Deiter's cells. *14.* Membrana basilaris. *15.* Crista basilaris. *16.* Prominentia spiralis. *17.* Stria vascularis. *18.* Ligamentum spirale. *19.* Upper plate of bony lamina spiralis. *20.* Lower plate of bony lamina spiralis.


S.V.
S.M.
S.T.
SS1
SSE
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

basilaris and covered by the membrana tectoria. The two central cells, rod-like in character, are called the *inner* and *outer rods of Corti*, and rest on the membrana basilaris, somewhat separated from one another. This intervening space is called the *zona arcuata* (see Plate III.). The very small canal thus formed by these arches and by the membrana basilaris, passes in a spiral direction throughout the whole length of the cochlea. Within the zona arcuata will be found two rows of cells, one in each angle formed by the rods of Corti and the membrana basilaris. They are called the *floor cells*. Two nerve-fibres will be observed in Plate III., crossing the zona arcuata to the outer hair-cells. The *outer hair-cells* are variously described as from three to five rows in number. They lie to the outer side of the outer rods of Corti, their broad ends directed upward. Between these will be found the cells of Deiter, the broad ends of which are directed downward. There is a single row of *inner hair-cells*, smaller than the outer, lying to the inner side of the inner rods of Corti. They have a layer of supporting cells on the inner side continuous with the epithelium lining the sulcus spiralis internus. Lying in close contact with the upper ends of the rods of Corti and the outer hair-cells will be found a thin reticulated membrane with openings for passage of the cilia of the outer cells. It is known as the *lamina reticularis*, or membrane of Kölliker, and seems to serve the purpose of binding together the cells over which it lies.

A mass of cells are present on the outer side of the organ of Corti which appear to be for the support of that organ. There are more of them, and they extend higher into the scala media in the rabbit (from which Plate III. is taken) than in the human subject.

The cells forming the upper portion are known as the supporting cells of Hensen, those forming the lower outer portion, with the large number of small nuclei, resting on the basilar membrane and extending outward to the crista basilaris, are called the cells of Claudius.

The *membranous labyrinth* consists of (1) the membranous canals, (2) the utricle, and (3) the saccule. The *membranous semicircular canals* are similar in number, arrangement, and general form to the bony canals, except that they are only about one-third the size of the latter. They open by five orifices into the utricle, one opening being common to two canals. The ampulla is much thicker than the other portions of the canal, and almost fills the bony encasement. The membranous canal consists of three layers, the inner being formed of polygonal nucleated epithelial cells which secrete the *endolymph*, a fluid which is contained within the membranous tubes. A fluid quite similar to the endolymph, called the *perilymph*, surrounds the membranous canals and is contained within the osseous canals. The ampullæ receive nerve-filaments from the vestibular nerve.

The *utricle*, oblong in shape, lies in contact with the fovea hemielliptica in the superior and posterior part of the vestibule. There is a yellow thickening on the anterior and lateral walls of the utricle, called the *macula acustica*. A clear, transparent semifluid substance covers the neuro-epithelium of the macula acustica, and contains crystals of carbonate of lime called the *otoliths*.

The *saccule*, globular in shape, is situated in the fovea hemispherica, and apparently its cavity is distinct from that of the utricle. It has a macula acustica, and otoliths similar to those in the utricle. There is a small

canal, the *canalis reuniens* (see Fig. 14), connecting the saccule with the ductus cochlearis of the membranous labyrinth. A number of stiff cilia project from the surface of the macula in both saccule and utricle, and are connected with the nerve-filaments of the auditory nerve.

The arteries supplying the labyrinth are: the (1) internal auditory, which is a branch of the basilar; (2) some branches from the occipital; (3) the stylomastoid,

FIG. 14.



from the posterior auricular. The *internal auditory artery* subdivides at the bottom of the internal auditory meatus into the *cochlear* and *vestibular* branches. The cochlear branch gives off numerous small vessels which pass through the canals in the modiolus and enter the substance of the lamina spiralis in the form of a capillary network. The vestibular branch forms a capillary network which is distributed to the membranous labyrinth.

The veins from the semicircular canals, the vestibule, and the cochlea unite with the superior petrosal sinus.

The *auditory nerve*, or nerve of hearing, also called the portio mollis, is the eighth nerve. Its superficial origin is at the lower border of the pons from a groove between the olivary and restiform bodies. There are two deep origins, both from the fourth ventricle. The nerve, after winding around the restiform body, passes, together with the facial nerve, across the posterior border of the middle peduncle of the cerebellum and enters the internal auditory meatus with the facial nerve. At the bottom of the meatus the nerve divides into the *cochlear* and *vestibular* branches.

The vestibular branch subdivides into the superior, middle, and inferior branches, which are distributed to the cristæ acusticæ of the semicircular canals and the maculæ acusticæ of the utricle and saccule. The cochlear branch divides into numerous twigs, and, after entering the base of the modiolus, passes between the plates of the osseous lamina spiralis, at which point the nerves form a plexus, the "ganglion spirale." From this plexus nerve-filaments are distributed to the sulcus spiralis and organ of Corti.

PHYSIOLOGY.

The function of the auricle is to collect the sound-waves and direct them to the membrana tympani. The concha is especially important for this purpose, as well as the cavity just beneath the tragus. After the sound-waves have reached the drumhead through the external auditory canal, they are transmitted by the chain of ossicles to the fluid of the labyrinth, the ossicles probably vibrating as whole bodies. The sound-waves then pass through the fluid in the cochlea and reach the

PLATE IV.

FIG. 1.



Section through human cochlea. (Dixon.)

FIG. 2.

Section through a portion of modiolus and lamina spiralis, showing (*A*) ganglion spirale, and (*B*) nerve fibres passing outward to be distributed to the organ of Corti. (Dixon.)

organ of Corti. From the nerve-filaments in the cochlea the sensations called sound are conveyed through the auditory nerve to the brain centre.

According to experiments made by Helmholz, it seems that the power of resonance of curved membranes is much greater than that of flatly stretched ones, a point of great importance in considering the concavity of the drumhead. Experiments have also shown that the vibrations of the malleus are twice as great as those of the incus and four times as great as those of the stapes. It is probable that by means of the cochlea the pitch of the sound is appreciated, and, further, that different notes are perceived by certain portions of the spiral scale. As to the function of the fenestra rotunda, it is probable that, owing to its elasticity, when the foot-plate of the stapes is forced in and the labyrinthine pressure is increased, such pressure is compensated for by the bulging outward of this membrane. When this delicate membrane becomes thickened through pathological changes, the patient is likely to complain of noises in the ear and vertigo, if the pressure within the labyrinth is suddenly increased.

It was formerly believed that the function of the semicircular canals had to do with the perception of the general direction of sounds, but it is more probable, and a fact generally accepted, that the sense of motion and of equilibrium is the function of the semicircular canals and the utricle.

By means of the Eustachian tube, air is allowed to enter the middle ear and thus keep the atmospheric conditions equalized on both sides of the membrana tympani. The Eustachian tube is not constantly open, as its permeability varies in different individuals. In some persons a current of air will pass from the pharynx to

the middle ear during ordinary respiration, while in others the act of swallowing or a powerful expiratory act, with nostrils open or closed, will be necessary to cause the air to enter the Eustachian tube. When constriction of the tube takes place, so that the air does not enter the tympanum except with difficulty, the result is that the drumhead and ossicles are forced inward, and congestion of the middle ear occurs, which condition, if not relieved, is likely to be followed by deafness.

CHAPTER II.

METHODS OF EXAMIMATION OF THE EAR.

When a patient presents himself for advice and treatment for some aural disease, a most complete history of the case should first be obtained. He should be questioned as to whether he has any hereditary tendency to deafness; then inquiry should be made as to his general health, particularly as to any predisposition to catarrhal disease. The patient should then be questioned as to the exact symptoms for which he comes for advice, and the surgeon should make careful note of the pain, if present, the tinnitus, the discharge, or vertigo.

After having obtained (1) a full and complete history of the case in all its details, the surgeon should then (2) inspect with the eye the auricle and surrounding parts, looking for any inflammation of the pinna or external meatus, or any redness over the mastoid process. By palpation some information can be obtained, especially if there is any tenderness on pressure over the mastoid process. It is always well to compare both auricles and both mastoid processes of a patient, for it must be remembered that in some persons, generally those of an hysterical temperament, pain is occasionally complained of when firm pressure is made over the apex of the mastoid, so that it is always well to compare the two sides. Redness and tenderness over the mastoid are symptoms suggestive of mastoid inflammation or

furuncular inflammation of the *external meatus*. After having obtained all information possible by inspection and palpation, the surgeon should then (3) make a most thorough examination, by means of the head-mirror and speculum, of the external auditory canal, the membrana tympani, and, if the latter is perforated, of the tympanic cavity.

A very good head-mirror is the one recommended by Politzer, and shown in Fig. 15. It is a concave mirror,

FIG. 15.



Indirect illumination of the ear.

with an aperture in the middle which allows the surgeon to look through the opening, if he desires to do so, and it should be fastened to the forehead-band or metal spring by means of a ball-and-socket joint. The mirror is two and three-quarters inches in diameter and has a focal distance of about six inches. In order to obtain the best view of the drumhead, the mirror should be pulled down over the eye, so that the surgeon can look through the opening; and to make a thorough examina-

tion of the external meatus and membrana tympani, it is most important to have good illumination, and for this purpose daylight from a northern exposure is the most satisfactory, since by this means the natural color of the drumhead will be preserved. Direct sunlight is usually too dazzling.

When good natural light cannot be obtained, artificial light should be substituted, such as that from an Argand burner, with or without a Mackenzie's bull's-eye lens

FIG. 16.



Mackenzie's bull's-eye lens and reflector.

and reflector. Some surgeons prefer the electric light, but for ordinary examination gas is still the most practical and desirable.

An excellent light is that obtained from a candle fitted in a holder and supplied with a lens. This can be carried by the surgeon in his satchel, and is particularly adapted for bedside examinations.

Illumination of the ear may be obtained by indirect illumination, as when an Argand burner or natural daylight is used, or by direct illumination, the latter method consisting of an electric lamp attached to the head-band.

FIG. 17.

Direct illumination of the ear.



Electric light and portable battery.

An excellent electric lamp with head-mirror is the one devised by Leiter, of Vienna. Another lamp is represented in Fig. 17.

In order to make use of the electric current in a city house, a rheostat has lately been devised, which regulates the current so that the surgeon is enabled not only to obtain a good illumination of the ear, but also to make use of a cautery and bone-drills at the same time (Fig. 18). For operations on the middle ear, and for

FIG. 18.



Electric motor with rheostat.

operations performed at night, on account of the danger of conflagration, the electric light is indispensable, and should be used whenever possible. For this purpose a storage battery, as represented in Fig. 17, may conveniently be carried by the surgeon. A very small storage battery, to which the Leiter lamp may be attached, has especially been designed to be carried in the physician's satchel. This will be found most convenient for bedside

examinations and for short operations. The objection to the photophore for general use is that it becomes very hot when employed for any length of time, and consequently very disagreeable to the surgeon.

Brunton's otoscope (Fig. 19) is but little used in this country, but is more popular in England. Its advantages, as claimed by the inventor, are its simple construction and easy application; it may be used with either sunlight or artificial light, and with or without the magnifying glass. The rays of light enter the funnel. An electric light has been adapted to this otoscope.

Fig. 19.



Brunton's otoscope.

Brunton's otoscope is obviously useless except for purposes of examination.

When making an examination with a head-mirror, it is of the greatest importance to place the patient in a good position. The ear to be examined should be away from the light. If gaslight is used, the Argand burner should be on a level with the ear, and the rays of light, striking the head-mirror at an angle of 45 degrees, should be reflected into the external auditory canal.

Speculum. The speculum that I find most satisfac-

tory is that known as Gruber's. It is constructed of silver, the narrow part being highly polished inside, while the dilated portion inside is blackened. There are four sizes of these specula. The end to be introduced is conical in shape (Fig. 20), which allows the

Fig. 20.

Gruber's speculum.

speculum to fit more accurately the exterual meatus, so that the surgeon is enabled to obtain a better view of both the external auditory canal and the drumhead.

Politzer's specula (Fig. 21) of hard rubber are also excellent ones, and will be found especially useful when

Fig. 21.

Politzer's speculum.

caustic applications are to be made to the canal or drumhead. The Gruber speculum is also made of hard rubber. Another speculum is that devised by Boucheron, of Paris (Fig. 22). I have modified the latter by having the extremity of the narrow end made oval instead of round. Before introducing the speculum it should be warmed, and particular care exercised that it is not

pushed in too far, as pressure on the bony canal is quite painful. After the speculum has been warmed, it should be held by the thumb and index finger of the left hand, and with a gentle rotary motion carefully introduced into the meatus, while the auricle, held between the index finger and middle finger of the same hand, should be drawn outward, upward, and backward. Occasionally, long hairs obstruct the view. These should be pushed aside, while any cerumen, scales, purulent or cheesy matter should be removed by syringing or by

FIG. 22.



Boucheron's speculum.

means of curettes or forceps, as the case may be. Mistakes are frequently made by beginners in not pushing the speculum beyond the hairs at the meatus, which at times are covered with cerumen, so that a diagnosis of inspissated cerumen is occasionally made. Again, if the long axis of the speculum is not directed toward the drumhead, but against the walls of the auditory canal, the student will make a mistake in diagnosis, as the walls are frequently smooth, glistening, and pinkish in color, resembling at times an inflamed drumhead.

OPERATING EAR SPECULA. For operative purposes specula have been devised with a slit-like opening in the

narrow part of the speculum, to allow greater play for the use of instruments. A short speculum is especially adapted for such purposes. Blake's operating otoscope "consists of a hard-rubber speculum (Politzer), fitted with a metallic rim, to which are attached a revolving prism and an arm bearing at its outer end a lens of about an inch focus; this arm is movable, but sufficiently firm to remain fixed at any angle at which it is placed."

FIG. 23.



Author's curette forceps.

All drops or solutions used in the ear should be warmed first, for otherwise they are liable to cause vertigo and inflammation. The syringe is the safest instrument that can be placed in the hands of the beginner, for the purpose of removing cerumen, purulent discharge, or scales, while forceps and curettes should never be used by the inexperienced, nor at any times,

Fig. 24.

Foreign-body forceps.

Fig. 25.



Aluminium cotton-carrier.

except under excellent illumination. The curette forceps (Fig. 23) I have used for many years, and find them especially adapted to removing hardened cerumen

when syringing alone has failed. Another pair of forceps that will be found of service are those represented in Fig. 24.

In addition to the forceps, it is necessary that the surgeon should be well supplied with cotton-carriers (Fig. 25). Those of aluminium will be found the most serviceable, on account of their lightness and flexibility. The surgeon will find it necessary to have several sizes of blunt curettes, for the purpose of removing scales, cerumen, etc., from the auditory canal.

APPEARANCES OF THE NORMAL EXTERNAL AUDITORY CANAL AND MEMBRANA TYMPANI.

After introducing the speculum and pushing aside any hairs in the meatus, one should observe the appearance of the auditory canal. The osseous part of the normal canal is usually of a delicate pink color and somewhat smooth and glistening. A reflex cough is frequently caused by introducing the speculum, due to irritation of the auricular branch of the pneumogastric.

The normal membrana tympani, when examined by daylight, is of a pearly-gray color with a slightly bluish tint, while gaslight gives it a reddish-yellow color. Other points to be observed are the long handle of the malleus, running downward and backward from the short process to the umbo or central concavity of the membrana. The cone or pyramid of light, as shown in Figs. 26 and 27, extends from the umbo almost to the lower periphery of the drumhead, with the apex at the anterior part of the umbo. It is supposed to be due partly to the concavity of the drumhead and to the angle that the membrana tympani forms with the

axis of the canal, and partly to the glistening or reflecting surface of the membrana.

From the short process, extending forward and backward, are two folds called the *anterior* and *posterior folds.* The portion of the drumhead between these folds above the short process is called the *membrana flaccida,* the portion below being the *membrana vibrans.*

In looking at a drumhead, one must note particularly (1) its color, whether normal, congested, opaque, or atrophic, and whether it contains opacities or chalky

Fig. 26.



Normal drumhead (right ear).

Fig. 27.



Normal drumhead (left ear).

(Politzer.)

a. Posterior fold. *b.* Short process. *c.* Anterior fold. *d.* Cone of light *e.* Long handle of incus. *f.* Umbo.

deposits; (2) the position of the drumhead, whether bulging, owing to a collection of fluid, or whether retracted. In the latter case the handle of the malleus, being retracted, will appear foreshortened; the short process will become more prominent, as well as the anterior and posterior folds, and the cone of light will be distorted or displaced. To decide as to the mobility of the drumhead, and whether bound down by adhesions, in addition to the Valsalvian method of inflation (to be described later), more or less can be determined by touching the different parts with a probe. For this

purpose, however, an instrument has been devised—viz., Siegle's pneumatic speculum or otoscope—consisting of a hard-rubber cylinder closed at one end by an obliquely placed thin plate of glass. There is a small opening on the side, connected by rubber tubing with a

FIG. 28.



Siegle's otoscope, with masseur of Delstanche attached.

Politzer bag, or, better still, with the masseur of Delstanche (Fig. 28). At the other extremity of the cylinder a speculum is attached, and it is well to cover the narrow end of the speculum with a small piece of rubber tubing, to make it fit the canal accurately and thus render the instrument air-tight. By manipulating the

piston with the thumb the mobility of the membrane can be determined. I have modified this instrument by substituting the Gruber metallic speculum for the hard-rubber ones, which are, as a rule, attached to this instrument (Fig. 28).

The use of the Siegle otoscope, for the purpose of exhausting the air in the external auditory canal, is indicated in cases of retraction of the drumhead due to catarrhal disease; to break up adhesions between the drumhead and walls of the middle ear; in certain cases of tinnitus, which frequently are much benefited after its use, and in cases of dizziness due to increased labyrinthine pressure from middle-ear disease; also in certain cases of perforation of the drumhead in which it is important to remove the secretion. The "rarefacteur," or masseur, of Delstanche is particularly useful in cases of adhesion between the membrana and walls of the tympanum, as rarefying and condensing the air in the external meatus effect what Delstanche calls "massage of the articulation of the ossicles." All instruments, for either rarefying or condensing the air in this manner should be used with great caution, and only when the surgeon can obtain a good view of the drumhead during such manipulation.

APPEARANCE OF THE PHARYNX, NASOPHARYNX, AND NASAL PASSAGES.

The surgeon should not consider the examination of the ear complete until he has inspected carefully the pharynx, the nasopharynx, and the nasal passages. For this purpose, in order to direct the light, a larger head-mirror is advisable.

First of all, the tongue should be depressed by means

of a spatula and the patient directed to make an inspi-

Fig. 29.



Colin's nasal speculum.

Fig. 30.

Mirror for posterior rhinoscopy.

ratory effort or to pronounce the word "ah," which latter procedure will raise the soft palate, and thus present an excellent view of the pharynx. Careful attention should be paid as to whether the pharynx is atrophic, hypertrophic, or granular, whether the tonsils are enlarged or not, and whether there are any ulcers.

As to the nasal passages, one should direct the patient to close one side of the nose and to force air in and out of the other passage, and to do the same on the other side, in order to determine the permeability of these passages. Afterward, inspection by means of a nasal speculum (Fig. 29) should be made, to ascertain the presence of polypi or spurs on the septum, or to detect malformations, disease of the bone, or hypertrophic or atrophic nasal catarrh, or ozæna.

After the pharynx and the nasal passages have been explored, the attention of the surgeon should be directed to the condition of the nasopharynx. A mirror (Fig. 30) should be warmed and carefully passed

behind the soft palate, and the nasopharynx should be so illuminated by means of the head-mirror that the surgeon can decide if there are adenoid growths present, or polypi or hypertrophy of the mucous membrane, or any other abnormal change in the tissues. In children it is particularly difficult to employ posterior rhinoscopy, and in such the index finger may be easily passed up behind the soft palate, and the presence or absence of adenoid growths may be ascertained. In nervous and excitable children this method should never be employed, as I have seen little patients in whom this method had been attempted rendered so nervous that it was extremely difficult for the aural surgeon to regain their confidence. Whenever possible, it is much more satisfactory and considerate to give the patient a whiff of chloroform.

EXAMINATION OF THE EUSTACHIAN TUBE AND MIDDLE EAR.

In order to ascertain the condition of the Eustachian tube, it is necessary to try one of three methods of inflation, viz., (*a*) Valsalvian method, (*b*) Politzerization, (*c*) Catheterization. A diagnostic tube (Fig. 31), consisting of rubber, from two and one-half to three feet in length, and provided with ivory tips, should always be used. One end is placed in the surgeon's ear and the other in that of the patient. When inflating the ear by either of the three methods, certain information can be obtained as to the condition of the Eustachian tube. In a normal case, when the middle ear is inflated a dull sound or click will be heard. The presence of fluid in the tympanic cavity will give rise to a bubbling sound, while a perforation in the membrana tympani

will be indicated by a blowing or hissing sound, according to the size of the perforation and the condition of the tube.

FIG. 31.



Diagnostic tube.

(*a*) VALSALVIAN METHOD. By this method the patient holds the nose closed with the thumb and index finger, and, the mouth also being closed, the condensed air in the nasopharynx is forced through the Eustachian tube into the tympanum when the patient makes a forced expiration. This method is useful only in cases in which it is desirable to make a diagnosis of the permeability of the Eustachian tube and to determine the mobility of the membrana. The drumhead should be inspected under good illumination whenever this method is employed; but the latter should never be recommended to patients, as it tends to produce congestion of the head, and when frequently practised, and for a long period of time, it is likely to stretch the ligaments of the ossicles, and thus cause a relaxed condition of the parts. This method is of limited value, and should never be used where there is much obstruction to the entrance of air in the tube.

(*b*) POLITZER'S METHOD. The method of Politzer is described best in his own words, as follows: "The

patient, being seated in a chair, takes a little water into his mouth—to facilitate swallowing—which he is required to swallow when told. The surgeon, standing on the patient's right, or in front, introduces the nozzle of the Politzer bag 1 cm. ($\frac{3}{8}$ inch) into the nasal orifice of the corresponding side at its posterior angle (Fig. 32),

Fig. 32.

Politzer's method.

and then compresses with the left thumb and forefinger the alæ of the nose closely round the instrument. The patient is next told to perform an act of swallowing, and at the same moment the surgeon expels the air from the inflating-bag with his right hand. By the condensation of air produced in the nasopharynx in this manner the air is forced into both middle ears, the closure effected by the soft palate is forced open, and its vibrations give rise to a dull, gurgling noise, which

frequently, if not always, may be taken as an indication that the air has entered into the middle ear." Politzer has partially modified this method by allowing the patient to inspire through the nearly closed lips, or through a piece of rubber tubing held between the lips while the air is being forced into the nasopharynx. By this modification of the original method the nasopharynx is not only closed, but at the same time the Eustachian tube is made more open by the soft palate being pulled backward. By inspiration through the mouth air enters the middle ear more readily when the catheter is used than by simple catheterization.

In inflating the middle ear, in cases where there is fluid, it is well to have the patient's head inclined forward and sideways, in order to allow the secretion to gravitate toward the nasopharynx.

In cases of perforative inflammation of the middle ear it seems to be proved that, by Politzer inflation, the secretion in the middle ear is forced through the opening in the drumhead and not into the mastoid cells. In cases of hyperæmia of the middle ear, with deafness resulting from inflammatory action, inflation of the middle ear in many cases will produce absorption of the exudation, and will also have a favorable effect on the circulation in the tympanic cavity.

The fact that a hissing sound is not always heard when the Valsalvian or Politzer method has been tried does not indicate that there is not a perforation of the drumhead, for there may be so much swelling or constriction of the tube that catheterization will be necessary to force air through the perforation.

The Politzer bag (Fig. 33) is pyriform in shape, and should hold about eight ounces. A rubber tube, six

inches in length, should be attached to the bag, and the nosepiece should be of glass, large enough to completely fill the nasal orifice, glass being easily cleansed and rendered aseptic.

It is best to have a small opening on the side of the bag, which should be covered over by the thumb when compressed. The thumb should be removed immediately from the opening after compressing the bag, so as to allow the air to enter the bag through this open-

FIG. 33.



Politzer's bag, with glass inhaler attached.

ing. Otherwise the air and possibly some secretion may be drawn through the glass attachment.

Instead of having the patient swallow water, another method, recommended by Gruber, consists in the pronunciation of the word "hock" by the patient at the moment that the surgeon forces air through the nose. With young children both of these methods are unnecessary, for if the mouth of the child is closed, it will

simply be necessary to use the Politzer bag. If the child cries, it will facilitate matters also.

(*c*) CATHETERIZATION. Both hard-rubber and silver catheters are used. I prefer, however, the latter (Fig. 34), of which there are four different sizes; it

FIG. 34.

Blake's Eustachian catheter.

is most important that each should have a probe-pointed extremity for introduction into the nose. The catheter should have a double curve, as recommended by Blake. The catheters formerly in use caused considerable pain and irritation when introduced. By using a catheter with a probe point very little discomfort is caused to the patient if the surgeon is skilful, and there is little if any danger of lacerating the parts and thereby causing emphysema of the connective tissue of the pharynx and neighboring parts.

The use of a 4 per cent. solution of cocaine sprayed into the nasal passages is advisable in some cases before passing the catheter, provided the patient is not unpleasantly affected by this remedy. The surgeon, however, should never resort to cocaine if he can possibly do without it, for fear the patient may ultimately contract the cocaine habit. The author has besides seen patients whose general health was seriously impaired by this practice on the part of the otologist, viz., of spraying the nasal passages with a cocaine solution before introducing the catheter. There are two methods generally employed in introducing a catheter:

First Method. The patient should be seated opposite the surgeon, and the former's head steadied by an assistant or allowed to rest against the back of a chair. The surgeon, either standing or sitting opposite the

FIG. 35.



Vertical section of the nasopharynx, with the catheter introduced into the Eustachian tube. (POLITZER.)

A. Inferior turbinated bone. B. Middle turbinated bone. C. Superior turbinated bone. D. Hard palate. E. Velum palati. F. Posterior pharyngeal wall. G. Rosenmüller's fossa. H. Posterior lip of the orifice of the Eustachian tube.

patient, then places the fingers of the left hand on the patient's forehead, gently pushing upward the extremity of the nose with the thumb of the same hand. With

the catheter held loosely between the thumb and forefinger of the right hand, the probe-pointed end is introduced very carefully, after being warmed and smeared with olive oil or vaseline. The end of the catheter should be pushed along the lower wall of the nasal passage until it touches the posterior wall of the pharynx, when it is turned obliquely outward, but not sufficiently to be in a horizontal position, and gently forced into Rosenmüller's fossa. By withdrawing the catheter and at the same time raising the probe-pointed extremity, the latter will be felt to glide over the posterior lip of the Eustachian tube. The point is then turned still further outward and upward, "so that the metal ring fastened to its posterior extremity is pointed toward the outer canthus of the eye of the same side" (Fig. 35). "The direction of the beak corresponds, as a rule, with the axis of the Eustachian tube" (Politzer). The catheter may be held in position by the surgeon, or, for this purpose, various clamps have been devised.

Second Method. By this method the posterior margin of the nasal septum serves as a guide. The catheter is introduced in the same way as in the first method, until the point reaches the posterior pharyngeal wall. Instead of turning the point outward, it should be turned inward, so that the extremity points toward the other Eustachian orifice. It is brought to a horizontal position, as determined by the metallic ring on the other end of the catheter. The catheter is then withdrawn until the curved portion reaches the posterior border of the nasal septum. The point of the catheter is now made to describe a complete semicircle by moving its point downward along the velum palati and then outward. It should thus enter the pharyngeal orifice, and be inserted afterward in the tube in the same manner

as by the first method. Any action of the muscles of these parts that pushes the point of the catheter backward or throws it out of position will interfere with the success of this method. It is occasionally impossible to effect an entrance into the tube by either of these methods, and when both have failed it is frequently possible to successfully catheterize the tube by resorting to a modification of the last described method, as follows:—introduce the catheter as in the first method, carrying the tip well down into the pharynx, and when it has reached the posterior wall draw it forward without altering its plane until the curve is felt to impinge on the velum, then turn it up and out when the tip will be found to have slipped into the orifice of the tube. Considerable experience is necessary to practice this method successfully, but when once proficient in the other methods this one will be found very simple and effective in difficult cases. Before the introduction of the catheter the external extremity should be connected with the Politzer bag by means of a piece of rubber tubing about a foot in length, and the bag should have an aperture on the side which can be closed by the thumb of the surgeon when it is compressed, so that it will be possible for air to enter the bag again without the necessity of disconnecting the tip of the Politzer bag after each compression of the bag, a procedure that is usually extremely disagreeable to the patient.

I prefer, in adults, catheterization to the inflation of the ear by Politzer's method, whenever practicable, for several reasons: 1. The air enters the middle ear much more directly, and much more is accomplished by this method when it is carefully performed. 2. Very frequently it is necessary to force air into only one middle

ear. Politzer's method should always be employed in the case of children and nervous individuals.

Before inflating the ear the drumhead should be examined, to see if atrophic changes may not have made it very thin. During the first compression of the air-bag very little force should be used. I have seen the drumhead ruptured by Politzer's method, when sufficient gentleness was not used, in a case in which the drumhead had become very thin.

The obstacles to the introduction of the catheter are various pathological processes that involve the nasal passages, such as hypertrophic catarrh, polypi, adenoid growths, and congenital deformities of the nasal septum and spongy bones. In acute inflammation of the middle ear, with swelling of the orifice of the Eustachian tube, catheterization is difficult and painful, as well as in some individuals in whom its accomplishment is prevented by spasmodic coughing or vomiting. When catheterization is impossible, owing to constriction of the nasal passages, air has been forced into the middle ear by the introduction of a catheter through the opposite nasal passage. Various catheters have been devised for this purpose, the method having first been advocated by Deleau in 1827.

The diagnostic tube should be used, as in Politzer's method, to determine the permeability of the Eustachian tube, as it is unwise to rely on the statement of the patient.

TESTS FOR HEARING.

The tests for the hearing-power are of the highest importance, for by such means the surgeon is usually enabled not only to locate the seat of the trouble,

whether in the middle or inner ears, but also to note the course of the disease and the results obtained by treatment. The ear perceives sound-waves communicated to it through the air as well as vibrations transmitted through the cranial bones, so that it is most important in all cases to make separate tests of this power of perceiving these sounds.

The watch and Politzer's acoumeter are generally used for determining the acuteness of hearing for sound waves transmitted through the air to the drumhead. It is best to have two watches for this purpose, the tick of one being higher than that of the other. First of all, the distance heard by the normal ear is determined and found to be, for example, fifty-six inches. This number will represent the denominator, so that if a patient hears the watch at ten or twelve inches his hearing-distance for the watch should be recorded as being ten-fifty-sixths or twelve-fifty-sixths of the normal distance. The patient's normal ear should be carefully stopped by inserting the moistened finger into the meatus; then the watch should be held at the normal distance from the affected ear and gradually brought toward it. The patient's eyes should be closed. It is well also to let the patient hear the watch tick before commencing the test. If the watch is placed in contact with the ear and gradually moved further away, the patient will hear the tick at a much greater distance than if the reverse is done.

As all watches vary very much as to the pitch and intensity of sound, tests made in this way are necessarily unsatisfactory. For this reason Politzer has devised an acoumeter, which consists of a horizontal steel cylinder, 28 mm. long and 4.5 mm. thick, con-

nected by means of a tightly fitting screw with a perpendicular vulcanite column, at the superior and inferior extremities of which are two semicircles for holding the instrument, the forefinger being placed in the upper and the thumb in the lower one. A percussion hammer is placed just above the attachment of the cylinder, being fastened in an oval orifice of the vulcanite column. This hammer, falling upon the steel cylinder, produces the sound. The posterior short lever of the hammer, when pressed, is not allowed to go beyond a check, which consists of a soft India-rubber plate, so that the hammer always falls the same height and should produce the same sound. There is a hole in the vulcanite column for the insertion of a pin (Fig. 36, 8) having a round metal plate attached. This pin is inserted and applied to the temple or mastoid process, in order to determine sound vibrations through the cranial bones, and also in cases of extreme deafness, when it is placed in contact with the external meatus. The hearing-distance for the acoumeter should be measured by means of a tape-measure, or, better still, the number of feet and inches should be marked on the wall or floor. The only objection to this instrument is the fact that it is heard by the normal ear at a distance of forty-five feet, so that in cases of slight deafness it is better to make use of the watch. The patient's hearing-power for speech should always be tested, viz., for ordinary conversation as well as for the whisper. Investigations made by Oscar Wolf go to show that vowels are generally perceived at a greater distance and more readily than consonants, and especially words in which the vowels a, e, and i occur. Frequently in patients the hearing-power for the watch is very good, while con-

versation is heard with difficulty, and *vice versa*. Whispered speech is heard by the average normal ear in an absolutely still room at a distance of about fifty feet. It should be remembered that the hearing-power in a patient is subject to great variations, not only on different days, but on the same day and frequently it varies with the condition of the weather. In those affected with atrophic catarrh this is particularly noticeable, the auditory nerve becoming fatigued in a very short space of time, even while the tests for hearing are being made. Note should be made in taking the history of a case as to whether the patient hears loud, medium, or ordinary tones of speech. Tests made with the voice are not wholly satisfactory, as it is difficult to raise the voice to the same pitch and give the forced whisper uniform intensity at each examination. Each ear should be tested separately, and the ear being examined should be directed toward the surgeon, while the other one should be closed with the moistened finger. With eyes shut so that he cannot see the lips of the examiner, the patient should be asked to repeat the words spoken by the surgeon, so as to make the test an accurate one.

Beside the watch and acoumeter, the aural surgeon should be equipped with (1) several tuning-forks, usually three in number, C—128, C^2—512, C^4—2048 (Politzer), or better still Hartmann's series of five forks C—C^4, (Fig. 36), (2) König rods, (3) a Galton whistle, (4) several shrill whistles and musical instruments for testing the hearing of those who are extremely deaf. Experiments prove that low-toned tuning-forks, when vibrating, are heard better when held close to the ear than the high ones in cases of labyrinthine disease, while

in middle-ear affections in which there is considerable inability to hear speech, low-toned tuning-forks are but faintly heard. It is well to have clamps attached to the extremities of the forks to prevent overtones. A fork should never be struck with or against a hard substance of any sort. A soft rubber percussion hammer such as is shown in Fig. 36 will yield a pure tone.

Fig. 36.



1-5. Hartmann's series of tuning-forks. 6. Politzer's low fork. 7. König's rods with hammer. 8. Politzer's acoumeter. 9. Hammer for striking forks. 10. Galton whistle.

By means of the tuning-fork a diagnosis can usually be made between disease of the middle and internal ears. In a normal case the tuning-fork, when vibrating and held close to the ear, will be heard more distinctly and louder than when placed against the mastoid process or on the side of the skull. In other words, the aërial

conduction will be better than the bony conduction. If the finger is inserted in the external meatus, the sound will be intensified on the closed side when the tuning-fork is placed against the mastoid process or on the top of the head or against the teeth. The same is true of disease involving the external or middle ear. In a case of deafness, when it is found that a patient hears the tuning-fork better by the air than by the bone, it is fair to assume implication of the sound-perceiving apparatus; and when the bone-conduction is lost, it is certain that labyrinthine disease exists. Where one tuning-fork alone is used, the tuning-fork of Blake, 512 v. s. (see

Fig. 37.

Blake's tuning-fork.

Fig. 37), is to be preferred. As a general rule, although there are exceptions, the higher tuning-forks are heard much better by aërial conduction than the lower ones in cases of disease of the middle ear.

The Galton whistle is especially suited for testing the perception of the high notes, and consists of a cylindrical whistle. By turning the screw the note can be changed, and around the side of the whistle is a scale to show the numbers. The whistle has a compass of more than the three highest octaves, with about 6000 to 80,000 vibrations per second. Beside the Galton whistle, the König rods will be found particularly useful. They consist of rods or cylinders of steel, which are of different lengths, producing vibrations of from

20,000 to 80,000 in the second, with intervals of 5,000 vibrations. They are suspended at each end by a loop of fine wire or silk thread and struck with a steel hammer as shown in Fig. 36.

The power of perception of sound transmitted through the cranial bones becomes very much diminished in old age. According to Politzer, after the sixtieth year the cases are rare in which a low-ticking watch is heard through the cranial bones.

RINNÉ'S TEST is a relative duration test, and consists in placing a vibrating tuning-fork against the side of the skull or mastoid process. When the sound is no longer perceived by bone-conduction the fork should be held close to the auricle without again striking it. If heard again, which is the case in the normal ear, this is called the *positive Rinné* test. The reverse is called the *negative Rinné* test. This test is of particular value in cases of considerable deafness due to middle-ear affections, the negative Rinné test being marked in these cases unless there is implication of the labyrinth. In middle-ear affections in which the deafness is slight, Weber's test is more accurate.

WEBER'S TEST. A vibrating tuning-fork, usually of 512 v. s., is held on the median line of the skull, or on the forehead just above the nose, or against the teeth, or on the median line of the lower jaw, and the conclusion can be reached as to whether the bone-conduction is better on one side or the other. With both Weber's and Rinné's tests in ordinary cases of deafness a diagnosis should be made between disease of the middle ear and labyrinth.

SCHWABACH'S TEST is an actual duration test in which Hartmann's series of five tuning-forks is used.

The forks are known as C = 128, C^{i} = 256, C^{ii} = 512, C^{iii} = 1024, and C^{iv}=2048 d. v. s.=double vibrations per second. The method consists in ascertaining the number of seconds the patient can hear each fork by air-conduction (A. C.) and by bone-conduction (B. C.). For example, a fork (say C.) is held close to the external auditory meatus; the time is noted at the instant of applying the vibrating fork to the meatus, and again when the patient raises his hand to indicate that the fork is no longer heard. This will give the time for A. C. The same fork is then vibrated again, the end is held firmly against the skull over the antrum of the same side, and the time taken and recorded as before, which will represent the time of B. C. for the C fork on that side.

The result of this test in a case of normal hearing is shown in the following table:[1]

	C	C^{i}	C^{ii}	C^{iii}	C^{iv}
A. C.	25	15	33	32	22
B. C.	13	7	13	13	14

Occasionally it happens, although rarely, that the vibrations of a tuning-fork sound one-quarter or one-half a tone higher in one ear than the other (*diplacusis*). This is probably due to increased tension of the labyrinthine fluids produced by pressure of the stapes, or it may be due to congestion of the labyrinth itself.

[1] After John E. Sheppard, M. D., in American Text-Book of Diseases of Eye, Ear, Nose, and Throat.

Only one ear is apt to be affected, and it is an easy matter for the surgeon to determine which one it is by holding the tuning-fork close to his own ear, so that when the patient sings the note heard he recognizes which side is affected.

I have observed this phenomenon in both middle-ear and labyrinthine diseases. In the former the treatment consists of inflation of the middle ear by means of the catheter, and of appropriate treatment for the naso-pharynx, if this region is affected. In labyrinthine disease I have seen it occur in a patient who had a severe attack of influenza. In this case the middle ear was not involved, and the symptoms persisted for a long time.

A case of otitis media catarrhalis chronica at the Vanderbilt clinic;

	C	C^{i}	C^{ii}	C^{iii}	C^{iv}
A.C.	0	0	7	12.5	10
B.C.	18	7	10	14	0

A case of labyrinthine disease at the Vanderbilt clinic;

	C	C^{i}	C^{ii}	C^{iii}	C^{iv}
A.C.	0	6.5	4	3	5
B.C.	6	0	0	0	0

CHAPTER III.

DISEASES OF THE AURICLE.

MALFORMATIONS.

MALFORMATIONS of the organ of hearing may exist alone or be associated with malformations of other

FIG. 38.

Case of polyotia.

organs, such as want of development of the eye or side of the face on the same side. There may be congenital

excessive development of the auricle, the development of two auricles (polyotia), or so-called auricular appendages. (See Figs. 38, 39 and 40.) According to Cassells, excessive development is usually limited to the

FIG. 39.

Case of polyotia.

auricle itself, while want of development is apt to occur with defect of development of the external meatus and tympanum, less frequently with defect of the internal ear. There may be entire absence of the auricle, mal-

formation or absence of the different portions of the ear, viz., the lobule, helix, and antihelix. Imperfect development (microtia) is more frequent. (Fig. 41.) The auricle is generally out of place in such cases, and may resemble a cat's ear; the lobule is at times adherent to the skin.

FIG. 40.

Showing result of operation for polyotia.

I have reported a case[1] of malformation of the auricle in which I performed a plastic operation for the cure of the deformity, which occurred in a girl of fourteen years. The auricle was attached somewhat higher than the normal one on the other side, and hung

[1] Archives of Otology, 1890, vol. xix. No. 1.

down as shown in Fig. 42. The hearing-distance was considerably affected.—The operation consisted in the removal of an eliptical portion of the integument and connective tissue, three-quarters of an inch in width and two and a quarter inches long, from the posterior and upper portions of the auricle, commencing the

FIG. 41.

Case of microtia.

incision at the beginning of the helix and carrying it along the fold at the attachment of the auricle almost to the lobule, so as to leave as little scar as possible. The integument and connective tissue having been dissected away, the cartilage was bent back upon itself, but not divided, and the different folds and depressions

on the auricle were brought out again by suturing the raw edges of the wound. It was necessary to keep a pad with an elastic band over the ear for a short time. The result was excellent and the hearing-distance was improved (Fig. 43). This operation may likewise be performed in cases of very prominent auricles. Congenital deformities are, according to Virchow, due to early disturbances in the closure of the first branchial cleft, and are often associated with cleft palate and other forms of arrest of development in the facial bones.

A *fistula auris congenita* is a short, blind canal, with a lining of epithelium which secretes a whitish or milky fluid. The orifice is usually visible just in front of or below the tragus, but there is no communication with either the tympanum or the external auditory canal.

INJURIES OF THE AURICLE.

Injuries of the auricle are not frequently observed by the aural surgeon, as patients so afflicted are more likely to consult the general surgeon. Cuts, bruises, burns, occasionally occur, and should be treated according to the usual surgical procedures. A case has been reported in which the entire auricle was bitten off by a horse. Traction on an ear-ring occasionally is the cause of a cleft lobule. The wearing of brass ear-rings at times sets up a severe inflammation of the lobule, and in a case that came under my observation there was intense inflammation, followed by sloughing of half of the lobule on each side. Since foot-ball has become so popular a game, injuries of the auricle are of more frequent occurrence. In stitching an incised wound, it is well not to include the cartilage, but simply to carry the sutures through the integument; otherwise there is a liability that necrosis of the cartilage may occur.

FIG. 42.

Deformed auricle before operation, showing elliptical incision with sutures in position.

FIG. 43.

Deformed auricle. (After operation.)

HÆMATOMA AURIS, OR OTHÆMATOMA.

An hæmatoma is the name applied to a swelling, of a bluish-red color, which is due to a sudden effusion of blood between the cartilage and perichondrium, and which is usually caused by a sudden blow on the ear, and is of frequent occurrence among boxers. Such a tumor is at times observed among the insane, and formerly it was supposed to be due to ill-treatment. This may have been the case in some patients, but it is more probable that the effusion in such instances is caused by tissue-changes in the auricle itself. By severing the restiform body in animals, Brown-Séquard was able to produce a hemorrhage in the auricle. The swelling, when observed immediately after its appearance, is of a bluish-red color, roundish in shape, somewhat hard, but with no distinct point of fluctuation. In some instances it is quite soft to the touch. Othæmatoma occurring spontaneously is rarely of large size, while in cases due to traumatism the swelling may involve the entire auricle, including both surfaces, and occlude the external meatus, thus producing considerable deafness. Generally, the swelling involves the concha and fossæ of the helix and antihelix, but not the lobule. In othæmatoma due to traumatism there are usually pain, heat, and more or less tenderness. The resulting deformity depends very much on the extent of the hemorrhage and the injury done to the cartilage. Othæmatoma should not be confounded with perichondritis auriculæ, as the former occurs suddenly, and is generally caused by a blow.

As to the treatment, when the swelling is slight, the application of cooling lotions, such as the lead and opium wash, may cause the hæmatoma to disappear. It is much better, however, especially when there is

much swelling, to incise the tumor at once and turn out the blood-clots, and treat the wound antiseptically. In a case treated recently in this manner the cavity of the tumor was thoroughly washed out with a 1 : 10,000 bichloride solution, and the edges of the wound sutured except at the lowest point, where a small tent of iodoform gauze was inserted. A pad and bandage were applied, with the result that there was little if any deformity. Massage is recommended by some, but should be used cautiously.

PERICHONDRITIS AURICULÆ.

This is a term applied to an acute inflammation of the perichondrium, giving rise to a swelling which is generally hot and painful to the touch, of a darkish-red color, and varying much in size in different cases. It is usually found on the anterior aspect of the auricle, and may close the external meatus, and thus interfere seriously with the hearing on the affected side. If an incision were made in one of these tumors, its cavity would be found filled with a thick yellowish fluid, less frequently with pus, and only occasionally are there observed a few red blood-corpuscles. The perichondrium becomes detached from the cartilage, and the latter and the surrounding parts are greatly inflamed, leading to more or less thickening of the tissues, with deformity of the auricle. If an incision is made in the tumor and a probe introduced, the cartilage will frequently be found denuded and roughened. The inflammation occasionally has its starting-point in the external meatus, and generally involves the concha and fossa helicis, but does not extend to the lobule.

Perichondritis should be distinguished from hæmatoma (1) by the fact that traumatism is not apt to be

the cause; (2) in perichondritis the inflammatory symptoms are more severe and the development of the tumor is more gradual; (3) the contents of the swelling are either a clear, thick, syrupy fluid, or occasionally pus, while in othæmatoma there is always blood at first. In the later stages a differential diagnosis becomes more difficult, as an othæmatoma may also contain a clear syrupy fluid or pus. The deformity is quite marked in some cases of perichondritis, although in some few instances it may be but slight. The duration of the disease varies from two to ten weeks. In the early stage the treatment should be entirely antiphlogistic, consisting in the application of cooling lotions, lead and opium wash. Later, the tumor should be incised as soon as fluctuation is detected, and thorough aseptic precautions should be observed. A pad and pressure bandage will be found beneficial in some cases as well as massage. Some authors advise the use of a hypodermic needle for the purpose of withdrawing the fluid, and afterward injecting into the cavity a solution of tincture of iodine one part, water two parts. After allowing it to remain for a few moments it should be withdrawn by means of the syringe.

CYSTS.

Under this heading cases have been reported in which tumors in the auricle have developed without apparent signs of inflammation. The contents consist of a clear fluid without any flocculi or blood-corpuscles. As for a differential diagnosis between cysts and perichondritis, Hartmann considers that "inflammatory phenomena are altogether absent in cases of cysts, while perichondritis is accompanied with redness, heat, and severe pain. The appearance of the secretion in peri-

chondritis is similar to that in cysts, yet the fluid has an unclean, purulent character from the admixture of pus-corpuscles." Certain changes which predispose to the formation of sanguineous tumors are found in the auricle, especially in the dyscrasic and insane as well as in the aged. Hartmann believes that the process of softening which has been described by different authorities is not only the predisposing cause of hæmatoma, but is also to be regarded as a preliminary stage of simple cyst-formation. Cases of hæmatoma due to traumatism occur quite frequently, however, where there are no changes in the cartilage.

HYPERÆMIA.

This is observed in some cases as a result of frostbite or eczema, or other cutaneous diseases; also in patients of a neuropathic and hysterical temperament, and in those suffering from valvular disease of the heart. Hyperæmia may also be due to the action of heat or cold. The application of poultices may also produce an hyperæmic condition and lead to the formation of small abscesses. As to treatment, the cause should be removed, if possible, by constitutional measures. Locally, the lead and opium wash is to be recommended.

DERMATITIS.

Inflammation of the skin of the auricle may be due to the sting of insects, the effects of heat or cold, or to blows, falls, piercing the ears for ear-rings, and mechanical irritation and scalding. The inflammation may be very slight, or severe enough to lead to gangrene. In the early stage antiphlogistic treatment should be employed, viz., by making applications of cooling lotions.

ERYSIPELAS.

According to some writers, erysipelas affecting the auricle may occur as a primary or idiopathic disease from cold or other causes, but more frequently it is due to an extension of the inflammation from the surrounding parts. Leech-bites sometimes cause erysipelatous inflammation, as well as injuries to the walls of the external auditory canal, in persons whose general health is impaired. In the light of modern science, it is now generally admitted that erysipelas is due to invasion by Fehleisen's streptococcus erysipelatis.

Two cases[1] of erysipelas of the auricle and face, occurring as a complication of disease of the ear and secondary to it, came under my observation. In one the patient had an acute suppurative otitis media, and two weeks later the antitragus became tender and inflamed. On the following day the entire auricle was involved in an erysipelatous inflammation, and was twice its natural size. The auditory canal was not involved, but the disease extended over the scalp and face. When the disease first appeared the discharge from the ear became scanty. The patient had a chill at first, and the temperature varied from 100.5° to 104.5°. He complained also of dizziness. In the second case erysipelas of the auricle followed an attack of acute suppurative otitis media and furunculosis of the external meatus. The erysipelatous inflammation spread from the tragus of the left ear to the scalp and face, and involved the right ear as well. This patient had a temperature ranging from 102.4° to 104.6°. In both cases the pulse was slow and full, and the general

[1] Archives of Otology, 1887, vol. xvi. No. 2.

health was below par. In the second case the erysipelas lasted for a week. It is also probable that erysipelas of the pharynx may spread to the face by means of the Eustachian tube, tympanum, and external auditory canal.

As to treatment, I have seen excellent results follow the administration of drop-doses of tincture of aconite and tincture of belladonna. Locally, the lead and opium wash should be applied, or 25 to 50 per cent. of ichthyol ointment.

FROSTBITE.

If the auricle is frozen, the ear becomes yellowish white in color, and in extreme cases, if actually congealed, it may be brittle. To restore the circulation in the parts, the patient should be placed in a cool room, and friction applied, very gently at first, to the auricle with snow, and afterward with cold water. After the re-establishment of the circulation warmer water may be used. In favorable cases the circulation becomes re-established, but the auricle is apt to be sensitive for some time afterward. Where the auricle has been frozen and proper treatment has not been carried out, there may result severe inflammation, such as perichondritis, leading ultimately to gangrene. In the latter case amputation becomes necessary. Redness and burning of the auricle resulting from frostbite should be treated with cooling lotions; when chronic, by the local application of iodized collodion. When the skin is excoriated, equal parts of zinc oxide and boric acid should be dusted over the parts. When ulcerations occur as well as gangrene, general surgical procedures should be adopted.

ECZEMA, ACUTE AND CHRONIC.

Eczema affecting the auricle is very similar to the disease occurring elsewhere, except that the hearing is liable to be affected, especially when the auditory canal and membrana tympani are involved. The varieties are the erythematous, vesicular, pustular, and squamous. The disease may be acute or chronic, although the former is more usual; it may attack all ages and both sexes, but is more frequent among children and females. Eczema occurring about the pinna may be idiopathic or secondary to that occurring about the head or face. In the acute idiopathic variety one or both auricles may be attacked, or the disease may be limited to one side. The auditory canal may or may not be involved. In the acute stage the ears are usually swollen, the seat of much burning and itching, and covered by vesicles or pustules, which may become dry and thus envelop the auricle with unsightly crusts. The auricle, in consequence, becomes deformed and unshapely; the meatus may be closed by the collection of crusts or through swelling of the parts, and, as a result, the hearing becomes impaired, so that a sense of fulness in the ears and tinnitus are often complained of. Painful rhagades may be found behind the ear. In chronic cases the auricle may be considerably thickened, leading to deformity and stenosis of the external meatus (Fig. 44). In children eczema is aggravated by uncleanliness, by picking at the ear, and by the use of unsuitable head-coverings, by which means the ears are pressed against the side of the head, thus causing an *eczema intertrigo* through friction of the surfaces. Anything

that tends to impair the general health, such as dentition, will act as an exciting cause.

An artificial eczema is frequently due to the instillation of improper ear-drops, an excoriating discharge from the ear, the effects of heat and cold, and, especially in children, to pediculi about the head.

In acute eczema good results often follow the administration of the fluid extract of viola tricolor in x-xx minim doses, especially in young subjects. Particular

FIG. 44.

Case of chronic eczema following frost bite.

attention should be paid to the patient's general health. The urine should be examined for albumin or sugar. Diuretics and alkaline waters are required for those of a gouty or rheumatic diathesis. In scrofulous children, cod-liver oil, syrup of the iodide of iron, and the compound syrup of the hypophosphites should be prescribed. In chronic eczema, Fowler's solution of arsenic should be administered. Nasopharyngeal catarrh, or middle-ear disease, if it exists, should receive careful treatment,

and the condition of the mouth should be noted, to see that there is no dental irritation. As regards local treatment, the officinal zinc ointment will be found all that is required, the unguentum diachyli of Hebra or Lassar's paste; ℞ zinc. oxid. amyl., aa. ʒij, acid. salicylic grs. v., vaselin alb. ℥ss.—M. The crusts should first be removed with sterile olive oil and then the parts well rubbed several times a day with the ointment. When the auditory canal is invaded, the crusts and scales should be removed gently with curettes, and then the zinc ointment or Lassar's paste applied, or, what is better, the canal may be dusted with a powder composed of equal parts of zinc oxide and boric acid. Occasionally it will be necessary to remove the scales from the meatus by syringing; but, if possible, no water should be used. Should the disease show a tendency to become chronic, especially if there is any thickening, the following ointment will answer well: Acidi tannici, ʒj; vaselini, ℥j.—M. or the following, ℞ zinc. oxid., amyl., aa. ʒij, acid. salicylic, grs. v., sulp. præcip., ol. rusci aa. grs. xxv. vaselin. alb. ℥ss.—M. In making applications to the external auditory canal, great care should be taken not to injure the membrana tympani. Another ointment for a dry and scaly eczema of the external meatus is the following: Acidi salicylici, gr. j; tr. benzoini, gtt. ij; ung. rosæ, ʒj.—M.

In such cases benefit will often follow the application of a solution of nitrate of silver, grs. x–xx to water ℥j, or of acetum cantharidis; but in painting the canal with the latter, great care should be exercised not to touch the membrana tympani. Ointments containing tar or white precipitate of mercury are occasionally indicated in chronic eczema.

The following skin diseases are also observed at times on the auricle, associated with the same diseases elsewhere on the body—viz., *herpes zoster, lupus, erythema, prurigo contagiosa, keloid, molluscum fibrosum, tinea tonsurans,* for the treatment of which the reader is referred to the standard works on dermatology.

Ossification of the auricle as a result of perichondritis has been reported, but is of very rare occurrence.

Deposition of the urates is noticed occasionally in the auricle of a gouty subject.

TUMORS OF THE AURICLE.

Under this heading may be mentioned *abscess, atheroma or sebaceous cyst, fibroma, myxofibroma, malignant growths, dermoid cysts, warts, papilloma, angioma, lipoma, nævus maternus, aneurismal dilatation of arteries, and cornu cutaneum.*

FIG. 45.

Case of atheroma.

ATHEROMA, or SEBACEOUS CYST, is due to the blocking of a sebaceous gland and the rapid increase of its contents. It is usually found in the integument

just behind the lobule and close to it, or in the lobule itself. The cyst should be removed with its sac entire, if possible. When the sac cannot be dissected out, the walls of the cyst should be curetted, so as to remove every trace of the sac as well as any purulent or cheesy matter. (Fig. 45.)

Fibroma. Cases of fibroma are observed at times, the growth, as a rule, involving the lobule. The disease may affect one or both lobules, and is occasionally due to piercing the ears, and is found quite frequently in negro women. A fibroma is made up of fibrous connective tissue, with an occasional admixture of mucous tissue; it is of a round or globular shape, somewhat movable, and of more or less firm consistency. After total removal the tumor is not apt to recur. I have reported a case of a fibroma[1] affecting the lobule of each auricle in the person of a girl, aged seventeen, who had had her ears pierced four years previously, and immediately afterward the lobules began to enlarge. Another case recently came to my attention at the Vanderbilt clinic of a negress, twenty years of age, who had a fibroma of each lobule, the right one about three-eighths of an inch in diameter, and the left considerably smaller. Both tumors were on the posterior surface and appeared to have had their origin from scar tissue due to piercing the ears for earrings two years previously. The tumors did not appear to develop sufficiently for the patient to notice them until eighteen months later. (See Fig. 46).

Dermoids occasionally occur. They have a capsule of connective tissue lined with epithelium and elements of the skin including hair follicles and fine hairs. The

[1] "Tumors of the Auricle," Annals of Ophthalmology and Otology July, 1893, vol. ii. No. 3.

contents consist of exfoliated epithelium, hairs, cholesterin crystals, fat and a liberal quantity of serum.

Fig. 46.

Case of fibroma of auricle.

EPITHELIOMA.

Epithelioma at times attacks the external meatus primarily but is more likely to spread to the canal from the auricle or face. It has been known to follow frost-bite of the pinna. In those cases in which excision has failed to effect a cure, the x-ray and radium have been used of late years with apparent success by some

observers while others have met with failures, and it is too early as yet to draw any definite conclusions as to the value of either method of treatment in establishing a permanent cure.

The other growths sometimes noticed in the auricle, and previously mentioned, should be excised, if possible, and treated on general surgical principles. Among the less frequent tumors that are occasionally observed may be mentioned *cylindroma*, or *myxoma cartilagineum*, originating in the cartilaginous *meatus*, and *enchondroma*, having its origin from the cartilaginous portion of the external auditory canal.

SYPHILITIC DISEASES.

Chancres and Gummata have been found on the auricle, but very infrequently.

Tubercular Syphilides on the auricle have been reported, and these may lead to deep ulcerations.

Papular, Pustular, and Squamous eruptions are observed, but usually in connection with similar eruptions about the scalp and forehead.

CHAPTER IV.

DISEASES OF THE EXTERNAL AUDITORY CANAL.

MALFORMATIONS.

Excesses of development occur among the malformations of the external auditory meatus, such as unusual dilatation and a double condition of the auditory canal. There may be two separate canals, or a blind canal behind the normal one.

Much more frequent, however, are the defects in the formation of the auditory canal which usually are associated not only with defects of the auricle, but also with defects of the middle ear, and frequently with arrest of development of the cranial bones. The canal may be very narrow or absent, with simply a depression where the meatus should exist. The contraction may be bony or membranous, usually the former. The hearing in such cases varies according to the amount of atresia, but even when there is complete occlusion, if the labyrinth is normal on the affected side, there will remain some hearing power.

As to the propriety of operating on such cases, the general opinion is decidedly against any such procedure, unless there is some urgent necessity for the operation, especially if the patient has one good ear. Even when there seems to be every indication that a bony canal exists beneath the integument, the surgeon will often

be disappointed if he attempts an operation. When an operation is performed, a crucial incision should be made through the soft parts down to the bone, and a silver tube should be inserted, to keep the passage open. In such cases there is always difficulty in accomplishing this, as granulations form rapidly and have a tendency *to force* the tube out.

OTITIS EXTERNA DIFFUSA ACUTA, OR ACUTE DIFFUSE INFLAMMATION OF THE EXTERNAL AUDITORY CANAL.

Acute diffuse inflammation of the external auditory canal may occur as a result of the instillation of improper solutions into the ear, and in a case under my observation "St. Jacob's Oil" caused acute inflammation of the external meatus, as well as of the middle ear. Scratching the canal with hairpins and other such instruments, to relieve itching, frequently causes diffuse inflammation. Sea-bathing is another cause. As an idiopathic disease, I believe this affection to be extremely infrequent, although cases do occur, followed by suppuration, in which the middle ear is not involved. Diffuse inflammation of the external meatus is of common occurrence in cases of middle-ear suppuration when the discharge is acrid; also in cases of furuncular inflammation of the external auditory canal.

Symptoms. In the early stage the *meatus* is greatly congested, especially so about the bony canal and external layer of the drumhead. The cartilaginous portion may in some cases be so swollen that but little of the osseous canal is visible. The external meatus becomes very narrow from the swelling of the tissues, and the dermal layer, being white and sodden from a serous

discharge, peels off, leaving the canal and drumhead extremely red and excoriated. Frequently the dermal layer peels off, and may be removed by syringing in the form of the canal. On microscopical examination of the dermal lining micrococci will be found. After removal of the finger-like tube the canal and drumhead will be found greatly congested and all landmarks of the latter will be absent, so that the boundary between the membrana tympani and canal is lost.

The *subjective symptoms* are radiating pains, especially when moving the jaw, and when the surgeon makes pressure on the tissues in front of the tragus. In some cases there are tinnitus and dizziness; in others, a sense of itching in the canal precedes the inflammation.

The duration of the attack varies from a few days to two weeks, or longer if the disease becomes chronic. The degree of deafness will depend very much on the severity of the inflammation. When the canal is very much narrowed and filled with dermal scales, or the outer layer of the drumhead is swollen, there will be considerable interference with the function of hearing.

Instead of recovery ensuing in due time, there may be several exacerbations, or ulceration may develop on the drumhead, leading to perforation of the membrane from without inward; or ulceration may occur on the osseous canal, and granulations may spring up from the exposed bone. If the disease becomes *chronic*, there is usually a thick, offensive discharge, containing scales of epidermis and pus-cells with micrococci. As the drumhead appears dullish red, difficulty will be experienced in making a differential diagnosis between diffuse inflammation and chronic otitis media purulenta, especially if there are granulations on the drumhead. There may be granulations and ulcerations on the walls of the canal,

which may lead to the formation of membranous bands, thus causing atresia of the external meatus.

As a result of chronic diffuse inflammation, there may follow hypertrophy of the cutis or periostitis of the bony meatus, as well as caries and necrosis. In some instances the inflammation may extend to the parotid gland.

Treatment. In acute cases in which the inflammation and pain are not severe, a saturated solution of boric acid should be instilled frequently into the ear. When the inflammation is severe, however, the artificial leech should be applied close to the tragus, and drops of a 4 per cent. solution of cocaine should be used, or drops containing liq. Magendie, ʒj; water, ℥j. If the canal becomes very narrow from swelling, free incisions should be made through the skin and periosteum. After the acute symptoms have subsided, especially if there is much discharge, and after removing all pus and scales from the meatus by means of the syringe and curettes, boric acid powder should be insufflated, or a powder containing equal parts of boric acid and zinc oxide. Urbantschitsch[1] has used with excellent results an ointment containing thigenol 3 to 4 parts; lanoline, vaseline, each 10 parts. It has a cauterizing effect so that its action must be watched. When the disease has become chronic, and especially if there are granulations, alcohol drops (either pure alcohol or diluted one-half with water and containing boric acid, grs. xx to ℥j) should be used. In some cases in which granulations arise from bare bone, these should be removed with a sharp spoon and the bone scraped. In other cases the granulations may be cauterized with nitrate of silver or chromic acid fused on the end of a probe.

[1] Monatschr f. Ohrenh, No. 4, 1903.

OTITIS EXTERNA CIRCUMSCRIPTA ACUTA, OR ACUTE CIRCUMSCRIBED INFLAMMATION OF THE EXTERNAL AUDITORY CANAL, OR FURUNCULOSIS.

This is quite a common complaint, especially among those who are suffering from impaired health, and it is occasionally observed among diabetic patients. Furuncles, however, are found at times in the case of those who are strong and healthy, but children are usually free from them. Mechanical irritation of any kind may be the exciting cause of a furuncle, such as too frequent syringing, scratching the ear with a hair-pin or some other instrument, to relieve itching of the external meatus. Furuncles may develop during the course of chronic purulent otitis media, or as a result of chronic eczema of the external meatus. There is undoubtedly a micro-organism present to account for the inflammation, most frequently the Staphylococcus pyogenes aureus and albus, which, according to Loewenberg, finds its way into the superficial layers or deeper parts of the skin by means of the hair-follicles or glands. The seat of the inflammation is, as a rule, in the cartilaginous portion of the external auditory canal.

Symptoms. Pain is a prominent symptom, varying in intensity according as to whether the furuncle is seated in the superficial layer of the cutis or in the deeper parts. The patient usually experiences great pain on moving the jaw and on making slight traction upon the auricle. The pain is generally very severe, and keeps the patient from getting any rest at night. There is usually loss of appetite, and fever may or may not occur. When the drumhead is involved, or

when the lumen of the canal is closed by a furuncle, tinnitus and deafness will be complained of. Another cause of deafness is the blocking up of the meatus with pus and scales. The seat of the furuncle is usually on the anterior inferior wall of the cartilaginous meatus, although there may be several furuncles. When situated on the anterior wall, pain is usually complained of when pressure is made on the tragus. When on the posterior wall, however, there are very often considerable swelling and redness of the tissues over the mastoid process, which symptoms are frequently mistaken for mastoid disease if a careful examination of the external auditory canal is not made.

In a case recently seen in consultation, which presented certain symptoms of mastoid disease, the tissues were hard, brawny, and œdematous, not only over the mastoid process, but also in the neck. The cartilaginous canal was slightly red and tender, and its calibre somewhat diminished. No furuncle could be seen. There was some redness of Shrapnell's membrane. The hearing was but slightly affected. The patient was a man, sixty-five years of age, broken down in health, and he had lately had several boils on the back of his neck. The hard and brawny tissues behind the auricle were freely incised down to the bone. A few days later a furuncle developed in the cartilaginous portion of the external meatus, proving the case to be one of furunculosis of the external auditory canal.

After from three to five days the furuncle will break down and discharge pus; the more superficial the furuncle the more readily will it discharge into the meatus. Occasionally the furuncular inflammation will subside without bursting. After evacuation of the pus there is

usually immediate relief, viz., the pain and other symptoms subside. This is not an invariable rule, however, for at times exacerbations occur, and other furuncles follow one another as a result of the invasion of the follicles or glands by the staphylococci. At times there will remain redness of the canal and infiltration of the tissues after the abscess has subsided, and occasionally granulation-tissue springs from the opening in the furuncle, which necessitates its removal with the curette or snare.

There is usually no difficulty in making a diagnosis of furunculosis, especially if a probe is used to touch the parts. A polypus is sometimes mistaken for a furuncle, and so is an exostosis, but the latter when touched with a probe will seem hard and bony.

Treatment. In the early stage, especially if the pain is severe, the artificial leech should be applied, and in some cases Leiter's coil will be found useful. Good results frequently are obtained by soaking a tampon of cotton in a solution of menthol 1–5, or carbolic glycerin 1–40, and introducing it in the canal to cause pressure. The tampon is kept in place for a day if possible and then should be removed. The following solution of Burow when warmed and dropped into the ear will often abort the disease; ℞ Liq. alumin. acetat., 12 parts, plumbi acetat., 5 parts, and distilled water 100 parts. When the furuncle is deep-seated and very painful, an early and free incision should be made in the abscess, and also in all cases in which fluctuation can be detected, as well as in those in which the abscess has burst spontaneously, but in which the opening is small. After incising the furuncle it is advisable to press out any core that it may contain and then to

syringe the external meatus with a 1 : 3000 bichloride of mercury solution, and to direct the patient to instil into the ear a warm saturated boric acid solution frequently during the day.

After the acute symptoms have subsided the patient should be directed to paint the canal with carbolic glycerin (1 : 30), or to instil drops of bichloride of mercury or boric acid solution, in order to prevent the bacteria from advancing further, and this treatment should be continued for some time.

Owing to the debilitated and anæmic condition frequently observed among these patients, particular attention should be paid to the general health. When the furuncle first appears, it is advisable to give the patient several small doses of calomel, to produce a slight laxative effect. The sulphide of calcium, in one-twentieth to one-tenth grain doses, administered every three hours, has seemed to me to have a curative effect in many cases. For chlorotic females large doses of iron should be prescribed, and the diet should be a nourishing one. Cod-liver oil and the syrup of the hypophosphites are also useful. After the disappearance of the furuncle the patient will often complain of considerable irritation and itching about the external meatus. To relieve this, ointments containing boric acid or white precipitate of mercury should be applied very lightly to the canal.

OTITIS EXTERNA HÆMORRHAGICA — HEMORRHAGIC INFLAMMATION OF THE EXTERNAL AUDITORY CANAL.

There seems to be a difference of opinion among otologists as to the significance of hemorrhagic vesicles on the walls of the bony auditory canal. The majority

of writers, if they mention the occurrence of these vesicles at all, attach no particular importance to them, considering them for the most part as evidence of middle-ear inflammation. That these vesicles are found in the canal in cases of severe inflammation, no one will deny who has had much experience in the treatment of aural cases. I have likewise seen several cases, which I have reported,[1] in which such effusions of blood occurred independently of any middle-ear inflammation, and to my mind constituted a hemorrhagic inflammation of the auditory canal. At first there are usually some pain, slight deafness, and tinnitus. On examination there will be found one or more elongated swellings in the bony canal, usually on its inferior wall, and extending to the drumhead on one side and to the cartilaginous meatus on the other. At times a vesicle is seen on the membrana tympani. The vesicle is due to inflammation of the cutis, the dermal layer being raised by the sanguineous fluid. The vesicles are soft when touched with a probe, and are easily punctured. The inflammation, as a rule, reaches its height on the third day, and the duration of the disease is from eight to fourteen days. The walls of the canal become covered with fine white scales. The treatment consists in puncturing the vesicles, evacuating their contents, and, after drying the canal with absorbent cotton on a probe, insufflating powdered boric acid.

OTITIS EXTERNA CROUPOSA.—CROUPOUS INFLAMMATION OF THE EXTERNAL AUDITORY CANAL.

This is a rare disease, in which the meatus and drumhead are covered by a layer of lymph similar to that found in croup occurring in the larynx. It is generally

[1] Archives of Otology, 1890, vol. xix. No. 1.

observed as secondary to otitis media or furunculosis of the canal. The exudate is made up of a fine network containing round cells, nuclei, and epithelium.

Prognosis. The prognosis is favorable. The false membrane should be removed by syringing and forceps, and boric acid powder should afterward be blown into the meatus.

OTITIS EXTERNA DIPHTHERITICA — DIPHTHERITIC INFLAMMATION OF THE EXTERNAL AUDITORY CANAL.

Diphtheritic inflammation of the external auditory canal is associated most frequently with diphtheria of the nasopharynx and middle ear, especially in cases of scarlet fever. Primary diphtheria of the meatus has been described by some writers as developing during an epidemic of that disease in a canal previously excoriated from diffuse inflammation. The exudation, of a dirty gray color, when removed leaves an excoriated and bleeding surface.

Prognosis. The prognosis of diphtheria of the canal alone is apt to be favorable, but diphtheria of the middle ear may lead to serious complications, such as necrosis and loss of ossicles, necrosis of the tympanum and external bony meatus. Great deafness from secondary involvement of the labyrinth may ensue. The glands at the angle of the jaw are frequently enlarged.

Treatment. Besides constitutional treatment, including the use of antitoxin, the ear should be frequently syringed with a bichloride of mercury solution (1 : 3000), and all scales, etc., removed with forceps. Afterward a powder consisting of equal parts of iodoform and boric acid should be insufflated.

SYPHILIS OF THE EXTERNAL AUDITORY CANAL.

Syphilitic affections of the canal are generally confined to condylomata, ulcerations, and at times gummata. Condylomata appear like granulations, but are differentiated from them by the existence of syphilitic manifestations on other parts of the body.

Gummata are seldom confined to the external meatus, but attack simultaneously the middle ear. A gumma may lead to deep ulceration and destruction of the concha. Syphilitic nodules make their appearance at times in the meatus and on the drum-head, and may develop into ulcers. If the patient is improperly treated, the deep ulceration may lead to stricture of the canal. Besides constitutional treatment, such as the administration of mercury and iodide of potash, the ear should be syringed with a bichloride solution (1 : 3000), and all secretion and scales carefully removed. The lotio nigra, or iodoform powder, should be used. In some cases it will be necessary to cauterize the gummata, if very large, with chromic acid fused on the end of a probe.

OTOMYCOSIS—PARASITIC INFLAMMATION OF THE EXTERNAL AUDITORY CANAL.

This is a name given to a parasitic inflammation of the external auditory canal. Aspergillus niger, or nigricans (see Fig. 47), is the fungus most commonly observed, although there are other varieties, viz., the Aspergillus flavus and Aspergillus fumigatus.

Etiology. The disease is said to be frequent among the Russians, who live in poorly ventilated and damp houses, dampness being favorable to the growth of the fungus. According to some writers, an abnormal secre-

tion in the meatus due to eczema, or a slight discharge in chronic purulent otitis media, favors the growth of the fungus, while others believe that fungous growths are developed after the instillation of oily solutions. Very frequently the cause of the disease cannot be ascertained.

Otomycosis is observed rarely in children or aged persons, but more frequently among the middle-aged, while the poor are more often affected than those in better circumstances.

FIG. 47.

Aspergillus nigricans. (POLITZER.)

A. Mycelium covered with numerous fallen spores. B. Hypha. C. Sporangium with ripe spores. B'. Hypha. D. Receptaculum. E. Sterigmata with spores.

Symptoms. When the superficial layers alone are affected there are but few symptoms; but later, when inflammation occurs, the patient complains of itching and pain, which may be quite sharp in some instances. There are usually some tinnitus and deafness.

Appearances of the External Meatus. The canal, espe-

cially the bony portion, will be covered with loose scales, which, in the case of Aspergillus niger and fumigatus, are studded with blackish spots. In cases of Aspergillus flavus the epithelial scales are covered with mycelium, which appears like a yellowish powder. After removing the epithelial scales the canal and drumhead are usually found red and swollen. The appearances of the disease are quite characteristic when observed through the speculum; but in order to be sure of a diagnosis a microscopic examination is necessary.

Besides the epithelial scales and network of mycelium, one observes under the microscope cylindrical filaments or fruit-stalks, which are supplied with septa or hyphæ, and which support the fruit capsule or head of the fungus.

Prognosis. The prognosis is favorable when the case is properly treated, but if neglected the disease may continue for a long time, with many relapses, until the fungus becomes exhausted.

Treatment. After removal of all scales, etc., from the canal, the ear should be filled with a solution of bichloride of mercury in alcohol (1 : 3000). Absolute alcohol should be used if it does not cause too much burning, and should always be instilled warm. Otherwise water should be added to the alcohol. This treatment should be carried out two or three times a day for several days, and occasionally after this for several weeks, in order to be certain of destroying the fungus. Other solutions which are recommended are 2 per cent. alcoholic solution of boric acid or salicylic acid. Some surgeons prefer boric acid powder instead of an alcoholic solution. Bar[1] of Nice believes that in order to cure the disease,

[1] Annales des mal. de l'oreille, Feb., 1905.

one must render alkaline and antiseptic the locality itself. With this object in view, he uses a 4 per cent. solution of bicarbonate of soda and a 1 per cent. solution of corrosive sublimate.

POLYPI.

Granulations and polypi are found at times in the external auditory meatus, as a result usually of furuncles or of caries or necrosis of the bony walls. For the treatment of such cases, the reader is referred to the chapter on Polypi of the Middle Ear.

ECZEMA, ACUTE AND CHRONIC.

Eczema of the auricle has already been described, and the reader is referred to that chapter for treatment of the disease affecting the external meatus. Reference has also been made in the same chapter to herpes zoster, lupus, and other skin affections involving the auricle.

In herpes zoster the vesicles may be found not only on the posterior surface of the auricle, and especially the lobule and the parts in front of the tragus, but also on the anterior superior wall of the external meatus. Vesicles have also been found on the membrana tympani.

SEBORRHŒA.

A seborrhœal condition of the auditory canal is observed at times, when it is apt to be associated with a seborrhœa of the face and scalp. It gives rise to more or less itching, deafness, and tinnitus, especially if the external meatus contains moist scales and seborrhœal fluid. After the scales have been removed, the best results will be obtained from the insufflation of boric

acid, or of boric acid and zinc oxide powder. In some cases ointments containing ammoniated mercury, or the red oxide, are to be preferred, while especial attention should be paid to the patient's general health.

SEBACEOUS TUMORS.

These tumors are found in the external auditory canal, and should be treated in the same manner as those in the auricle.

CERUMEN INSPISSATUM OR INSPISSATED CERUMEN.

Inspissated cerumen is a frequent cause of deafness, and when found in one ear it is very apt to be present in the other one as well, so that it is always wise to examine both canals. Cerumen may remain in the canal for years, and in consequence become as hard as bone. The cause of the accumulation seems to be due, in some instances, to a scaly condition of the walls of the canal, which prevents the cerumen from escaping as it should do. Once it begins to accumulate, a plug is formed rapidly. In other cases there seems to be an overabundant secretion of cerumen, especially in persons affected with acne and seborrhœa of the face. Anything which prevents the cerumen from coming out of the orifice, such as atresia of the canal, constriction from swollen tissues or bony growths, or numerous hairs about the tragus, will act as a cause. Another cause is due to the fact that patients frequently try to cleanse the ear by pouring soap and water into the canal, and then attempt to dry the ear with the end of a towel, thus pushing the mixed cerumen and soap further in.

Symptoms. Cerumen in large quantities may be present in the external meatus for a considerable period without causing any inconvenience, as long as it does not rest against the drumhead, and provided there is a passage for the transmission of the waves of sound to the membrana. When the canal is closed or the plug falls against the membrana tympani, the patient will then complain of a sense of fulness and deafness. Other symptoms are autophony, or a hollow sound to one's voice like "talking into a barrel"; occasionally, giddiness, vertigo, reflex cough and hallucinations of hearing. Epileptic attacks have been caused by the presence of hardened cerumen in the auditory canal. When the plug is very large, it may irritate the walls of the canal and give rise to more or less pain. In aged persons the external meatus becomes flattened, so that the cerumen is more likely to accumulate. If cerumen collects and forms a plug, a similar accumulation is very likely to recur.

Laminated epithelial plugs, or *keratosis obturans*, may be a cause of deafness in some cases, the plugs being due to a gradual collection and accretion of epithelial laminæ derived from the external auditory canal. They give rise to the same symptoms as inspissated cerumen.

Diagnosis. Frequently a diagnosis can be made with the eye if the auricle is pulled upward and backward. After introducing a speculum, with light reflected from the head-mirror yellowish or brownish masses will be seen, and by touching the same with a probe a diagnosis of cerumen or epithelial plug can usually be made. Sometimes an error in diagnosis is made where there are numerous stiff hairs in the canal covered with cerumen. It is necessary to push the speculum beyond

these hairs. Again, the speculum may be directed against a layer of cerumen on the wall of the canal, or a wad of cotton or a foreign body may be smeared over with cerumen and be mistaken for a plug, or the canal may be filled with dry cholesteatomatous masses or dried blood. Tests made with tuning-forks will show increased bone-conduction on the affected side unless the inner ear is affected.

Prognosis. When the deafness has come on suddenly, complete restoration of hearing will usually result after removal of the cerumen. One should be guarded in ex-

Fig. 48.



Author's ear syringe.

pressing an opinion, however, before removing the plug, for the case may be complicated by adhesive catarrh or labyrinthine disease.

Treatment. When the cerumen or laminated epithelial plug is hard, it should be softened by the instillation of the following: Sodæ bicarb., gr. xxv; glycerin., ʒj; aquæ, ℥j. The drops should be allowed to remain in the canal for fifteen minutes before syringing. Frequently the plug is so hard that it is necessary for the patient to use this solution three or four times a day for several days before the mass can be removed with

the syringe. A syringe and warm water (containing in solution a teaspoonful of bicarbonate of soda to the pint) are the best means of removing cerumen. A convenient syringe is the one represented in Fig. 48, which is made of glass, with hard-rubber attachments and metallic rings. The hard rubber at each end can be unscrewed and the glass barrel can be kept very clean. The syringe holds about two ounces. The point of the syringe is quite small, to enable the surgeon to direct the stream of water between the wall of the canal and the plug. This is not a safe syringe to place in the hands of the

FIG. 49.



Metal syringe with solid packing.

patient. The soft-rubber bulb is generally recommended for patients, or the hard-rubber syringe with a large tip. Another excellent syringe, (Fig. 49), made entirely of metal with solid packing is easily sterilized.

Before using the syringe, the surgeon should be sure of good illumination, and should direct the stream of water between the plug and the canal at different points, in order to have the return current force the obstruction outward. The ordinary pus basin (see Fig. 50), held against the neck and just below the auricle, makes an excellent basin with which to catch the return flow

from the meatus, or the vessel (see Fig. 51) especially devised for the purpose. A graduated glass vessel, containing a solution of bicarbonate of soda, should be used for holding the clean water, and under no circum-

FIG. 50.

Pus basin.

stances should the same vessel be used for both purposes. Very little force should be employed in syringing the ear, for fear not only of doing damage to the drumhead, but also on account of the danger of causing

FIG. 51.



vertigo or faintness in the patient. The surgeon should also inspect the ear from time to time, and frequently it will be necessary to remove the plug with curette forceps (Fig. 23) or with blunt curettes, of which there are three sizes. The syringe should always be used, if pos-

sible, and only in skilled hands and under good illumination from a head-mirror should curettes or other instruments be employed. After the cerumen has been removed the canal should be dried carefully with absorbent cotton on a cotton-holder, made of aluminium (Fig. 25), with a roughened end, and afterward the ear should be inflated by Politzer's method, as the drum-head will usually be found more or less retracted. Then a piece of absorbent cotton should be inserted in the meatus and worn for the rest of the day. In some cases the walls of the canal may be found excoriated, and in such it will be necessary, after drying the canal, to insufflate powdered boric acid and zinc oxide (equal parts). When the plug consists for the most part of layers of epithelium, frequent syringing will often be insufficient, so that it will be necessary to use forceps.

FOREIGN BODIES.

Foreign bodies found in the external meatus are of every shape and variety. There are :

1. Those that are animate — consisting of various insects, viz., fleas, bed-bugs, cockroaches, etc.

2. Those that are inanimate, such as wads of cotton, pieces of paper, ends of slate-pencils, fruit-kernels, beads, cherry-stones, etc.

3. Those that are liable to swell and thus become enlarged, as beans, onions, kernels of wheat, etc.

Foreign bodies that do not swell and which do not cause irritation or inflammation of the canal may remain *in situ* for many years without causing any discomfort. Children are much more apt to put foreign bodies in their ears than adults.

Symptoms. There is usually more or less disturbance of hearing to almost complete deafness, depending on the size, shape, and position of the foreign body. There may be a sensation of pressure or tension and subjective noises, accompanied by more or less pain, according to the irritation and inflammation of the canal, drum membrane, or middle ear. The most serious symptoms often arise from unskilful attempts at removal of foreign bodies, when the auditory canal is lacerated and the drumhead is injured, and severe inflammation results. There may be certain reflex phenomena, such as pain in the head, vertigo, dizziness, vomiting, and coughing. In one case reported, a persistent cough with vomiting was caused by the pressure of two beans in the auditory canal, while in rare instances attacks of epilepsy have been reported as cured after the removal of foreign bodies.

Prognosis. The prognosis is usually favorable except in cases in which attempts on the part of an ignorant person have resulted in serious injury to the canal, drumhead, and middle ear; or in which the foreign body has become wedged in the deeper parts of the canal or middle ear, and severe inflammation has followed; or in which pus has not found an outlet, and brain complications have developed.

Treatment. In regard to treatment, much depends on the nature of the foreign body, whether impacted or not, the length of time that it has been in the ear, and the amount of irritation and inflammation of the parts. The first and most important step for the surgeon is to examine the ear with a speculum and good illumination, and determine for himself, and not rely on the patient's statement as to the presence of a foreign body in the

ear, for I have seen serious harm done when this precaution had been neglected and instruments had been unskilfully used. When this unfortunate accident has happened, it will be necessary at first to reduce the inflammation before attempting the extraction of the foreign body. When, however, there is but little, if any, swelling or inflammation of the canal, and the foreign substance is not wedged in, the syringe and warm water are usually all that are necessary to remove the foreign body. The auricle should be drawn backward and upward, while the stream of water is directed between the foreign body and the wall of the canal, and it is well at the same time to incline the patient's head toward the affected side. Some writers advise that the patient be placed in the recumbent position, so that the head will hang somewhat over the end of the table, and in this way the upper posterior wall of the auditory canal will be below the membrana tympani, and thus form with it an inclined plane.

In the case of foreign bodies likely to swell, oil or glycerin should be poured into the canal, and the ear should then be syringed, or after syringing with water enlargement of the foreign body can be prevented by immediately pouring alcohol into the ear. When the drumhead is perforated or the foreign body has been forced into the middle ear, some writers advise inflation of the ear to assist in the removal of the object. When an insect has gotten into the ear, it is best to kill it first by instillations of oil, when it can usually be removed by syringing or by means of forceps. Occasionally a very small insect will enter the ear and give great annoyance to the patient, while it may escape the observation of the surgeon. In such cases it is always well to syringe the ear.

When the patient comes for advice, and on examination it is found that the canal has been lacerated by attempts made at extraction, and especially if the foreign body cannot be seen, it is always best to allay the inflammation, if possible, before proceeding to any operation. If serious symptoms develop, such as severe pain and meningeal irritation, the surgeon must decide as to the propriety of an immediate operation. In such cases, where the syringe has failed, it will be necessary to resort to instruments, of which all sizes and shapes have been devised, viz., forceps, blunt and sharp hooks, curettes, loops of wire, scoops, etc. No one instrument is suitable for all cases. It is necessary at times to anæsthetize the patient before resorting to the use of instruments, especially when the body has become wedged in the deeper portions of the canal, and in the case of small children. The forceps shown in Fig. 24 and Fig. 52 are excellent ones. In using them, however, great care should be exercised not to push the foreign body further inward, if it cannot be grasped at once. A bent probe or a curette can frequently be passed around the foreign substance.

An old method of removing foreign bodies, which has been revived, consists in dipping the point of a camel's-hair brush into a solution of glue and touching with it the foreign body, which has previously been dried, and leaving it in position until the brush sticks, and then withdrawing it with the foreign body attached. This method is best adapted to round, smooth substances and to those liable to swell, and also to those cases in which there is no discharge from the meatus, which would prevent the glue from drying. When the membrana tympani is perforated, injections of warm

water through the Eustachian tube by means of the catheter will sometimes dislodge a deeply impacted foreign body. The galvano-cautery has at times been used to divide a foreign body piecemeal, which had become wedged in, before attempting to remove it. Occasionally it becomes necessary to detach the auricle and posterior wall of the cartilaginous canal by making an incision behind the pinna close to the canal, when

Fig. 52.

Allen's foreign-body forceps.

the foreign body has become impacted in the bony auditory canal or has been forced into the middle ear. This procedure allows the surgeon much more space for the use of instruments.

Nothing but the syringe and warm water should ever be used by the surgeon, in the case of a foreign body in the ear, unless he understands thoroughly the use of the speculum and head-mirror and has had considerable experience in the treatment of aural patients.

INGROWING HAIRS.

Hairs from the auditory meatus at times are so thick and long that they may interfere with the hearing, or they may even rest against the drumhead, giving rise to noises in the ear, especially when the patient moves his jaw. Epilation is required in extreme cases.

HEMORRHAGE FROM THE EAR.

Hemorrhage from the ear, beside being due to direct injuries from foreign bodies, may be caused by fracture of the bony canal following a blow on the jaw. Bleeding from the ear, or vicarious menstruation, sometimes occurs in females during the menstrual period.

EAR-COUGH.

This phenomenon is noticed frequently in patients when examination is made of the ear by the introduction of a speculum. Foreign bodies and impacted cerumen in the external meatus have been known to cause severe attacks of coughing. The explanation is that the cough, reflex in character, is due to irritation of the auricular branch of the pneumogastric (which latter supplies the auditory canal).

ADHESIONS AND CONTRACTIONS OF THE MEATUS —EXOSTOSIS.

Swelling and infiltration of the lining membrane of the external meatus may occur during the course of chronic suppurative otitis media, and in some cases of inflammation of the external auditory canal, and thus lead to contraction of the external meatus. During the course of long-standing suppuration from the middle

ear in cases due to diphtheria or syphilis, or to injuries of the canal from instruments or chemicals, cicatricial strictures may result, and give rise to alarming cerebral symptoms, if operative procedures are not immediately undertaken, so that pent-up secretions may escape. When the contraction is due simply to hypertrophy of the tissues from inflammatory infiltration, much may be accomplished by treatment. Attention should first be directed to the cure of the chronic otorrhœa if present, while the infiltration of the tissues will often subside under the use of ointments containing mercury. In the case of strictures, some are circular while others are annular in shape, and the opening, being as small as a pin's head, may be mistaken by the inexperienced observer for a perforation in the drumhead. The contraction may in other cases extend from the bony canal to the external orifice of the meatus. Bony strictures also are observed, which are due to periosteal ossification of the walls of the external meatus or to *hyperostoses*. The latter are found generally on the anterior and posterior walls of the external auditory bony canal. They are more frequent among "the rich than the poor" and in certain families there is undoubtedly a predisposition. Fracture of the temporal bone, involving the bony canal may also be an exciting cause. Osseous strictures are associated often with caries and necrosis of the temporal bone, and develop after sequestra have been removed. The deafness in these cases will depend altogether on the condition of the middle ear, and also on the amount of secretion or epithelial scales that is retained behind the stricture. In making a diagnosis of stricture, a careful examination should be made with the probe, to ascertain whether the stricture is due to membrane or to bone.

Treatment. The most important point in the treatment should be the cure of any otorrhœa or inflammation of the external auditory canal. If the inflammatory hypertrophy does not diminish under such treatment, the contracted parts may be dilated by inserting plugs of absorbent cotton soaked in a carbolic acid or boric acid solution, especial care being taken not to leave them in too long at a time, as secretions are apt to be retained behind them. An annular stricture should be excised and the canal further dilated by the use of cotton plugs or the insertion of sponge-tents. If these means fail, laminaria tents are sometimes employed, but, as a rule, they are too painful. I have in some cases inserted rubber tubing, through which a quill toothpick had been passed, the idea being to prevent collapse of the rubber tube, and thus allow the pus to escape, and at the same time to exert continuous pressure against the walls of the external meatus. In all cases in which there is suppuration, as in chronic otorrhœa, the pus should be removed carefully by syringing with a bichloride of mercury or boric acid solution before inserting the tube. Hard-rubber and silver tubes are used also for the same purpose. When the stricture is osseous, very little can be accomplished in the way of dilating the canal except by operation. The latter should never he undertaken unless pus or some other secretion is prevented from escaping, or there is extensive caries or necrosis. In such cases the auricle should be displaced forward and the lining of the posterior wall of the canal dissected away, to give a good view of the seat of the operation.

Beside *hyperostoses*, already mentioned, *exostoses* are found in the canal. These are new formations of bone.

Their structure may be very dense like ivory, or spongy, or even hollow; they may have very broad bases or be pedunculated; and they may exist singly in one or both ears, or may be multiple. I have seen cases in which there were many white nodules situated on the walls of the external auditory canal and close to Shrapnell's membrane. Those that are spherical in shape are usually found on the upper posterior wall of the external bony canal. They may remain moderately small or grow to such a size as to close the lumen of the canal. Koerner[1] found in one hundred and forty-seven cases of exostosis or hyperostosis of the external auditory canal that one hundred and fourteen were men and only thirty-three women. Also that people living by the sea were more frequently affected than those in inland countries, and that the disease was more frequent in England than on the continent. There is also an undoubted inherited tendency to the disease in certain families. Exostoses may be due to chronic periosteal inflammation of the bony canal as a result of fracture, or may result from chronic suppurative otitis media. Polypi and new cartilaginous growths occasionally ossify. Syphilis, gout, and rheumatism are mentioned as causes of these bony tumors, but, according to my experience, sea-bathing plays a much more important rôle.

Diagnosis. By means of the probe the diagnosis can usually be made, as the tumor will be found hard and resisting, and frequently very tender to the touch.

Treatment. When the exostoses are small and do not interfere with the hearing-power, and do not prevent the escape of cerumen and other secretion, no operative

[1]Zeitschr. f. Ohrenh, December, 1904.

measures are usually required. A pedunculated growth can easily be removed with the chisel and mallet, but an exostosis with a broad base, and of compact tissue, is not easily excised.

When it becomes necessary to operate because considerable deafness has resulted, or because alarming symptoms have ensued from retention of pus, the auricle should first be displaced forward and the tumor removed by means of the mallet and chisel, or gouge, or by the burr, which can be rotated by a motor. The most thorough aseptic precautions should be followed and all instruments thoroughly sterilized.

After the removal of an exostosis the hemorrhage should first be controlled, and then the canal should be filled with a strip of iodoform or boric acid gauze.

The use of mercury and iodide of potash is indicated in all cases in which there is a history of syphilis. Some surgeons recommend painting the exostosis with tincture of iodine, but such treatment is of little if any avail, while there is great danger of causing a severe inflammation of the external auditory canal from its application.

CHAPTER V.

DISEASES AND INJURIES OF THE DRUMHEAD AND MIDDLE EAR.

MYRINGITIS, ACUTE AND CHRONIC.

MANY otologists do not believe that inflammation of the membrana tympani (myringitis) ever occurs without simultaneous involvement of the middle ear. In the acute stage the drumhead will be found congested in part only or in its whole extent. In the later stage there is apt to be an effusion of serum in the dermal layer, as well as slight effusions of blood, while small vesicles, filled with serum, often develop. As a rule, the dermal layer alone is involved; but in severe inflammation abscesses sometimes form in the deeper layers of the drumhead, when it becomes much more difficult to distinguish this form of inflammation from acute otitis media.

Symptoms. Pain, usually moderate in character, is complained of, although it may be sharp and stinging in severe cases. Other symptoms are a pulsating noise and more or less tinnitus. The disturbance in hearing is usually much less than in cases of acute otitis media, and one is justified in making a diagnosis of acute myringitis when, in spite of more or less swelling or bulging of the membrane, the hearing-distance is not materially affected. Frequently, however, the two diseases are associated, or a case beginning as an inflam-

mation of the drumhead is followed by acute otitis media, so that in many instances it becomes difficult to separate the two. The duration of the disease is usually four or five days. After the vesicles have burst, the membrana assumes a macerated appearance and the dermal layer peels off, leaving a congested drumhead. In the chronic form there is more or less suppuration, with the formation of granulations or ulcerations on the drumhead if the disease has run a protracted course.

Treatment. In severe cases the artificial leech should be applied, while in the mild form of this inflammation the vesicles generally burst and recovery follows in a few days. The insufflation of boric acid powder is indicated in the latter cases. In the chronic form boric acid alone or combined with zinc oxide (equal parts) should be insufflated, or drops should be instilled, containing zinc sulphate, copper sulphate, or lead acetate, in the proportion of four grains to the ounce. Granulations should be cauterized with nitrate of silver fused on the end of a probe.

INJURIES OF THE MEMBRANA TYMPANI.

Injuries of the drumhead are usually caused (1) by the introduction of a foreign body, (2) by a blow or explosion producing a sudden condensation of air in the external meatus or tympanum, or (3) by a fracture of the cranial bones, thus involving the drumhead.

Diving, sea-bathing, boxing the ears, the discharge of cannon, and gunshot wounds are frequent causes of rupture; also violent paroxysms of coughing, vomiting, and sneezing. Again, ruptures are often met with in patients who have tried to remove wax from the ears with toothpicks and hair-pins, and have pushed them in too

far, so that these and other unskilful attempts at the removal of foreign bodies often cause serious trouble. Rupture of the drumhead has been known to follow a too violent inflation of the ear by Politzer's method, and also the sudden exhaustion of the air from the external meatus by the use of Siegle's otoscope. In the cases where rupture occurs from the sudden condensation or rarefaction of air, it will generally be found on examination that catarrhal changes have taken place in the membrana tympani, and that there is more or less obstruction to the entrance of air through the Eustachian tube.

In direct injuries the point of rupture is most frequently observed in the anterior half of the drumhead, and the size and shape of the rupture or perforation depend to a great extent on the nature of the instrument used. A sharp-pointed instrument is apt to cause a wound somewhat round (see Plate V., Fig. B), the edges being covered with dark-red blood, and frequently there is more or less blood in the canal, with laceration of the meatus. Inflammation of the middle ear, followed by suppuration, may result from direct injuries. The shape of the rupture produced by indirect violence is, as a rule, gaping, so that the inner wall of the tympanum will be seen. (See Plate V., Fig. C.) This wall will appear reddish yellow in color, in direct contrast to the bright-red hue observed in cases of acute or chronic otitis media. The most frequent seat of the rupture is in the posterior half of the membrana, and the shape of the perforation is usually round or oval, with the long axis of the oval running parallel with the direction of the radiating fibres. There is, as a general rule, but one rupture. When the rupture is due to

a fall or blow, there may occur at the same time concussion of the labyrinth, with marked deafness, and it is in such cases, as well as in those in which suppurative inflammation of the middle ear has developed, that permanent disturbances of hearing frequently remain after the perforation has healed.

Beside the causes already mentioned, a case[1] has been reported of injury to the membrana tympani from a twig. The patient was walking through the woods when a twig of a branch penetrated the right auditory canal, rupturing the drumhead. In another patient the injury to the ear was due to a pistol-shot wound. The most serious cases are those in which there is concussion of the labyrinth from a fall or blow. The prognosis in such instances must always be grave, and on account of their medico-legal significance they become of great interest to the aural surgeon, who is frequently asked to give expert testimony as to the diagnosis and prognosis of such accidents. In order that the surgeon may express an opinion as to whether the rupture is of traumatic origin, it is absolutely essential that the drumhead be inspected immediately after the alleged injury. Just after an injury to the drumhead from a sharp-pointed instrument, as previously mentioned, the edges will be covered with dark-red coagulated blood; and if the rupture is large, the middle-ear wall will appear yellowish red. If suppuration has already occurred, it is impossible to make a diagnosis of traumatic rupture. Usually when the perforation has been due to disease of the middle ear, even when of large size, a sharp, hissing sound is heard when air is forced through the Eustachian tube by the Valsalvian or Politzer method; while in

[1] Bacon: Transactions of American Otological Society, 1888.

a normal ear which has been ruptured through traumatism, the escaping air causes a broad, deep-breathing sound. Injuries to the drumhead from sharp instruments heal rapidly, as a rule, if inflammation of the middle ear does not follow.

If suppurative inflammation occurs, the disease may become chronic and lead to complications, or the drumhead may be thickened and adhesions take place, or there may be concussion of the labyrinth with or without rupture of the drumhead. In all such cases permanent disturbances of hearing may remain. In cases in which deafness results from a blow or fall, and the drumhead and external meatus appear normal, while there is loss of bone-conduction, as shown by the tuning-fork, the surgeon should be very guarded in expressing an opinion as to the cause of the deafness, for in such instances much depends on the answers given by the patient, who may be a malingerer. Tuning-fork tests show that the bone-conduction is much diminished or even lost in labyrinthine disease. while the patient is apt to complain of subjective noises, giddiness, deafness, and a dull feeling in the head.

A case[1] has been reported in which a suit was brought for damages on account of an alleged blow on the ear. In this instance the boy was a catarrhal subject, and the hearing was as much affected on one side as on the other.

Treatment. It is most important that the instillation of drops and syringing the ear should carefully be avoided in all cases of simple rupture. The patient should be kept quiet and placed on a low diet for a few days, and a piece of absorbent cotton should be inserted in the meatus. When inflammation of the membrana

[1] Bacon: New York Medical Journal, December 12, 1888.

occurs, the artificial leech should be applied in front of the tragus (see Treatment of Acute Otitis Media); if suppuration ensues, the same treatment should be followed as that indicated for acute suppurative otitis media. When concussion of the labyrinth has occurred, permanent disturbance of hearing and tinnitus will occasionally remain in spite of all treatment. In some cases, nerve tonics containing strychnine sulphate are indicated. Others improve under the administration of iodide of potash and hypodermic injections of pilocarpine. The constant electric current is beneficial in other instances.

A rare form of injury to the drumhead is *fracture of the handle of the malleus.* The lesion may be due to a fall or to a direct injury to the membrana tympani. It is possible to make a diagnosis of this condition if the surgeon can discover a false point of motion in the long handle while the patient inflates the ear by the Valsalvian method.

OTITIS MEDIA ACUTA, OR ACUTE INFLAMMATION OF THE MIDDLE EAR.

Acute inflammation of the middle ear may be either catarrhal or purulent. Very frequently, however, one disease runs into the other, so that it is extremely difficult in some cases to make any distinction.

OTITIS MEDIA CATARRHALIS ACUTA, OR ACUTE CATARRHAL INFLAMMATION OF THE MIDDLE EAR.

There are in the first stage considerable hyperæmia and swelling of the lining membrane of the tympanic cavity and of that covering the ossicles and inner surface of the drumhead. There is probably a more or less

hyperæmic condition of the lining membrane of the mastoid cells in many, if not all, cases of this disease. After the acute stage there follows more or less effusion of serum containing exudation cells. In severe cases there may be ecchymoses. The epithelium lining the tympanum peels off and becomes macerated. Thick mucus, pus-cells, and numerous red blood-corpuscles are found in the tympanic cavity. In some instances the mucus is very tenacious and stringy, while in others there is a preponderance of pus-cells. This affection is what is commonly called "earache" in children, and some children are especially liable to frequent attacks. It is more often observed during the spring and winter months, when sudden changes in temperature are apt to occur.

Causation. This disease most frequently follows a pharyngitis or "cold in the head," or it may be due to long exposure to wet or cold; or it may be a complication of the acute infectious diseases, such as measles, scarlatina, whooping-cough or pneumonia, bronchitis, tuberculosis, puerperal fever, cerebrospinal meningitis and syphilis. Another cause is the use of the nasal douche, by means of which water entering the Eustachean tube sets up an acute inflammation. In many instances the same accident follows the pernicious practice of "sniffing up" salt and water for the relief of nasal catarrh.

Teething children are liable to attacks of catarrhal otitis media, which frequently develops into the purulent form with perforation of the drumhead.

Sea-bathing is one of the causes of this disease as well as nasopharyngeal catarrh, or it may be due to the presence of adenoid growths and enlarged tonsils. In sea-

bathing there is a liability that water filling the external meatus may cause an inflammation of the canal or middle ear, while, on the other hand, an acute otitis media may result from water entering the middle ear through the Eustachian tube. Parents should forbid children to "duck" the head or to dive while bathing, and should always insist that they wear in the ear a piece of non-absorbent cotton-wool.

Symptoms. There is a dull feeling or fulness about the ear, or the pain may be sharp and stinging in character and quite severe at night. Infants are liable to scream and to put the hand to the side of the head. Headache is sometimes complained of. The pain frequently radiates about the ear and side of the head, and shoots toward the teeth, and the patient may complain of pain on opening the mouth and when pressure is made just below the lobule. In children, more especially, this form of disease is accompanied by fever, while in rare instances delirium and convulsions are observed. In infants meningitis is sometimes suspected until an examination of the ear reveals the cause of the disease. The pain, though much less severe in acute catarrhal than in acute purulent otitis media, is due to the severity of the inflammation itself, to the pressure exerted on the drumhead by the exudation, or to great retraction of the drumhead.

Constriction of the Eustachian tube interferes with the proper ventilation of the tympanic cavity; owing to absorption of the air in the tympanum having taken place, the drumhead becomes retracted from the atmospheric pressure and the patient complains of pain.

In the milder cases the pain is of short duration, but in the more severe ones it is apt to continue for several

days, while during convalescence the patient has a feeling of fulness and pressure in the ear.

Tinnitus, or noise in the ear, of a hissing, rushing, singing, or pulsating character, is a source of considerable annoyance to the patient. The deafness in the early stage as well as the tinnitus is due to the intense hyperæmia and to inflammatory exudation. The disturbance in hearing is apt to be more marked during the stage of exudation, but the degree of deafness does not depend so much on the amount as on the seat of the exudation in the middle ear. If both fenestræ in the middle ear are clogged, there is apt to be great deafness. A patient frequently complains of autophony—*i. e.*, a muffled sound of his own voice, which is likened to the hollow sound heard when talking with the head in a barrel. Another source of annoyance is the hearing by the patient of the same note double (diplacusis)—that is, a tuning-fork when vibrating and held close to each ear will seem a tone or half a tone higher usually on the affected side. This same phenomenon also occurs in labyrinthine disease and in some cases of exostosis.

In children the pain is especially apt to come on during the night, while in the daytime it is not generally so severe. Parents very often attach but little importance to such attacks, and consider that it is not necessary to seek professional advice for a mere earache. Repeated attacks, however, frequently lead to changes in the drumhead and adhesions, and later in life to deafness. There would be far fewer cases of incurable chronic middle-ear catarrh if these children during early life had received proper treatment. The duration of the disease varies from a few days to a week or ten days. There may be several relapses. Much depends

PLATE V.



on the general condition of the patient and on the causation.

Appearances of the Membrana Tympani. In less severe cases the drumhead appears congested, especially about Shrapnell's membrane and the long handle of the malleus. (See Plate V., Fig. D.) The bony canal may or may not be involved. In severe cases the whole membrane is of a reddish, angry-looking color, and frequently has a gray appearance on account of the infiltration of the dermal layer. The only landmark, the short process, appears as a red or yellowish-white point. The posterior superior quadrant of the membrane may present a bulging appearance due to inflammatory infiltration. In some cases there may be vesicles and interlamellar abscesses. Bulging and exudation sacs in connection with the tympanic cavity are occasionally observed. In other cases there may be a bulging outward of Shrapnell's membrane or of the posterior inferior quadrant of the membrana vibrans, due to the

---

DESCRIPTION OF PLATE V.

A.—Normal membrana tympani. Pages 40–77.

B.—Injury to drumhead from a sharp-pointed instrument. Congestion in Shrapnell's membrane and along the handle of the malleus. Page 150.

C.—Rupture of drumhead due to indirect violence. Page 150.

D.—Acute catarrhal otitis media. Hyperæmic appearance of the membrana tympani in the early stage. Page 157.

E.—Sero-mucous catarrh of the middle ear. Fluid in tympanum seen through transparent membrana tympani. Page 162.

F.—Acute purulent otitis media. Deeply congested membrana tympani. Perforation in the lower anterior quadrant. Page 168.

G.—Acute catarrhal or purulent otitis media. Stage of resolution. Membrana tympani gradually assuming a grayish appearance. Tortuous vessels extending from the periphery to the handle of the malleus. Pages 158 and 170.

H.—Chronic catarrhal otitis media. Great retraction of drumhead, with a chalky deposit, crescentic in shape, in posterior portion. Page 226.

K.—Chronic purulent otitis media. Large perforation. Handle of malleus, resting against the inner tympanic wall. Page 257.

presence of fluid in the tympanum. The congestion gradually disappears when resolution begins, and the membrana at first assumes a grayish appearance, with small, tortuous vessels running from the periphery toward the handle of the malleus. (See Plate V., Fig. G.) The epidermis peels off from the membrana and bony canal, while the congestion remains longest along the handle of the malleus and in Shrapnell's membrane.

Treatment. Acute catarrhal otitis media in children is often due to the presence of enlarged tonsils or of adenoids, so that it is of the greatest importance to make a thorough examination of the nasopharynx in all cases. Usually in young subjects it is impossible to obtain a satisfactory view of the nasopharynx with the rhinoscopic mirror, a fact which compels the otologist to make a digital examination of this region. With a rubber band on the index finger, to prevent it from being bitten, an examination can readily be made; and, further, while the finger is in the nasopharynx, if adenoid growths are present, and particularly if they are soft and flabby, the surgeon can scrape away some of these tumors with the finger-nail and break them up, so that something can be accomplished at this time. There are many little patients, however, of a nervous temperament whom it is necessary to anæsthetize even when making an examination. In such cases chloroform or nitrous oxide gas is to be preferred. (For the further treatment of adenoids, the reader is referred to the chapter on Nasopharyngeal Diseases.) In the first or hyperæmic stage of acute catarrhal otitis media the local abstraction of blood by means of the artificial leech is a valuable method of treatment, and if applied early will often

cut short the attack and relieve the pain. The leech should be applied in front of the tragus. (See method of application under Treatment of Acute Purulent Otitis Media.) In children warm water gently poured into the ear will often stop the pain, or the ear may be douched with warm water (containing boric acid, ʒj to Oj) every two or three hours. Some surgeons prefer dry heat applied by means of a hot-water bag or a hot bag of salt. In the mild cases the instillation of a saturated boric acid solution will be all that is necessary if careful attention is paid to the condition of the pharynx and nasopharynx.

According to Kirchner, 10–20 per cent. carbolic glycerine warmed and dropped into the ear may in two to three applications cut short the inflammation. When a perforation has occurred this concentrated solution will cause severe pain. To abort the inflammation he recommends most highly Burow's solution (Liq. alumin-acetat. 12 parts, plumbi acetat. 5 parts, and water 100 parts.). The ear should be filled with this solution warmed three to four times a day and it should be allowed to remain in the canal twenty to thirty minutes.

In children when the Eustachian tube is swollen and the drumhead is very much retracted, a condition of affairs resulting from a nasopharyngeal catarrh, (tubal catarrh), the treatment should consist in inflation of the middle ear by Politzer's method or catheterization, and the application of astringent sprays to the nasopharynx.

After the inflammatory stage has passed, the catheter or Politzer's method should be employed, in order to prevent adhesions, to restore the membrana tympani to its normal position, and to remove any mucus or other fluid

from the tympanum. If the disease follows a cold, small doses of calomel should be given, to produce slight catharsis, and aconite should be administered in small and frequently repeated doses. The patient should be kept quiet in the house and given a light diet. Gargles are generally indicated, and attention should be paid to the nares as well as to the naso-pharynx. Instillations of a 4 per cent. solution of cocaine, or of a solution containing liq. Magendie, ʒj; aqua, ℥j, are indicated in somes cases. It is always best not to administer opium unless absolutely necessary. Patients who frequently have "colds" should be cautioned against taking large doses of quinine, as deafness is undoubtedly caused by this practice in some instances. Catarrhal subjects should be impressed with the importance of paying strict attention to the general health. They should lead an out-of-door life as much as possible, protect their feet in stormy weather, and should sponge the neck and chest every morning with cold water, and follow this by vigorous rubbing with a coarse towel. Living in overheated rooms is a frequent source of catarrhal disease. If there is bulging of the membrane, suggesting the presence of fluid in the tympanic cavity, the drumhead should be incised immediately. (See Treatment for Acute Purulent Otitis Media).

SUBACUTE CATARRHAL OTITIS MEDIA.

This is a form of inflammation similar to acute catarrh, except that it is milder in character, the pain and inflammation being less marked. The patient is usually subject to frequent attacks, while the tonsils are apt to be en-

larged, and the disease is liable to become chronic. This disease is often found among school-children, who formerly were thought to be inattentive. Proper hygienic surroundings, tonics, etc., should be prescribed, and treatment directed to the nares and postnares, in addition to inflation of the middle ear by Politzer's method.

What is commonly called *tubal catarrh* is an inflammation of the Eustachian tube, leading to great retraction of the drumhead. The membrana is usually but slightly congested. As the swelling of the walls of the tube allows but little air to enter the tympanic cavity, the air in the middle ear becomes absorbed and retraction of the membrana follows. This affection is usually secondary to nasopharyngeal catarrh.

Seromucous catarrh, or *hydrops ex vacuo*, is in reality another form of acute catarrhal otitis media. In this affection the faucial end of the Eustachian tube is usually congested and swollen, leading to closure of the tube. Acute inflammation of the nasopharynx is generally the cause of the disease. The drumhead may be slightly congested as well as retracted, and other changes of color will be noticed, depending on the amount of the exudation, the condition of the middle ear, and the transparency of the drumhead. When the tympanic membrane is translucent and there is a seromucous exudation, a line of demarcation between that portion of the tympanum containing air and that containing fluid will be distinctly seen. The latter portion usually appears greenish yellow in color and the former darkish gray. (See Plate V., Fig. E.) When the patient moves his head the line of the fluid will change if it is serum. After inflation of the middle ear, air-bubbles, as rings,

will be seen through the membrane if the latter is transparent.

Symptoms. The symptoms generally observed are very slight pain, a sense of fulness or pressure in the ear, tinnitus of an intermittent character, autophony, and a cracking sound in the ear, which latter is probably due to a sudden separation of the walls of the Eustachian tube, and more or less deafness. Tests with the tuning-fork show increased bone-conduction on the affected side unless the labyrinth is involved.

Prognosis. The prognosis is generally favorable when the inflammation is due to a cold and the patient's general health is good. The disease in some instances becomes chronic, and may lead to adhesions and deafness.

Treatment. When there is fluid in the middle ear, it may sometimes be made to disappear by Politzer's method of inflation or by catheterization. A much more satisfactory method, however, is to puncture the drumhead in its posterior inferior quadrant, and then force the fluid out by inflation or by the use of Siegle's otoscope. Careful attention should be paid to the patient's general health, and treatment directed to the nasopharynx, the nasal passages, and the mouth of the Eustachian tube. Catarrhal patients should be advised to follow the same line of treatment as that recommended in the chapter on Acute Catarrhal Otitis Media. Preparations containing iron and strychnine should be given. Whenever possible, catheterization is preferable to Politzer's inflation. In children, however, and in adults of a nervous temperament, and also in all cases in which there is much inflammation and irritation about the faucial end of the Eustachian tube, Politzer's method should be employed. As to the frequency of inflating

the ears, a good rule is to inflate the ear at first every day for several days, then every other day, and less frequently as the condition requires. A method advocated by Politzer, of removing the fluid from the tympanum, is as follows: After the patient has taken a little water in his mouth he is directed to turn his head well forward and somewhat toward the opposite side, a position which causes the pharyngeal orifice of the Eustachian tube to point directly downward. The patient should keep his head in this position for a minute or two, in order to allow the secretion to flow toward the orifice. Air is then forced into the middle ear by Politzer's method. In some instances in which the mucus in the middle ear is tough and stringy, it is necessary, in order to remove the secretion, to make a free incision in the drumhead, and then inject into the tympanic cavity a very weak solution of bicarbonate of soda.

CHAPTER VI.

OTITIS MEDIA PURULENTA ACUTA, OR ACUTE PURULENT OTITIS MEDIA.

Acute purulent otitis media differs from acute catarrhal otitis media in the greater intensity of the stage of inflammation, and the occurrence of a purulent discharge with perforation of the membrana tympani. In the early stage of the disease there is marked congestion of the tympanic cavity and tympanic end of the Eustachian tube. If the inflammation has extended from the nasopharynx, which is usually the case, the Eustachian tube in its whole extent is more or less congested and swollen. It is probable that the lining membrane of the mastoid cells is likewise congested at the same time in many, if not all, cases of acute purulent otitis media. In some instances there is hyperæmia of the labyrinth with serous exudation. The inflammatory stage is followed by considerable swelling of the lining membrane of the middle ear. The discharge is frequently of a serous character at first, or it may be mucopurulent. Later, however, it becomes decidedly purulent. It may be scanty or very profuse. Acute suppurative otitis media frequently follows the acute catarrhal variety, so that, clinically, the two diseases should be considered as different stages of the same disease, viz., acute otitis media. In cases of tuberculosis there is, as a rule, but little hyperæmia, the mucous

membrane of the middle ear appearing of a pale, reddish-yellow color, while the discharge or exudation is apt to be thin and watery.

Occasionally the discharge may be acrid and give rise to severe and extensive excoriation about the canal and pinna in a very short time, sometimes within two or three days but fortunately this does not occur frequently in acute cases, and generally in ill-kept and poorly nourished children.

In cases due to *influenza*—and these constitute by far the greatest number—the discharge is at first serosanguinolent and very sticky in consistency. In some cases there is more or less blood, and the stage of inflammation is marked by great hyperæmia and swelling of the mucous membrane. These symptoms are so constant that they are very characteristic of *influenza otitis*, while the mastoid cells are very often involved at the same time, and in rare instances the labyrinth is affected.

Etiology. Purulent inflammation of the middle ear usually occurs in connection with acute or chronic nasopharyngeal catarrh, and may be due to the same causes that produce acute catarrhal otitis media. Beside the influenza and colds, which are responsible for a great many cases, the exanthematous diseases play a most important rôle in the causation, especially measles and scarlet fever. Other causes are diphtheria, exposure to wet and cold, smallpox, typhus and typhoid fevers, tuberculosis, pneumonia, bronchitis, whooping-cough, syphilis, erysipelas, diabetes, and the puerperal state. Acute suppuration of the middle ear is due at times to operations on the drumhead, such as paracentesis, to blows on the head or ear, or to falls, to the improper

use of instruments when attempts are made to extract a foreign body from the external meatus, and to the use of caustic applications, and occasionally it follows operations on the nose. Water occasionally enters the Eustachian tube and middle ear from the nasal douche, and from "sniffing up" salt water and other solutions for the relief of nasal catarrh, a most pernicious practice, and causes severe inflammation of the middle ear and occasionally of the mastoid cells. During dentition there is apt very frequently to be a purulent discharge, and it is most important in such cases to examine frequently the ears during this period. *Sea-bathing* is another cause of this disease. The drumhead may be ruptured in diving or from the force of a wave striking against the side of the head ; or the cold water filling the external meatus may set up an inflammation of the drumhead and later of the middle ear. Frequently, however, I believe that an earache from sea-bathing is due to the fact that water enters the middle ear through the pharynx and Eustachian tube, the result frequently of forcibly blowing the nose after diving, the uniform act of every bather.

Children are more frequently subjects of this disease than adults, and it is observed more often in winter and spring, especially during sudden changes of weather. Children are apt to have both ears affected from scarlatina, measles, or diphtheria.

Appearances of the Membrana Tympani. In the first stage of inflammation the drumhead in the region of Shrapnell's membrane is congested, and the hyperæmia extends along the posterior border of the handle of the malleus. The drumhead becomes gradually pinkish in its entire extent, or of a bluish-red and angry appearance,

according to the intensity of the inflammation. Ecchymoses or vesicles filled with serum or blood are sometimes observed on the membrana tympani or on the walls of the external meatus. In cases of intense inflammation marked congestion of the bony auditory canal occurs, so that it is difficult to distinguish where the bony wall ends and the drumhead begins. Exudation of serum with swelling of the tissues gives the drumhead a sodden and lustreless appearance. When the epidermis peels off, the drumhead may be mistaken for a polypus, on account of its smooth, glistening, and red appearance. Bulging of the membrana tympani is observed most frequently in its posterior half, and this may be due to swelling of the tissues or to interlamellar abscesses, or, more frequently, to a collection of fluid in the tympanum. The most frequent seat of the perforation is in the lower anterior or posterior quadrant of the drumhead. It is occasionally, though very infrequently, found in Shrapnell's membrane. When the latter does occur, the disease is mostly confined to the attic, exudation of fluid being found about the articulation of the malleus and incus, and in Prussak's space. The appearances of the drumhead in such cases are intense redness in its upper portion, in Shrapnell's membrane, and in the wall of the canal adjacent to it, followed by bulging of the membrana flaccida. If the tissues are not incised at an early date, perforation is apt to result and pus is discharged. The perforation in such cases may heal and the hearing be restored, or adhesions between the ossicles and walls of the tympanum may form and give rise to considerable disturbance of hearing, or the disease may become chronic, when caries of the ossicles is the usual result. In *influenza cases* the

drumhead is markedly injected and angry looking, as a rule, and in some cases the perforation has a nipple-shaped projection in the posterior half of the drumhead, out of which the pus escapes with difficulty. Marked pulsation of the fluid will be noticed at this point. In some cases not only the bony canal, but the cartilaginous portion as well is inflamed. Especially is this the case when otitis media results from injuries to the canal and drumhead from the improper use of instruments or caustic applications. At times the lymphatic glands behind the ear are enlarged. After the discharge has filled the canal the landmarks are not easily made out, so that it becomes necessary to remove the secretion by syringing and by using absorbent cotton on the end of a cotton-holder before the site of the perforation can be discovered. Cases in which the discharge consists largely of stringy mucus are, as a rule, extremely difficult ones to cure.

Symptoms. The disease usually commences with pain of an excruciating and lancinating character in the ear, not limited, however, to the ear, but shooting in the direction of the teeth, the neck, and in the occipital, parietal, or temporal region. The pain is more intense at night than during the daytime. As previously mentioned, perforation may occur without pain in the case of tuberculous subjects. In some persons the pain is not very marked, but such cases are infrequent. The temperature is usually considerably elevated in the case of young children—as high as 105° F. In adults the temperature varies very much, usually from 100° to 102° F. After perforation has occurred the pain, as a rule, becomes much less severe, and occasionally ceases at once.

In young subjects, meningitis sometimes is suspected on account of a very high temperature, and from the fact that the child puts its hand to the side of the head. In some children the disease is ushered in by a convulsion or severe vomiting. A child that has been screaming with pain, and that has been on the verge of a convulsion, will suddenly fall asleep, and the temperature will drop several degrees, as soon as perforation has taken place.

Other symptoms complained of are tinnitus, which is described by the patient as being of a roaring, hissing, hammering, or pulsating character, such noises probably being due to secretion covering the ossicula and fenestræ ovalis and rotunda, or to a hyperæmic condition of the labyrinth, with possibly exudation; giddiness and fulness in the head, and deafness, which may be very slight or well marked, depending on the intensity of the disease, the nature of the secretion, and especially as to whether the labyrinth is implicated or not. After perforation the hearing-distance usually is improved. Patients generally hear the watch or acoumeter through the cranial bones. In severe cases of scarlet fever or diphtheria in which the labyrinth is involved or in which acute caries follows, or in cases of syphilis, this power of hearing the watch or acoumeter will be lost. In Weber's test the vibrations of the tuning-fork usually are perceived better by the affected ear.

Course and Complications. Perforation may occur in a few hours, or it may be three or four days or longer before the discharge makes its appearance. Even after perforation has taken place, the pain at times is severe. In such cases it is probable that the periosteal layer is involved or that the mastoid cells are implicated.

As a rule, after perforation, all symptoms, such as pain, fulness in the head, and giddiness, are considerably modified. The duration of the suppurative stage varies from several days to three weeks, or the disease may become chronic. A cicatrix may or may not mark the seat of the perforation. During the stage of resolution the membrana gradually loses its deep-red color and takes on a grayish-red appearance; the landmarks gradually reappear, viz., the handle of the malleus and the short process. (See Plate V., Fig. G.) In some cases adhesions form between the ossicula and the tympanic walls, and disturbances of hearing result. Opacities, calcareous deposits, atrophic changes, and thickening of the drumhead are observed in other instances.

Acute purulent otitis media is apt to follow a much more protracted and severe course in those who are debilitated or scrofulous, or who are suffering from one of the exanthematous diseases, diphtheria, bronchitis, etc., than when it occurs in a patient who otherwise is healthy and strong. In cases occurring during the course of scarlatina, diphtheria, measles, and influenza, the inflammation is frequently of an intense and virulent nature, and unless a free incision is made at an early date in the drumhead caries of the ossicles and tympanic walls is the result. To emphasize this point, I would refer to a case of scarlet fever that came under my care a few years ago, and in which acute suppurative inflammation of the right ear had already taken place at the time of my first visit. The left ear subsequently became inflamed and painful: a free incision was made in the drumhead, with the result that the left membrana tympani subsequently healed and the hearing on this side became normal, while in the right

ear, in which the perforation had taken place spontaneously, the greater portion of the drumhead was destroyed.

Permanent disturbances of hearing occur in cases in which the ossicula are bound down by adhesions, or in which there is an adhesion between the drumhead and middle-ear walls, or in which thickening or atrophic changes or calcareous deposits are observed in the drumhead. Deafness is often well marked when great destruction of the drumhead and ossicula takes place, and especially when the labyrinth is affected.

Beside caries of the ossicula and walls of the tympanic cavity, as a result of acute suppurative otitis media, other complications that occasionally follow are acute mastoiditis, facial erysipelas, meningitis, pyæmia and septicæmia, thrombosis of the lateral or other sinuses, hemorrhage from the internal jugular vein or carotid artery due to acute caries of the temporal bone, and epidural and cerebral abscesses. I have reported two cases in which erysipelas developed suddenly as a complication of acute purulent otitis media, the disease involving the scalp and face and giving rise to considerable constitutional disturbance. When erysipelas first makes its appearance the discharge from the middle ear becomes much less in quantity or ceases altogether. Erysipelas is undoubtedly due to bacterial invasion of a wound by Fehleisen's streptococcus erysipelatis. In both cases referred to, the auricle first was affected. I have reported also a case of pyæmia occurring in connection with acute purulent otitis media in which recovery ultimately took place.

That teething is a cause of acute purulent otitis media was clearly demonstrated in the report of a case[1] of a

[1] Annals of Ophthal. and Otol., January, 1894, vol. iii. No. 1.

child who came under my observation, and who had very high temperature and symptoms of brain complication. The mastoid antrum was carious, and it was necessary to scrape it with sharp spoons. The disease was undoubtedly due to sympathetic irritation from the cutting of four molar teeth; the child recovered. When pain persists after the appearance of the discharge, and especially if the temperature is high, such indications generally point to mastoid disease or some other complication.

Diagnosis. Before perforation has occurred, it is impossible to make a differential diagnosis between acute catarrhal and acute purulent otitis media, as in both diseases the first stage is marked by hyperæmia of the drumhead. As previously mentioned, the hyperæmia is usually more marked and the pain more intense in acute suppurative otitis media. Acute myringitis may be mistaken for acute otitis media. In the former, however, the pain is not so severe, nor is the disturbance of hearing so great as in the latter disease. If there is stringy mucus in the discharge from the ear, one may safely diagnose acute purulent otitis media, as mucus is never present in a case of otitis externa or furuncle of the canal. After removal of the discharge by syringing or by means of absorbent cotton on a cotton-holder, inspection with head-mirror and speculum will usually reveal the seat of the perforation.

In cases in which the latter cannot be seen on account of the curvature of the external auditory canal, a perforation whistle may often be heard by having the patient try the Valsalvian method of inflating the ear, or by Politzer's method. This experiment will not suc-

ceed, however, when there is great swelling of the Eustachian tube or when the perforation is in Shrapnell's membrane. It should not be forgotten that in some cases of acute purulent otitis media the discharge is

FIG. 53.

Case of carcinoma of the temporal bone, giving rise to facial paralysis.

composed almost wholly of pus. Malignant disease having its origin in the temporal bone may give rise to great difficulty in the diagnosis during the early stage. I have reported the case[1] of a woman (see Fig. 53) who

[1] Archives of Otology, 1888, vol. xvii. No. 1.

complained for some time of pain with tinnitus in the right ear, followed by a slight serous discharge. Facial paralysis caused her considerable alarm, so that she sought advice. There was marked induration of the tissues in front of the ear, extending toward the jaw. The pain became very severe, as is usually the case in such patients, an important point to consider,

FIG. 54.



a. Trabeculæ. *b*. Mastoid sinuses. *c*. Carcinoma.

for when it is persistent and cannot otherwise be accounted for, malignant disease should be suspected. This case proved to be a carcinoma, as shown in Figs. 54 and 55. Another case with a similar history also came under my care.

Prognosis. In simple cases of acute purulent otitis media occurring in healthy individuals when due to

cold or nasopharyngeal catarrh, the prognosis is good if the case is treated properly. The prognosis is, however, unfavorable in cachectic, scrofulous, tuberculous and syphilitic patients; also in severe cases of diph-

FIG. 55.



a. Mastoid sinuses. *b.* Carcinoma. *c.* Bone trabeculæ.

theria, scarlet fever, measles and typhus fever. In some cases of influenza a favorable prognosis should not be given. One should consider the following as unfavor-

able symptoms, viz., a distressing tinnitus, fever; severe pain, which has been more or less continuous for several weeks; a profuse discharge, which has persisted in spite of treatment; hemorrhage from the ear, caused in most instances by granulations or polypi; great destruction of the membrana tympani; symptoms of mastoid disease; swelling of the bony auditory canal; caries of the

FIG. 56.



Malignant growth of temporal bone.

ossicula and walls of the middle ear; facial paralysis, and enlargement of the lymphatic glands in the neck. Facial paralysis sometimes, although very infrequently, manifests itself in simple acute otitis media even when perforation does not occur, but in such instances even the paralysis generally disappears in a few days.

Treatment. In uncomplicated cases the patient should be kept quiet in the house, while in severe cases he

should be put to bed and given a light diet. A trit-urate tablet of calomel, gr. one-tenth to one-quarter, should be administered every hour for six doses, or until a slight laxative action is produced. Aconite in drop doses is also a most valuable remedy when there is fever, and especially in cases due to cold. Tincture of pulsatilla, likewise given in drop doses, is indicated also in cases in which there is a profuse discharge from the nares or nasopharynx, and may be administered alternately with aconite. In the early stage, when there is great hyperæmia, the artificial leech should be applied at once in front of the tragus and close to it. When the mastoid is simultaneously involved, it should be placed

Fig. 57.

Author's scarificator.

behind the auricle just over the antrum or the mastoid tip. By such treatment the disease will, in simple, uncomplicated cases, frequently be cut short, or it will at least run a much less severe course. The artificial leech consists of a scarificator (Fig. 57) and cupping-glass (Fig. 58). The scarificator has seven lance-shaped needles, and the depth of the punctures made in the soft tissues can be regulated by screwing or unscrewing the cap on the end of the instrument. The latter being placed in front of the tragus or over the mastoid, according to the indications, the spring is then pulled back and the tissues scarified. The cupping-glass is immersed in hot water for a few seconds and then

applied over the bleeding surface. The artificial leech is much to be preferred to the natural one, for the following reasons: 1. The scarificator and cupping-glass are always at hand, while natural leeches are frequently very difficult to obtain, especially at night. 2. Leeches are very repulsive and disagreeable to most patients, and

Fig. 58.

Author's cupping-glass.

especially to children. 3. After the artificial leech has been removed, the bleeding ceases at once, while with the natural leech it is often difficult to control the hemorrhage. I have known of cases in which alarming hemorrhage occurred after the surgeon left the patient.

In the case of children, hot water instilled into the

ear often affords great relief, as well as douching the ear every hour or two with a warm boric acid solution. Some patients derive much comfort from the application of dry heat by means of a hot-water bag or hot bag of salt. Lucæ uses hot compresses about the ear soaked in a solution of acetate of aluminium, two drachms to the pint of water. Drops of a 4 per cent. cocaine solution may be used with benefit, as well as a solution composed of—

℞. Atropiæ sulphat.,	gr. ij;
Acid. boric.,	gr. x;
Glycerin.,	ʒj;
Aquæ,	℥j.

All drops and solutions for the ear should always be warmed.

The condition of the pharynx, nasopharynx, and nares should be noted, and treatment directed to the parts whenever necessary. (See Chapter VII.)

As micro-organisms travel from the pharynx and nasopharynx to the middle ear by means of the Eustachian tube, it is of the greatest importance in cases of measles, scarlet fever, diphtheria, and influenza, to try to prevent the disease from extending to the middle ear. To accomplish this, sprays of boric acid or carbolic acid or bichloride of mercury (1 : 5000) are indispensable, while the ears should be examined frequently for the first evidences of inflammation. As the micro-organisms in these diseases are apt to cause acute caries, it is very important that the surgeon should make an *early incision* in the drumhead, and not wait until perforation has occurred through ulceration. A *free incision* should be made with a bistoury

Fig. 59.

Author's knife for incising the membrane tympani and for dividing adhesions.

(see Fig. 59), and *not a paracentesis, as formerly was practised,* the line of incision extending from just behind the stapes to the lower border of the drumhead and close to the bony auditory canal (see Fig. 60). In some cases, particularly where there is bulging of the membrana tympani in its upper and posterior quadrant, complicated with symptoms of acute mastoiditis, the knife should be carried well up to the posterior fold of the drumhead and into the attic, so as to drain more fully that cavity, as well as the aditus ad antrum. The knife should then be brought out along the upper posterior wall of the external auditory canal, so as to relieve the tension frequently observed in this locality. Before incising the drumhead, the auditory canal should be syringed with a bichloride solution (1–5000), and thoroughly dried with sterile absorbent cotton. As incision of the drumhead is an extremely painful operation, it is generally advisable to administer an anæsthetic. In young children a whiff of

Fig. 60.

Line of incision.

chloroform may be given, but great care should always be used in its administration. Nitrous oxide gas is to be preferred in the case of adults. A convenient apparatus for this purpose has been manufactured by the S. S. White Dental Manufacturing Company, which can be conveniently carried by the surgeon. Ethyl chloride has also been used lately in an inhaler especially designed for its administration. After the membrana has been incised, the object of treatment should be, first of all, to promote the discharge as much as possible by douching the ear with the normal salt solution or a solution of boric acid, ʒj to water ʒviij, or a 1–2 per cent. solution of aluminium acetat., (Kirchner); and secondly, to try to destroy the micro-organisms. A bichloride solution (1–5000) can be used to advantage after the discharge has become profuse, and especially in cases in which the infection is of a virulent character. After douching the ear, which in some cases is necessary every hour or two, according to the amount and character of the discharge, it is advisable to insert a strip of boric acid or iodoform gauze down to the membrana tympani by means of a probe, as this procedure helps to drain the ear.

In all cases of acute otitis media due to cold the drumhead should be incised (after the leech has been applied) as soon as any bulging of the drumhead is apparent, as well as in those cases in which pain and fever persist in spite of all antiphlogistic treatment. In other instances in which spontaneous rupture has taken place, and in which the perforation is very small, the latter should be enlarged. In some cases, after incising the drumhead, no discharge will appear for twenty-four to forty-eight hours, while in others it is so thick and tenacious that it is necessary to force it from the middle

ear into the external meatus by means of Politzer's method. Great care should be exercised in inflating the ear, especially in the early stage, for fear of increasing the inflammation, and under no circumstances should this be done unless there is a large perforation in the drumhead.

If the bulb of the internal jugular vein runs an anomalous course and projects into the middle ear, as sometimes but very infrequently occurs, it is very easy for the surgeon to wound it when incising a drum membrane as occurred in a case reported by Max[1]. The patient had an acute otitis media with a bulging drumhead which was incised. At first a few drops of bloody serum escaped but it was followed immediately by a stream of dark blood. The canal had to be tamponed. The boy was four years old. A month later an examination showed that the whole posterior inferior and a portion of the anterior inferior quadrant of the drumhead appeared bluish white and behind the drumhead one could see a rounded body which became fuller and the bluish white color more intense when pressure was made on the internal jugular vein. There was an identical appearance of the drum membrane on the other side.

After the more acute symptoms have begun to subside, a saturated solution of boric acid should be warmed and instilled into the ear several times a day after the discharge has been thoroughly removed by syringing or by absorbent cotton on a cotton-holder. Some cases do well under the so-called dry method of treatment, consisting in the insufflation of various powders into the external meatus, principally boric acid.

[1] Monatschr. f. Ohrenh. May, 1904.

After the canal has been thoroughly cleansed of all secretion and dried as before, boric acid powder is insufflated, by means of a powder-blower, into the external auditory canal, but should never be packed in, as pyæmia has followed such practice. The dry method should not be employed until all acute inflammatory symptoms have subsided, and never in cases in which there is a small perforation, nor in which there is considerable thick and stringy mucus. Hydrastis canadensis may be combined with boric acid as follows: A paste is made of boric acid and tincture of hydrastis canadensis, and the mass, after it is dry, is thoroughly pulverized and mixed again with equal parts of boric acid powder. Astringent solutions are indicated in cases in which the treatment by boric acid solution or powder has not succeeded. For this purpose, sulphate of zinc, or sulphate of copper, or acetate of lead, in the proportion of three to eight grains to the ounce of water, is generally prescribed.

Injections of warm water through the catheter into the middle ear are recommended by Politzer in cases in which the pain is persistent and does not yield to the usual remedies; also in those cases in which the discharge cannot be arrested by local treatment; and especially in cases in which there is a small perforation (nipple-like) in the membrana tympani, and likewise in cases in which the mastoid process is involved. Whenever the mastoid cells are implicated, the treatment described in the chapter on Mastoid Diseases should be followed. All polypi should be removed by curettes, the snare, or caustic applications. It is of the same importance to attend to the patient's general health in this disease as in acute catarrhal otitis media. After the perforation

has closed, it is advisable at first that the patient wear a small piece of cotton in the meatus, especially during unseasonable weather. The middle ear should be inflated several times a week, and gradually less frequently until the hearing is entirely restored; and the patient should be cautioned to be very careful about exposing himself to draughts and sudden changes in the weather.

CHAPTER VII.

ADENOID GROWTHS, ENLARGED TONSILS, DISEASES OF THE NASAL PASSAGES.

As diseases of the ear are associated so frequently with affections of the nasal passages, the nasopharynx, the Eustachian tubes, and the tonsils, it not only falls within the province of the aural surgeon, but also becomes his duty to examine most carefully into the condition of these structures and passages as well as to apply proper treatment. When the Eustachian tubes are constricted or the patient is unable to breathe through the nose, owing to some obstruction in the nasal passages or to the presence of adenoid growths in the nasopharynx, so that proper ventilation of the middle ear is interfered with, it is useless for the surgeon to practise repeated inflation of the ears by Politzer's method, with the expectation of effecting permanent relief, without first removing the cause of the occlusion.

Great attention has of late years been paid to the treatment of the nose and nasopharynx, and much benefit has resulted from this practice.

When a patient has been deaf for a considerable length of time and serious changes have taken place in the drumhead, with adhesions, and especially if there is tinnitus, one should be very guarded in expressing an opinion as to the benefit to be derived from treatment of the nasopharynx and nares. Too frequently

one sees patients, the victims of chronic catarrhal otitis media, who have undergone operations on the nose and nasopharynx without any improvement in the hearing. It is in the more recent cases that one is apt to meet with success, before pathological changes have taken place in the *membrana tympani.*

ADENOID GROWTHS.

To Dr. Wilhelm Meyer, of Copenhagen, is due the honor of having first directed the attention of the medical profession to the great importance of removing adenoid growths in the nasopharynx, especially when their presence interfered with the passage of air through the nose and Eustachian tube.

Others have followed in Meyer's footsteps, so that at the present time the operation of removal of these growths is performed frequently, and with much success. Not only has deafness been cured in many instances by the excision of an adenoid growth, but diseases of a nervous character, such as asthma, false croup, as well as malformations of the chest, nose, and throat, have been benefited by this operation.

The examination of the ear should never be considered complete until the nose and nasopharynx have been most carefully examined, more especially in the case of children. In about 74 per cent. of Meyer's cases of adenoid disease there was more or less deafness.

As usually observed, adenoid hypertrophy at the vault of the pharynx is of two varieties. In one there is more or less fibrous tissue associated with the lymphoid mass, while in the second variety it is very scarce. The growths vary much as to size and number. There may be simply a chronic enlargement of the pharyngeal or

Luschka's tonsil, situated at the roof of the naso-pharynx, which may fill the space entirely, or the growth may be so small that the surgeon is not justified in removing it. The vegetations may extend from the vault along the posterior and lateral walls of the pharynx, or one large mass or several smaller growths may be found on the posterior pharyngeal wall. The adenoid

Fig. 61.

Typical facial expression of a patient with adenoid vegetations in the nasopharynx.

tissue is, as a rule, bright in color, and bleeds readily when examined with the finger or probe.

Symptoms. The expression of the face is exceedingly characteristic of this disease, as well as the manner in which the patient pronounces certain words. There is difficulty in pronouncing sounds of m, n, and ng, so that the patient talks, as it were, "through the nose."

As a result of mouth-breathing, the patient presents a stupid and listless appearance (see Fig. 61); he is anæmic; has drooping eyelids, prominent teeth, and arched palate. He may have what is known as a "pigeon-breast"; he is apt to complain of frontal headache, a feeling of fulness in the nasopharynx, and the sense of smell is impaired. There is apt to be a persistent catarrh with considerable secretion. The nose becomes pinched, and the alæ have a sunken appearance. Snoring is a prominent symptom. As a result of the obstruction to respiration through the nose, deformity of the septum may occur. The appearance of the face, in consequence, may be changed, owing to the imperfect development of the osseous structures of the nose and hard palate. The hearing becomes affected when the normal ventilation and nutrition of the middle-ear tract are interfered with. The mucous membrane at the mouth of the Eustachian tube becomes swollen as a result of frequent head colds, or the vegetations themselves may encroach upon the orifice of the tube. In consequence, the patient complains of earache, tinnitus, and impaired hearing; and if this condition of affairs is allowed to continue without treatment, permanent changes take place in the structures of the middle ear, so that the drumhead becomes retracted, adhesions develop, or the drum membrane becomes perforated from a suppurative process. Enlarged tonsils often are observed at the same time, especially in children.

Causation. Adenoid hypertrophy is much more frequently seen in children than in adults. In fact, after puberty one meets only occasionally with these growths, although they may be found in rare instances in elderly persons, as well as in infants at birth. Hereditary in-

fluences undoubtedly play an important rôle in the causation, as well as residence in damp places. They occur sometimes as a result of scarlet fever, measles, diphtheria, debilitating fevers, scrofula, and hereditary syphilis.

Diagnosis. In adults, one can usually make a diagnosis of adenoid hypertrophy by means of the head-mirror and posterior rhinoscopy. With children, however, it is almost always impossible to obtain a satisfactory view of the nasopharynx by such means. The surgeon should make a digital examination, and for this purpose should protect his index-finger with a rubber band, to prevent the child from biting his finger. If the examiner has a long finger-nail and the vegetations are soft and flabby, he can scrape away or break up many of these vegetations, and thus accomplish something at this time.

Treatment. In older children who will submit to local treatment, such as the application of chromic acid or the galvano-cautery, these growths may be removed if they are small and soft. Much may be accomplished by the administration of tonics and alteratives, especially if the general health is impaired. The syrup of the iodide of iron, the syrup of hydriodic acid, and cod-liver oil, are remedies that are especially indicated. The most satisfactory and efficient method, however, is the surgical treatment of this condition. In a certain number of patients, who are of a nervous temperament and who are "bleeeders," it is absolutely necessary to use caustics or the galvano-cautery. As regards operative interference, many and various instruments have been devised for this purpose.

Meyer himself used an annular knife, one centimetre

in diameter and twenty-one centimetres in length, which he inserted with the right hand between the septum and the spongy bones as far as the nasopharynx. By introducing the left forefinger through the mouth and behind the soft palate until he felt the instrument, he pressed the vegetations inside the ring. By drawing the knife outward, the growths were cut off. He further recommended a second method, viz., cauterization with solid nitrate of silver.

In adults the adenoid hypertrophy may be removed without an anæsthetic in many instances; but in children, according to my experience, it is always advisable to use ether or chloroform. Ether is to be preferred, as being safer. Whichever one is used, the patient should not be completely anæsthetized, for fear of blood entering the trachea. The instruments that have been most satisfactory in my hands are the Hooper or Knight forceps, although in some instances the artificial steel finger-nail is used with good results.

Hooper's Method. The child is placed in a good light and anæsthetized, and held upright in the lap of an assistant. The operator sitting opposite, after a small-sized mouth-gag has been inserted between the teeth on the right side, removes any mucus that has accumulated in the pharynx. After having made sure of the presence and situation of the growths, the soft palate should be pulled forward and upward by means of a palate-hook held in the left hand. The forceps (Fig. 62) are then introduced, closed, into the nasopharynx, and after touching the growths with the closed end of the forceps, the blades are opened and the growth is seized and removed by direct traction or by a twisting motion of the forceps; force should

never be used. If there is difficulty in removing the growth, the blades should be released and another

Fig. 62.

Hooper's forceps.

Fig. 63.

Knight forceps.

Fig. 64.

Gottstein curette.

attempt made to grasp a smaller portion. The surgeon should examine with his fingers from time to time

to see if the nasopharynx is free. The child's head is then held forward so that the blood may flow out of the nose, while the operator scrapes away with the finger-nail of the left hand all ragged edges in the naso-pharynx. Although Hooper generally relied on the forceps and finger-nails, he found it necessary in some cases to use other instruments.

The surgeon who operates very frequently will find that he requires different instruments for different conditions, although he may prefer one particular instrument.

In using the artificial or steel finger-nail the surgeon should be very careful that it does not come off in the patient's mouth.

The Recumbent Method. The patient is anæsthetized and placed in a recumbent position on an operating-table, and a mouth-gag is inserted on the right side of the mouth. The operator, standing on the left side of the patient, introduces the forefinger of his left hand into the nasopharynx, to locate the position of the growth as well as to guide the point of the instrument.

The Knight forceps (see Fig. 63) are admirably adapted for the removal of these growths. They are introduced closed, and afterward opened, so as to include all of the growth if possible. The patient should be turned to the right side from time to time, to allow the blood to flow out of the mouth and nose. Some blood is swallowed and frequently vomited after the operation. There is less danger that fragments of the growth will fall into the larynx by this method than when the patient is in the upright position. After the operator has satisfied himself that he has removed the growth as thoroughly as possible, he should, for fear of leaving some fragments, either scrape the cavity thor-

oughly with the *steel finger-nail* or else with the Gottstein curette (Fig. 64). By means of good illumination the curette is introduced behind the soft palate and kept well forward until the vault is reached. At this point the instrument is pushed firmly backward and then drawn downward over the posterior pharyngeal wall. To control the hemorrhage, the nares should be thoroughly sprayed with ice-water while the patient is in the recumbent position.

The patient should be prepared beforehand for the operation. The evening before, a mild purgative should be administered; he should be given a light diet on the day of operation, and there should be at least four hours' interval between the last meal and the time of the operation.

Cases of severe hemorrhage after operation have been reported, but they are infrequent, and when they do occur are probably due to the hemorrhagic diathesis. Burger[1] reports the case of a child, eleven years of age, who suffered from leukæmia and who died of hemorrhage after removal of adenoids. According to this writer the danger of hemorrhage increases with age as in seventy-eight cases reported, fourteen were over fifteen years of age. He regards the particles of adenoid tissue which remain after excision as the most important source of the bleeding. On this account, it is important for the surgeon to be well informed as to the patient's general condition. The operation should never be undertaken if the patient has a cold or symptoms of otitis media, for in a case[2] that I reported, in which the operation was performed when the patient was suffering

[1] Presse oto-laryngologique Belge, 1904, No. 4.

[2] Transactions of the American Otological Society, 1892.

from an acute pharyngitis, not only did purulent otitis media follow, but it became necessary to open the mastoid cells. Great care and gentleness should also be employed when operating, for fear of injuring the mouths of the Eustachian tubes.

After the operation the patient should be watched until the hemorrhage has ceased, and should then immeditately be put to bed and kept in a recumbent position for at least twenty-four hours, and for several days should remain quiet in the house. During this time the diet should be a *simple* one. The nares and post-nasal cavity should be sprayed with a Dobell or a mild Listerine solution for forty-eight hours after the operation, if necessary.

In some cases a second operation is required, even if all the growth was removed the first time. In other cases, especially in adults, the remnants of the adenoid growth should be cauterized with solid nitrate of silver after cocaine has been used.

If the tonsils are enlarged, it may be necessary to excise them, as well as to perform some operation on the nasal passages, in order that the patient may breathe easily with the mouth closed.

When the child has been in the habit of breathing with the mouth open for a long period of time, it may be necessary to apply a bandage, to prevent the jaw from dropping at night, and this should be worn for some little time or until the habit has been corrected.

In case of severe hemorrhage, applications of per-sulphate of iron should be made, or it may be necessary to plug the posterior nares. Great care should be exercised so that the patient does not take cold after the operation. Usually after the operation there is a de-

cided improvement in the hearing, provided the latter has not been affected for a long time and permanent changes have not taken place in the drumhead. Even in cases in which there is a discharge from the middle ear the suppuration generally becomes diminished at once. The patient's general health is, as a rule, much improved after the operation. I have seen children who were sickly, poorly nourished, and anæmic, show a decided change for the better after removal of these growths. The following case illustrates this point: A boy, aged four and one-half years, weak-minded, poorly nourished, breathes with mouth open and is very restless at night. There has been a discharge from the ear for about two weeks. Under ether an examination of the posterior nares was made with the index-finger, and the vault was found filled with adenoid masses. These were removed with the Gradle forceps. There was considerable hemorrhage, which soon ceased. Two weeks later there was a very decided improvement in the child's general appearance, as shown by his breathing and sleeping better, and being in much brighter spirits. The hearing was much improved and the discharge less. His mother was very enthusiastic over the good results of the operation.

It does not follow that improvement in the hearing will take place after the operation without treatment directed to the ear itself, for frequently the drumhead is considerably retracted and bound down by adhesions, so that it will be necessary to restore the drumhead to its normal position.

ENLARGED TONSILS.

Enlarged tonsils, or chronic hypertrophy of the tonsils, are undoubtedly an indirect cause of deafness, in

that they interfere often with nasal respiration, and by pressure upward prevent the proper ventilation of the tympanic cavity. When removed, the hearing-distance is often much increased, but the surgeon should always examine the nasopharynx, for frequently in children adenoid hypertrophy is associated with enlarged tonsils.

Hypertrophy of the tonsil consists of a proliferation of its fibrous stroma, together with an unusual increase of its lymphoid tissue. The proliferation of its fibrous stroma is observed more frequently in cases of long standing.

Causes. Enlarged tonsils are frequently due to malnutrition, scrofula, syphilis, tuberculosis, rheumatism, and indigestion. Exciting causes are frequent attacks of acute inflammation from cold, scarlet fever, diphtheria, measles, whooping-cough, and recurring attacks of quinsy sore-throat.

Enlarged tonsils have been observed in children soon after birth. After puberty the tonsils, as a rule, begin to grow smaller, and in some cases resume a normal size and appearance.

Symptoms. The symptoms are very similar to those observed in cases of adenoid hypertrophy. The patient is apt to be poorly nourished, has difficulty in breathing with mouth closed, and there is more or less difficulty in speaking. In some cases the senses of hearing, smelling, and tasting are impaired. Nervous symptoms are sometimes observed, such as spasmodic cough, chorea, and epilepsy.

Treatment. It is advisable to try to reduce the size of the tonsils by internal medication and by local treatment, whenever practicable, before resorting to surgical interference. I think that surgeons are often inclined

to operate too early, and in some cases I believe that much can be accomplished without resorting to excision. In some instances cheesy matter may be forced out of the crypts by pressure made on the tonsils. In cases of malnutrition the child should be given cod-liver oil, syrup of the iodide of iron, or syrup of hydriodic acid, for some time before resorting to operative measures. Astringent sprays of ferric alum are often useful, and in some cases chromic acid may be applied by being fused on the end of a probe and the latter passed into the substance of the tonsil at several different points.

In cases of the hemorrhagic diathesis it is necessary for the surgeon to make use of escharotics, especially chromic acid or the galvano-cautery loop, or of the cold wire.

There are various methods of removing the tonsil. Beside cauterization and écrasement by means of the galvano-cautery, another method is to excise it with forceps and a probe-pointed bistoury. Some prefer scissors, but such practice has for the most part been abandoned. When the galvano-cautery loop is employed, on account of its being a painful procedure, several drops of a 4 per cent. solution of cocaine should be injected into the substance of the tonsil, to produce local anæsthesia. This may only be done in the case of adults. Great care should be taken to include the greater portion of the tonsil within the loop, and this can usually be best accomplished by means of forceps. Traction should never be made upon the loop except when the electric current is passing through it, and the current should be used intermittently.

The most satisfactory method, and the one universally adopted, is tonsillotomy. In the case of young children an anæsthetic is indispensable, and for this purpose ether is undoubtedly the safest, although many prefer chloroform. In order to perform this operation successfully, the services of an assistant or trained nurse are absolutely essential. In the case of adults and children old enough to undergo the operation without ether, the assistant should steady the head of the patient, who sits in a chair with his back to the light and facing the surgeon. With a head-mirror the operator should throw a good light into the pharynx. The assistant, standing behind the patient, should not only support the head with both hands, but should also make pressure with the fingers of each hand immediately below the angle of the jaw over the tonsillar region, so that the tonsil will be grasped more readily by the tonsillotome. In the case of young children in whom it is necessary to use an anæsthetic, the child should be placed on an operating-table, and the same precautions adopted to prevent blood from entering the larynx as in the adenoid operation. A mouth-gag should be inserted after the child is anæsthetized, and while the surgeon throws a light into the pharynx by means of a head-mirror, the instrument should be introduced flatwise, like a tongue-depressor, until it touches the pharynx. It is then turned so that the entire tonsil or as much as possible of it is included within the ring. With the Mackenzie instrument (Fig. 65), the blade is then pushed firmly and steadily through the hypertrophied tissue. The growth usually adheres to the ring, so that it comes away when the instrument is withdrawn. A fork has been added to this instrument, for fear that the growth

might drop into the larynx in cases in which an anæsthetic had been administered. There are several sizes of the Mackenzie tonsillotome as well as of the Mathieu instrument (see Fig. 66). The latter I prefer, but the

FIG. 65.



Mackenzie's tonsillotome.

former is more commonly used by operators. With the Mathieu tonsillotome the ring is slipped over the tonsil, and instead of pushing, as in the Mackenzie instrument, a cutting blade concealed within the ring is

FIG. 66.

Mathieu tonsillotome.

pulled forward. A fork is applied to this, and the growth is pierced by it before excision.

As a general rule, there is but slight hemorrhage after excision, which usually ceases spontaneously. If the bleeding continues, it may be checked by the patient

sucking cracked ice or gargling the throat with iced water, or by making applications of persulphate of iron.

In case of severe hemorrhage where several small arteries have been divided, the bleeding may be controlled by torsion of the vessels, or by application of styptics, or by the galvano-cautery. Pressure is sometimes made over the bleeding surface by means of the finger. Venous and capillary hemorrhages are generally stopped by the application of ice or of some styptic.

In alarming cases, in which the hemorrhage is severe, it may become necessary to ligate the common carotid artery, but such instances are exceedingly infrequent. According to Henking[1] in five cases of severe hemorrhage occurring after the excision of hypertrophied tonsils, the bleeding came from the posterior palatal arch, which had been incised during the operation. After the operation, the patient should be kept very quiet in the house, and should be given a light diet for several days. It should not be forgotten, however, that secondary hemorrhage may occur a week after the operation, so that this fact should be constantly borne in mind. Milk and beef-tea should be prescribed for the first twenty-four hours. A gargle or spray of boric acid is very soothing, and should be used, or a gargle containing permanganate of potash.

EXAMINATION OF THE NOSE.

Anterior Rhinoscopy.

The patient should be seated before the surgeon, and by means of a head-mirror and Argand burner, or by an electric lamp, the anterior nares should be thor-

[1] St. Petersburg med. Wochenschr, No. 11, 1904.

oughly illuminated. A speculum with two blades (see Fig. 29) should be introduced, so that one blade rests

Fig. 67.



Sectional view of the nose, mouth, pharynx, etc. (Gray.)

against the septum, while the other one is pushed upward and outward. The patient's head should then be tilted in different positions, to enable the surgeon to

see the different sides of the nasal cavity. When the patient's head is thrown backward, the portions of the superior and middle turbinated bones will be brought into view, together with the middle meatus, while with the head inclined forward the anterior portion of the inferior turbinated bone will be seen.

Posterior Rhinoscopy.

The patient should be seated before the surgeon with his head slightly thrown back, and the light for illumination placed on a level with his ear. The light is then

Fig. 68.



Tongue-depressor.

reflected from the head-mirror into the patient's pharynx and the focal distance determined. The tongue being held down by means of a spatula (see Fig. 68), the rhinoscopic mirror should first be warmed and introduced, with the mirror surface upward, between the uvula and the soft palate. The patient should be instructed to breathe through the nose, so that there will be as much space as possible between the soft palate and the posterior pharyngeal wall. One should change the position of the mirror, so as to bring into view the vault of the pharynx, the posterior edge of the septum,

the orifices of the Eustachian tubes, and the posterior extremities of the turbinated bones.

In the treatment of aural diseases it is important for the surgeon not only to make a thorough examination of the anterior and posterior nares, but also to remove nasal polypi, and to perform such other operations as may be necessary to establish proper nasal respiration, and thus thoroughly ventilate the tympanic cavity. Nasal catarrh should receive appropriate treatment by sprays and attention to the general health.

The surgeon should be prepared to prescribe for all cases of acute nasopharyngeal catarrh or acute rhinitis, for aural diseases are very often caused by these affections, especially when due to influenza. In a work of this character it is impossible to do more than discuss in a brief manner the more important diseases of the nose and nasopharynx. The reader is, therefore, referred to the standard text-books on diseases of the nose and throat.

The diseases that are of especial interest to the otologist are chronic nasopharyngeal catarrh (including hypertrophic and atrophic rhinitis), nasal and nasopharyngeal polypi, and deflections of the nasal septum.

CHRONIC CATARRH.

In this disease there is a parenchymatous swelling of the mucous membrane, which is due to the presence of enlarged bloodvessels, to the infiltration of serum, and to an increase of the connective tissue—hence results hypertrophy. The cavernous tissue on the turbinated bones becomes excessively swollen, and the patient for a time may be unable to breathe through one side of

the nose. In a short time the same side may become pervious, while the other one will be closed.

As a result of chronic catarrh, the posterior extremity of the middle and lower turbinated bones may become so much hypertrophied that the tissue may protrude into the nasopharynx and be mistaken for a polypus. True hypertrophy of the mucous membrane is due more especially to an increase in the fibrous or connective tissue. Increase of the cavernous tissue generally affects the anterior portions of the turbinated bodies.

Chronic Hypertrophic Catarrh, or Hypertrophic Rhinitis.

This affection may follow frequent attacks of acute catarrh, or may be due to impairment of the general health, gout, rheumatism, the excessive use of stimulants and tobacco, and hereditary causes. It is very seldom that one meets with a normal condition of the mucous membrane of the anterior and posterior nares. The tendency of the present day is, I think, to operate more frequently than is absolutely necessary. If a number of persons should be examined, who did not even complain of any symptoms referable to the nose or throat, it is probable that evidences of catarrhal inflammation, or spurs on the septum, or hypertrophy of the mucosa covering the turbinated bodies would be found in many instances. There is undoubtedly an excessive inclination to operative interference on the part of many rhinologists, especially when the deafness seems to be due to catarrhal inflammation. Many promises to improve the hearing by operative measures

on the nose have not been fulfilled, for the reason that catarrhal changes had already taken place in the drum-head, and in some instances in the internal ear. A surgeon should not operate on the nose simply because the condition of the mucosa covering the turbinated bodies is not normal, unless the symptoms are such as to warrant interference.

Symptoms. There is a certain amount of nasal obstruction owing to swelling of the tissues, but this varies in intensity from time to time according as to whether the erectile tissue covering the turbinated bodies is involved. The voice may have a nasal twang, while the patient, breathing with open mouth, presents a stupid appearance. The patient complains of a dull feeling in the head, difficulty in smelling and tasting, and sneezes frequently. There is usually a marked hypersecretion of mucus of either a serous or a viscid character.

An examination reveals an abnormal redness of the mucosa with a certain amount of hypertrophy. The anterior extremity of the inferior turbinated body may be enlarged, or the mucous membrane covering the entire body may be thickened and corrugated. The inferior turbinated body is more frequently affected than the middle one, although the latter may be decidedly enlarged and press upon the septum. By posterior rhinoscopy, one may find the posterior extremities of the middle and inferior turbinated bodies very much hypertrophied, so that when projecting into the naso-pharynx the growth may present the appearance of a mulberry-like polypus.

When the stenosis is due to simple enlargement of the erectile tissue, applications of a 4 per cent. solu-

tion of cocaine will cause a marked shrinking of the tissue, while in cases of true hypertrophy cocaine will have but little effect.

Treatment. In cases in which there is but slight hypertrophy, much may be accomplished by having the patient spray the anterior nares with Dobell's or Seiler's antiseptic solution. Astringent sprays should be used by the surgeon, and in cases in which there is but slight hypertrophy, the latter may be cauterized with chromic acid. If the hypertrophied mass is sufficiently large so that a loop can be passed around the growth, it should be removed by means of the galvano-cautery. If snaring is inapplicable, the flat point of the cautery may be used to destroy the hypertrophied tissue.

The superficial sloughs will fall off in a few days' time or within a fortnight. The application of the cautery or of caustic agents should be repeated until the nose is pervious, but a week or two should intervene between the applications. In the meantime the patient should be careful not to take cold, and should spray the nasal passages with a boric acid solution.

Beside local treatment, which is most important, benefit will also follow from attention to the patient's general health, by correcting any digestive disturbance, and prescribing for any gouty or rheumatic disease. Change to a dry atmosphere should be recommended in many instances.

Chronic Atrophic Catarrh, or Atrophic Rhinitis; Ozæna.

In this disease atrophic changes occur, so that the nasal passages are usually more pervious than in a normal case. The atrophy, as a rule, is most apparent

in the inferior turbinated body. The nasopharynx and pharynx are apt to be involved as well, and when the hearing is impaired it is usual to find the Eustachian tubes dilated, so that air is easily forced into the tympanum by Politzer inflation or by catheterization. The same atrophic changes may take place in the middle ear and drum membrane, so that the latter may be exceedingly thin. Great care should be used in inflating the ear when this has occurred, for I have seen the membrane ruptured when forcible inflation was practised. There is very little secretion in the nasal passages, and the latter are generally exceedingly dry. When green crusts form, which give rise to a very offensive odor, the disease is called *ozæna.* In simple atrophic rhinitis attention should be paid to the patient's general health, and alkaline sprays, or a spray of albolene, containing 2.5 per cent. each of camphor and menthol, should be used.

In ozæna the proper treatment is to remove the crusts as soon as they form, and advise the patient to use daily a spray of permanganate of potash, boric acid, carbolic acid, or some other disinfectant. Some rhinologists recommend sprays of aceto-tartrate of aluminium, or of tincture of iodine. Formalin in the proportion of two to two and a half per cent. in an alkaline spray, such as Seiler's, is very useful in a certain proportion of cases. Others advise the insufflation of antiseptic powders, boric acid, iodol, etc., while the application of the electric cautery has been recommended as well as the use of the constant current.

Chronic Nasopharyngeal Catarrh.

The nasopharynx participates in the same changes as are found in chronic hypertrophic or atrophic rhinitis and chronic pharyngitis.

An examination with the rhinoscopic mirror will reveal the nasopharynx as being simply hyperæmic, with more or less swelling of the mucosa, and occasionally a swollen condition of the lips of the Eustachian tubes, or there may be a granular condition similar to that observed in granular pharyngitis, or atrophic changes may have taken place.

The usual symptoms are a feeling of fulness and discomfort in the nasopharynx, with an accumulation of mucus, which the patient tries to get rid of by hawking or swallowing. At night the patient is apt to complain of a dropping of mucus from the nasopharynx. The above symptoms are observed only in cases in which there is a hypersecretion.

Nasopharyngeal catarrh is a disease that is very prevalent in the United States, especially in the larger cities. It is undoubtedly due to the sudden and extreme changes of temperature that are so common, and also in a large measure, I believe, to a fine dust, especially in New York City. Persons in impaired health are subject to the disease as well as those suffering from a rheumatic and gouty diathesis. Other deleterious influences are living in steam-heated rooms, the abuse of alcohol and tobacco.

Catarrh of the nasopharynx is often much benefited by removing some obstruction in the anterior nares, so that the surgeon should be sure to secure permeability of the nasal passages in all cases.

The nasopharynx should be sprayed with alkaline solutions in order to remove all secretion. Afterward, astringent sprays should be used, or applications made of nitrate of silver, in simple uncomplicated cases.

Where there is a granular condition, it becomes necessary to use caustics, viz., nitrate of silver or chromic acid, or the galvano-cautery. Curetting the pharyngeal wall will often effect a cure of granular pharyngitis.

In cases of ozæna the crusts should be removed and the same treatment followed as when the disease affects the anterior nares.

Some authors advise scraping the remains of the pharyngeal tonsil, or the local application of the electric cautery after the crusts have been removed.

NASAL POLYPI.

The mucous polypus is the form of tumor most frequently met with in the nasal passages, and generally is found attached to the middle turbinated body. It is usually round and smooth in appearance, frequently pear-shaped, and of a bluish-gray color. It may, however, not form a distinct tumor, but simply a fringe, constituting what is called "polypoid hypertrophy." These polypi are, as a rule, bilateral and multiple, and vary in size from a very small pea to a walnut. Beside the mucous polypus, there are other varieties of tumor which are of rare occurrence, but nevertheless are occasionally observed, viz.:

1. A polypoid growth, usually single, reddish in color, and containing fibrous tissue.

2. Papilloma, either single or multiple, which is apt to be attached to some portion of the inferior turbinated body or the adjoining parts.

3. Angioma.
4. Adenoma.
5. Cysts.
6. Osseous tumor.
7. Malignant growth.

Treatment. The most successful method of removal of nasal polypi is by means of the cold wire or galvano-cautery snare. The surgeon should have thorough illumination of the parts, and after applying a 4 per cent. solution of cocaine should examine the nasal passages with a blunt probe. In some cases it is necessary to introduce the finger into the nasopharynx. After removing the polypus with the snare, the root should be cauterized with chromic acid or the galvano-cautery. The older plan of using forceps and torsion is at the present time generally discarded.

It may require several sittings before the patient is cured.

Polypi of the Nasopharynx.

Polypi found in the nasopharynx are usually of a fibro-myxomatous character, but occasionally are very hard, and consist of fibrous tissue.

Fibro-myxomatous Polypi.

These growths are somewhat firmer in consistency than those found in the anterior nares. They are said to be more frequent in women than in men, and, as a general rule, are attached by a slender pedicle near the borders of the choanæ.

The symptoms are those due to nasal obstruction and pressure on the palate. A bluish growth, quite freely movable, will be seen with the rhinoscopic mirror. The

polypus may be distinguished from a true fibroma on account of its color, mobility, and softer consistency. It should not be mistaken for a cyst, owing to absence of fluctuation.

The growth can generally be removed by means of the cold steel wire, which is slipped around the pedicle. Some prefer to use the galvano-cautery, while others consider evulsion by forceps to be a more satisfactory method.

The remains of the pedicle should be treated with chromic acid or the galvano-cautery.

FIBROMATA.

A fibroma springs from the periosteum and bone, and is attached most frequently to the body of the sphenoid and the basilar process of the occipital bone. It is firm to the touch, of a pale reddish color, and consists of fibrous tissue, round cells, and frequently of dilated vessels. It occurs more often in males than in females, and the development of the growth is not very active after the twenty-fifth year. The tumor is not, *per se*, malignant, but, on account of the destruction of adjacent tissue caused by its growth, it is of a dangerous character. It may extend into the nasal passages and cause the nasal bones to expand, giving the patient a deformed appearance called "frog-face." Adhesions may take place in the nose and pharynx, whereby its nutrition may be increased. When ulceration follows there is apt to be a fetid discharge. The growth may extend to the orbit and give rise to exophthalmos. Deglutition and respiration may be interfered with when it projects downward toward the larynx. The cranial cavity may be invaded, and great pain follow from pressure on cer-

tain nerves. Deafness results in some cases from pressure on the orifices of the Eustachian tubes.

Exceptionally cases occur in which spontaneous cure is established by a sloughing off of the growth. The prognosis is always grave, especially when adhesions have taken place and the patient is suffering from pressure-symptoms.

The diagnosis is generally made by posterior rhinoscopy and palpation. A fibroma is hard and firm to the touch, and usually of a pale or dark red color. It is very important to find out first of all about the attachment of the tumor, in order to decide if the growth can be removed by means of the cold snare or the galvano-cautery loop. When the base is attached to the pharynx alone, the electric loop can be passed around the growth either through the nose or the mouth. Only a moderate degree of heat should be applied when the galvano-cautery loop is used. The galvano-cautery should also be employed to destroy the remnants of the tumor after its removal. Some surgeons prefer the method of evulsion, while electrolysis is recommended by others. When the growth extends beyond the pharynx, various operative measures have been devised for excising it—namely, an operation performed through the mouth, through the nose, or through the superior maxillary bone. It is, however, beyond the province of this book to discuss such operations. Beside the growths already mentioned, malignant tumors are met with occasionally.

DEVIATION OF THE CARTILAGINOUS NASAL SEPTUM.

Deviation of the cartilaginous septum, when of recent origin, is probably due to a blow or fall, while in cases

of long standing the cause is probably one of faulty development, congenital in character. The cartilaginous septum may be but slightly deflected, or the nasal passage may be so constricted that the patient can with great difficulty force any air through by an expiratory effort. As one side becomes narrowed the other one is correspondingly dilated.

Symptoms. The patient is apt to breathe with mouth open, especially at night. There is obstruction to respiration on one side, so that when the patient is requested to blow through the obstructed side, the nostril on the dilated side being closed, very little or no air will pass through. By closing the constricted nostril and repeating the same procedure a free current of air is obtained. Other symptoms are: disturbed rest at night, a nasal tone to the voice, epistaxis, and, what is of special concern to the otologist, more or less deafness, or a sense of fulness in the ear, or tinnitus, due to interference with the proper ventilation of the tympanic cavity. The nasal passage being obstructed, it is easy to understand how the drumhead becomes retracted in the act of swallowing; unless this condition is relieved at an early date serious changes are liable to occur in the drumhead, so that a permanent disturbance of hearing results. According to my experience, a very common symptom is a feeling of fulness in the ear, which is very annoying to the patient. It is usually temporarily relieved by inflation of the middle ear, but soon returns. This disease also gives rise at times to headache and reflex symptoms. A diagnosis can readily be made by inspection with a nasal speculum (Fig. 29). The deviated portion is easily seen, constricting the nasal passages, while on the other side the nasal space is unusually

dilated. The latter condition will assist the surgeon in differentiating from tumors and exostoses.

Numerous operations have been proposed and per-

Fig. 69.

Asch's separator, with blunt edge.

formed for relieving this condition, the galvano-cautery being used by some, while the saw is preferred by others. The operation that seems to me to be most

Fig. 70.

Asch's separator, with sharp edge.

successful is the one devised by Dr. Morris J. Asch.

The Asch Operation. The instruments used by Asch are two separators (Figs. 69 and 70), one with a

Fig. 71.



Asch's scissors.

blunt edge and the other sharp, to break up any adhesions existing between the septum and the turbinate on the affected side, as also to destroy any posterior obstructions. The scissors (Fig. 71) are of two sizes, being

powerful instruments, curving outward from the point of crossing and meeting again in front. Of the blades,

FIG. 72.



Asch's scissors.

one is blunt, while the other is sharp with a triangular cutting surface, the apex of which is at the tip. The

FIG. 73.



Asch's compressing forceps.

other pair of scissors (Fig. 72) are made with the blades bent at right angles, and are occasionally necessary for

FIG. 74.



Asch's tubes.

those deviations that lie low down on the floor of the nose. Of two compressing forceps (Fig. 73), one has a long

and the other a short beak. Hollow tubes of vulcanite are used (Fig. 74), which have a slight curve on the under surface, corresponding with the normal curve at the entrance of the vestibule of the nose. The tubes are perforated, to prevent them from slipping when inserted, and vary in size and shape. The Asch tubes (Fig. 74) are five in number, and are so constructed that the heat of a sterilizer does not destroy their shape.

The patient should be thoroughly anæsthetized, as the local application of cocaine is not sufficient. The nose should be sprayed first of all with an alkaline antiseptic solution. All instruments, etc., should be sterilized. After the patient is completely anæsthetized the cone is removed and is not usually re-applied, as the operation is quickly performed, as a rule. The head should then be drawn backward, so that blood will not enter the larynx, and the table should be so arranged that the surgeon can use either direct or artificial light. Any adhesions existing between the septum and the adjacent mucous membrane are then broken up by means of the blunt separator, which has been introduced sideways on the deviated side, or by using the little finger. The operator is thus enabled at the time to detect the presence of bony obstructions posteriorly, in which case, and also if the adhesions cannot be broken up, the separator with sharp edges is substituted. Hemorrhage may occur at this step of the operation, but is generally controlled by an iced spray or by pressure made with cotton on an applicator. The scissors are then introduced open and parallel to the floor of the nose, the sharp blade in the concavity or dilated nasal passage, while the blunt edge is over the point of greatest convexity. The instrument is next closed, so that the blade

cuts through the cartilage into the opposite side. The instrument is then opened and withdrawn, but immediately reintroduced, the direction of the instrument being upward and pointing to the frontal bone, the cutting blade again in the cavity and crossing the line of the first incision at as nearly a right angle as possible, and at its centre. The incisions will thus intersect each other. The second incision is made by closing the scissors firmly. There are, therefore, four segments as a result of the crucial incision, and these are pushed forcibly into the dilated passage by means of the finger introduced into the deviated side. One blade of the powerful blunt forceps is next introduced into each nostril and firmly closed, so as to straighten the septum and force the broken segments to override each other on the concavity. After withdrawal of the forceps, an iced solution is sprayed into each nostril and sterilized tubes introduced into the nose, a closely fitting one on the constricted side and a smaller one on the dilated side, in order to cause equable pressure, this being usually sufficient to control all hemorrhage.

After the patient is placed in bed, cold antiseptic sprays are used every half hour, and iced cloths are applied externally. The tube in the former concave side is withdrawn permanently twenty-four hours after the operation. Twenty-four hours later, the tube on the constricted side is removed and cleansed, while the nose is sprayed and cleansed with cotton applied to applicators, cocaine being used if necessary. The same tube is generally reinserted. The patient at this time is allowed to sit up. During the next three or four days, the tube is removed each day and cleansed, and during this time the nose is cleansed, and the patient is in-

structed how to extract and reintroduce the tube. After this the patient is seen once or twice a week for four weeks, and the tube is withdrawn permanently five weeks after the operation. Occasionally the lower segment of the septum remains thickened, and presents the appearance of a spur. This should be removed by the electro-trephine or galvano-cautery after the parts have been cocainized.

CHAPTER VIII.

OTITIS MEDIA CATARRHALIS CHRONICA, OR CHRONIC CATARRHAL OTITIS MEDIA.

THE acute form of catarrh, or frequent attacks of the subacute variety, lead to a chronic inflammation of the middle ear. In the chronic form, after exudation has occurred, adhesive processes in the middle ear are apt to develop, leading to adhesions between the drum membrane and tympanum, rigidity of the chain of ossicles, and anchylosis or fixation of the stapes. It is not necessary, however, that there should be first of all an exudative catarrh before the adhesive process commences, for very frequently the latter starts in as an interstitial inflammation in the mucous membrane of the tympanic cavity, with simultaneous involvement of the labyrinth. Some writers believe that, in these cases, the same pathological process begins at the same time in both the labyrinth and middle ear. This form of interstitial inflammation with labyrinthine implication should in many instances be classified as a disease *per se*, since it has no connection whatever with the exudative catarrhal process. It has been called "dry catarrh of the middle ear," or "sclerotic otitis media," and usually gives rise to considerable deafness and tinnitus, while the pathological changes are usually most marked about the fenestræ ovalis and rotunda.

In the exudative or moist variety of chronic catarrh the mucous membrane is congested and swollen from

infiltration with serum. There is an increase of connective tissue which covers the incus, malleus, and attic, as well as the fenestræ ovalis and rotunda. There is more or less hypertrophy of the lining membrane of the tympanum, due to an increase of round cells and also to new connective-tissue formation. At this stage partial resolution may take place through fatty degeneration of the round cells. The tympanic cavity

FIG. 75.



Internal surface of the left membrana tympani. (Enlarged.) (POLITZER.)

A. Head of the malleus. B. Neck of the malleus. C. Tendon of the tensor tympani muscle and anterior fold of the membrana tympani. D. Inferior extremity of the handle of the malleus. E. Anterior portion of the membrana tympani. F. Posterior fold of the membrana tympani and chorda tympani. G. Incus. H. Short process of incus. I. Long process of incus.

frequently contains thick mucus. The mobility of the ossicular chain becomes impaired, and in consequence there results more or less deafness, which may be relieved at this time by treatment, especially if adhesions have not already taken place. After cessation

of the discharge the mucous membrane loses more or less of its red color, and becomes pale, although it is still thickened. The different folds of the lining membrane of the tympanum, and the ligaments which bind the ossicles together and to the walls of the middle ear, become rigid and thickened. In rare instances the mucous membrane undergoes a partial calcification. Adhesions take place between the inner wall of the middle ear and the drumhead by means of bands or

FIG. 76.

Section showing the inner or mucous surface of the drumhead and ossicles (normal).

striæ. The ossicles may be attached to the tympanic walls, and anchylosis may occur, so that the power of vibration of the drum membrane is seriously interfered with. The attic at times undergoes more or less obliteration from the excessive formation of connective tissue and from becoming joined with the ossicles and the inner surface of the drumhead. The conduction of the sound-waves by means of the ossicular chain becomes seriously interfered with at this stage, owing to the formation of such adhesions and the occurrence of anchylosis. The three bones may be anchylosed, or there

may be a separate attachment between each bone and the tympanic wall. Anchylosis between the stapes and fenestra ovalis gives rise to considerable deafness and to serious results. The edge of the foot-plate may be united with the margin of the fenestra ovalis, or the crura may be bound down by adhesive bands or striæ.

Fig. 77.

Horizontal section through human stapes in foramen ovale. 1 anterior and 2 posterior crura; 3 foot plate of stapes; 4 articulation of incus with head of stapes; 5 stapedius muscle; 6 facial nerve; 7 connective-tissue adhesions (pathological).

Such anchylosis may follow an inflammation of the entire tympanic cavity, and be associated with more or less anchylosis of the malleus or incus, or of both of them. There may be at the same time numerous striæ and bands running through the tympanum, with thickening of the membrane covering the fenestra ovalis. Instead

of the entire drum cavity being involved, the disease may be limited to a circumscribed inflammation of the sclerotic or interstitial variety, and be confined to the region of the fenestra ovalis. Bony union between the stapes and fenestra takes place in some instances, and when this occurs the deafness is incurable.

Certain changes are observed frequently in the fenestra rotunda, such as thickening and calcification. The changes in the Eustachian tube depend very much on the nature of the disease in the tympanum itself. There is frequently constriction of the tube in the early stage, when the mucous membrane of the tympanic cavity is swollen and filled with round cells, and before shrinking of the tissues and sclerosis have taken place. There is usually at the same time catarrhal inflammation of the nasopharynx. It is in this stage of the disease, before adhesions have produced fixation of the ossicles and drumhead, that much benefit follows from inflation of the tympanum, and appropriate treatment of the nasopharynx and the mouths of the Eustachian tubes.

In the circumscribed sclerotic form of inflammation, limited mostly to the neighborhood of the fenestra ovalis, the tube is, as a general rule, normal as to its permeability, although there are occasional exceptions, when there is some constriction. Occasionally the ventilation of the middle ear is interfered with, owing to certain changes having occurred in the muscles of the Eustachian tube. The intratympanic muscles are likewise subject to atrophy, fatty or colloid degeneration. Habermann[1] has examined the temporal bones of a number of patients who had had the sclerotic form of

[1] Arch. f. Ohrenh., Bd. 60, Heft, 1 u 2.

otitis media and he found the characteristic lesion to consist of small circumscribed points of ostitis, to the number of one, two, three, rarely more, which traversed the bone, the point of departure being probably about the borders of the round or oval windows. According to this writer the disease attacks the labyrinthine capsule but hardly ever the membranous labyrinth.

From this description of the pathological changes that are usually observed in this disease, it will be seen that *chronic catarrhal otitis media* is a term that should include cases of *secretive catarrh*, in which there are considerable hyperæmia and swelling of the mucous membrane, followed by the formation of adhesions and anchylosis of the ossicles, as well as cases of *selerotic* or *interstitial inflammation*, which is confined to a limited portion of the tympanum, usually the neighborhood of the fenestra ovalis or fenestra rotunda. It very frequently happens that the patient does not seek advice during the first stage, when the surgeon could accomplish very much by treatment, but only comes when adhesions and anchylosis have occasioned more or less deafness, which interferes with his occupation. The tinnitus at this time is often most annoying and distressing.

Heredity plays an important rôle in the etiology of chronic catarrh of the middle ear, and especially in that form of circumscribed sclerotic inflammation, although very often patients are unable to ascribe the disease to any such influences. Other causes are frequent attacks of acute or subacute catarrh of the middle ear especially in childhood, chronic nasopharyngeal catarrh, diphtheria, the exanthematous diseases, tuberculosis, syphilis, Bright's disease, anæmia, pregnancy, gout, and rheumatism,

residence in a damp climate, sea-bathing, and the excessive use of alcohol and tobacco. Young persons and children are not so frequently affected with chronic catarrh (adhesive form) as the aged and those in middle life. The former are more frequently the subjects of the so-called moist variety of catarrh, which, as a rule, affects both ears.

As chronic catarrh of the middle ear usually is considered as an extension of catarrh from the nasopharynx, it is important, in such cases, to examine thoroughly the latter cavity as well as the nasal passages and pharynx. In some instances the patient has much difficulty in breathing through the nose with the mouth closed, owing to an hypertrophic rhinitis, the presence of a spur or polypus, or a deviated septum. In other cases the nasal passage is unusually dilated, owing to atrophic changes, and frequently at the same time the Eustachian tube is found in a similar condition. In such individuals inflation of the ear is apt to cause little if any improvement in the hearing-distance, while tests with the tuning-fork often show loss of bone-conduction. An examination of the throat in one case reveals a thickened and dull-red mucous membrane (hypertrophic); in another, the mucous membrane is unusually dry and has a glazed appearance (atrophic). Then, again, the pharyngeal wall may be studded with small elevations, with enlarged vessels leading to them (follicular pharyngitis).

Appearances of the Membrana Tympani. The membrana appears dull, lustreless, and more or less opaque. Certain portions of the drumhead may be normal in appearance, or the entire membrana may be thickened and milky white (see Fig. 78). The opacities vary greatly

in shape, being semilunar at times (see Plate V., Fig. H), or there may be a peripheral or circular opacity along the lower border of the membrane, which is compared to the arcus senilis of the cornea. Chalky or calcareous deposits, often crescentic in shape, are found at times in front of the handle, although they are observed also occasionally in the posterior segment. Again, there may be chalky spots in both anterior and posterior segments, or a chalky deposit in one portion, while other parts of

FIG. 78.



Section of the temporal bone, showing marked retraction and thickening of the drumhead and anchylosis of the ossicles.

the membrana are opaque from thickening. The drumhead, beside being milky white or yellowish, like parchment, may assume a bluish-white appearance or even reddish-gray, the latter color being due frequently to a congested condition of the middle ear, while the drumhead is more or less transparent from atrophic changes. In some instances the membrana becomes very thin, so that the different portions of the middle ear may be clearly seen, as well as the incudo-stapedial articulation.

In such a case the drumhead may be ruptured when forcible inflation of the ear is practised according to Politzer's method. The Eustachian tube is apt to be freely pervious, so that one should always be most cautious in forcing air into the tympanum whenever such conditions are present.

There may be atrophy of the membrane in one portion, while in another portion there is considerable thickening. On account of the retraction of the drumhead, the handle of the malleus appears foreshortened and drawn inward and upward, and especially so when the inferior extremity is adherent to the inner wall of the middle ear, while the short process stands out prominently, as well as the anterior and posterior folds of the membrana (see Fig. 79). The membrana flaccida is apt to be sunken in. There may be deep depressions in the drumhead, where the membrane is attached to the tympanic wall. In consequence of the inward and upward displacement of the long handle of the malleus, the cone of light becomes distorted and frequently very much contracted. In cases of sclerotic inflammation limited to the region of the fenestra ovalis, an important symptom is the appearance of congestion of the promontory, which is seen through the membrana. In this same class of cases, however, there may be simply anchylosis of the stapes, while the drum membrane shows every evidence of being normal.

Symptoms. The first symptom to cause annoyance to the patient and make him seek advice, as a general rule, is a *tinnitus or subjective noise* in the ear, which, though slight at first, gradually becomes more intense, so that he is often very much disturbed by it, while in exceptional cases where the patient is of an extremely

nervous temperament, and the disease has lasted for a long time, melancholia has followed. Tinnitus, on the other hand, in cases of chronic catarrh, in which there is some secretion, is usually of a transitory character or else absent altogether. In the adhesive and sclerotic varieties, while the noises may be intermittent at first, they generally become continuous as the disease progresses, especially when anchylosis of the stapes with the foramen ovale has occurred. The noises are apt to be most distressing when the weather is damp and muggy, or after the patient has indulged freely in alcoholic drinks, or is mentally fatigued, or otherwise exhausted from some indisposition. They are most likely to be the first premonition of commencing deafness, but this is not the rule, for in some instances the noises and the deafness develop at the same time, while in other cases the first symptom is the deafness. The noises are apt to grow louder as the deafness increases, but they may, on the contrary, become less distressing, and even cease altogether when the deafness becomes extreme.

The deafness depends very much on the changes that have taken place in the drumhead, on the extent of the adhesions, and more particularly on the condition of the foramen ovale and foramen rotundum. The hearing often varies at different times of the same day, being more impaired at night, when the patient is fatigued. Change of climate, especially to a high altitude, often causes considerable improvement in the hearing as long as the patient remains there, only to be followed by loss of hearing when he returns to a damp region. Other symptoms are a sense of weight, fulness, or pressure in the head, occasional vertigo and giddiness, which

may be permanent, unsteadiness in the gait, with nausea and vomiting. A patient suffering from chronic catarrh of the middle ear at times complains of pain in or about the ear, which is generally due to neuralgia, as this disease is often observed in neurasthenic subjects. Some persons are disturbed by certain sounds, particularly shrill tones, even when the degree of deafness is quite marked (hyperæsthesia acustica). The hearing of ordinary conversation, especially at a dinner or reception, when several persons are talking at the same time, is extremely difficult for a patient suffering from chronic catarrh of the middle ear. He frequently hears the higher notes, especially musical ones, extremely well, and this is often the case when the labyrinth is not affected. If, however, the latter is involved, as shown by tests made with the tuning-fork, not only will there be deafnesss for speech, but also for the watch and acoumeter, and for musical sounds, in the higher register.

The ability to hear better in a noise, more particularly in a carriage or railway train, *paracusis Willisiana*, so-called, is a symptom that frequently is noticed in this class of patients. A patient under my care could hear only very loud conversation very close to the ear, but she could understand easily every word spoken in an ordinary tone of voice when riding in a railway train. In cases in which the auditory nerve is affected without involvement of the middle ear this phenomenon is not observed. Some otologists believe this symptom to be due to an increased excitability of the auditory nerve brought about by the concussion, while another explanation (Politzer) is "that the improvement in hearing is due to the movement of the inflexible auditory ossicula, so that they are placed in a more favorable condition for

the conduction of the waves of sound, and that at the same time the endings of the auditory nerve are set in vibration, producing an increased sensibility." Although this symptom is undoubtedly due to middle-ear catarrh, it may be observed in patients in whom the labyrinth is secondarily involved, for it is noted in cases in which bone conduction is normal as well as in those in which it is impaired or lost. It generally is considered as an unfavorable symptom.

When left to nature, the deafness generally increases, so that eventually the patient has great difficulty in hearing at all. Some cases progress slowly, others very rapidly. If the patient is subject to frequent head-colds, the disturbance of hearing is apt to increase rapidly, also in cases in which the labyrinth is suddenly involved, or in which the patients are suffering from some exhausting disease, such as syphilis or tuberculosis. Profound deafness is not very frequent. After steadily growing worse for a number of years, the disturbance of hearing may remain quite stationary for a time. When the disease is confined mostly to the region of the fenestra ovalis, the tinnitus is apt to be troublesome for some years before the patient really knows that he is deaf. Occasionally the deafness is limited to one ear, but this is only for a time, because when the previously normal ear becomes involved the deafness progresses very rapidly as a rule.

Diagnosis. Inspection of the drumhead reveals the existence of chalky deposits or a thickened condition of the drumhead. On the other hand, the surgeon may find atrophic changes. The membrane is more or less retracted, with prominence of the short process and of the anterior and posterior folds. The cone of light is

distorted. By having the patient try the Valsalvian experiment, as well as by using Siegle's otoscope, one can observe the mobility of the drumhead. By such means the atrophic portions bulge outward when the air in the canal is exhausted, while the adherent portions remain stationary. By means of the diagnostic tube, when the ears are inflated, the permeability of the Eustachian tubes can be determined. When the labyrinth is not involved, the bone-conduction, as shown by Weber's and Rinné's tests, is better than the aërial conduction, and the higher tuning-forks are better heard by aërial conduction than the lower ones. If the disease is complicated with marked labyrinthine involvement, then the lowest tuning-forks are heard by aërial conduction, while bone-conduction is very much impaired or even lost.

The probe may also be employed to advantage in order to determine as to whether the membrana is adherent to the tympanic wall or not.

Prognosis. Much depends on the patient's general health and condition, as well as on the changes that have taken place in the middle ear and labyrinth. The prognosis is much less favorable in those insidious forms of sclerotic catarrh limited to the neighborhood of the fenestræ, and particularly when the labyrinth is involved, than in cases in which the catarrh is of the so-called moist variety. When a patient seeks advice who has had a continuous subjective noise for several years, and which he likens to the sound of escaping steam, and moreover if the Eustachian tubes are pervious, and inflation by Politzer's method and the use of Siegle's otoscope do not produce any change for the better in the hearing-distance, the prognosis in such

cases as to any permanent improvement is extremely unfavorable. Such patients are likely to be neurasthenic or hypochondriacal, and frequently inherit some tendency to deafness. The prognosis is much more favorable when the tinnitus and deafness show even a slight improvement after inflation of the middle ears. If the disturbance in hearing has come on very gradually, the chances are that the patient will not be profoundly deaf, even if he receives no treatment whatever. On the other hand, if the patient has suddenly become hard of hearing, and there is a continuous subjective noise which is not modified by treatment, and particularly if the labyrinth is involved, the probability is that the disease will advance rapidly, and the prognosis is bad, especially if the patient is suffering from some constitutional disease, such as syphilis, anæmia, or tuberculosis. A patient's occupation also has much to do with the prognosis. Gout and rheumatism play a very important part in this disease, for one meets with chronic catarrh of the middle ear very frequently in such patients. Subjective noises, which sound to the patient like escaping steam, or are pulsating in character, and which are continuous and have existed for a long time, are unfavorable symptoms. On the other hand, if the noise is intermittent, or if the surgeon is able to arrest it even for a short space of time, and at the same time cause a slight improvement in the hearing-distance, the chances of some permanent improvement are fairly good.

The question often arises as to what is the duty of the otologist under conditions just mentioned, in which the prognosis is unfavorable, namely, as to whether he should hold out any hope to the patient of improve-

ment. It is possible for the surgeon to determine, in a very short time, whether the hearing is likely to be improved by treatment, and until the patient has been seen at least two or three times, and the middle ears have been inflated, the prognosis should be a guarded one. If at the end of this time there is any improvement in the hearing-distance or the subjective noise, the treatment should be continued. On the other hand, if there is no change whatever, the patient should be informed of the fact that treatment is not likely to be of much if any benefit; but, as the disease is likely to progress steadily, it is advisable that he should receive regular treatment for a period of six weeks at a time during the spring and fall, in order to arrest if possible the progress of the disease. He should in the intervals consult the surgeon if there is any change for the worse in the symptoms. As previously stated, many of these patients are extremely nervous in temperament, and require encouragement as well as treatment directed to their general condition. Otherwise, if told that the disease is incurable and that treatment is useless, such patients become very despondent, and the deafness is liable to become much worse. They need, therefore, some moral support, and should be under the supervision of the otologist, who should not only see them occasionally, but should also encourage them by stating that possibly by treatment the disease may be arrested. Iron and strychnine and the hypophosphites are remedies to be prescribed in many instances.

Treatment. The treatment of chronic catarrh of the middle ear should be so directed as to obtain free ventilation of the middle ear, and the restoration of the membrana tympani to its normal position if possible.

Catarrh of the nasal passages and nasopharynx should receive appropriate treatment, while if there is any obstruction in the anterior nares or nasopharynx it should be removed if it interferes with the entrance of air into the middle ear. The mouths of the Eustachian tubes should be inspected, and their permeability tested by Politzer's method of inflation or by catheterization. (See chapter on Diseases of the Nose and Nasopharynx.) In the so-called secretive or moist form of catarrh, particularly if of short duration, and if attended with more or less swelling of the Eustachian tube, considerable improvement will follow the injection of air into the tympanic cavity. According to my experience, the catheter is much more efficacious for this purpose than the method of Politzer, while the improvement is more marked. Inflation in cases of adhesive catarrh is not followed by the same improvement as when the disease is of the secretive or exudative variety. The surgeon, when forcing air into the tympanum, should at first use but little force, either by Politzer's method or by the use of the catheter, for, if the tube is very dilated and the membrane flaccid, rupture of the latter may follow when too much pressure is used. In those cases in which the tubes are freely pervious, and improvement in the hearing does not follow inflation, there is no object in continuing such treatment. Instead of using air alone through the catheter, certain vapors are to be preferred, more particularly a mixture of chloric ether and iodine (Blake). This consists of: tincture of iodin. gtt. xv; chloroform., spir. vini rectif., āā ʒss. Five or six minims of this mixture should be dropped into the Politzer bag through the hole in the side, which is then covered by the thumb, and the bag is com-

pressed. The catheter, before introduction, should be connected with the Politzer bag by means of a piece of rubber tubing at least a foot long, and then should be warmed and smeared over with vaseline. It is important for the surgeon to determine, by means of the diagnostic tube, the exact condition of the Eustachian tube. When the vapor is forced into the ear, the patient usually experiences a cool and pleasant sensation. In some instances the hearing-distance is decreased immediately after this procedure, but is improved several hours afterward. When the ears are inflated by Politzer's method, and it is desirable to use the vapor just mentioned, a glass inhaler should be attached to the Politzer bag. A few drops of the mixture should be poured on absorbent cotton placed in the inhaler. Some surgeons recommend to their patients the use of a large inhaler in which chloride of ammonium or other fumes are generated. After filling the nasal passages and nasopharynx with the vapor, they are directed to perform the Valsalvian experiment. The principal objection to this method is that the patient is likely to inflate the ears too frequently, and thus eventually produce a relaxed condition of the membrana tympani. Vapors of sal ammoniac, acetic acid, turpentine, the different balsams, as well as hydrogen gas and iodide of ethyl, are also used. George Stoker,[1] of London, has used ozone in cases of chronic catarrh of the middle ear with tinnitus. He has devised an apparatus for generating the ozone and by means of a catheter it is applied directly to the middle ear every day for four or five minutes at a time for at least six weeks or longer. He claims that in cases of nasal catarrh, it

[1] Annales des Mal. de l'oreille, July, 1905.

reduces the hyperæmia and renders the secretion normal. Beside vapors, some otologists recommend the injection through the catheter of various solutions into the middle ear.

Politzer has seen "favorable results follow in a series of cases from the injection of five or six drops of a 2 per cent. solution of pilocarpine." The catheter is first introduced and fastened by means of a clamp. A small syringe containing the solution is warmed, and a few drops are then injected into the catheter. By means of the Politzer bag the solution is forced into the middle ear. Liquid vaseline (Ch. Delstanche) or albolene is sometimes used in the same manner as the pilocarpine solution.

Some benefit follows from the use of albolene containing 2 per cent. of pilocarpine. If a few drops only are injected into the middle ear, there is not likely to be any discomfort to the patient. As the hearing immediately after the injection is frequently made temporarily worse, it is better to vary the treatment by simply forcing air through the catheter every other time. It is occasionally advisable to treat such a patient every day for a week or ten days; but, as a general rule, it is better to inflate the ears not oftener than three times a week. The patient should discontinue treatment after six weeks' time, and have a period of rest. After the first two weeks it may be necessary to catheterize the ears but twice a week. A very mild solution of bicarbonate of soda is recommended by some otologists, and is not likely to cause much irritation of the mucous membrane of the tympanic cavity. Solutions of borax, bichloride of mercury, and iodide of potash, which are recommended by some, are contraindicated, since they are liable to cause considerable reaction.

Some authorities claim that hypodermic injections of strychnia in the neighborhood of the mastoid process have been followed by improvement in the hearing and diminution of the tinnitus.

Good results have of late been claimed from the use of thiosinamin, which is derived from the volatile oil of horse-radish or more commonly from the volatile oil of mustard seed. The dose, according to Hare is 4–6 grains given once a day. Sugar[1], of Budapest, uses it in cases of chronic otitis with adhesions and cicatrices, and in cases of sclerosis where there is connective-tissue binding the oval and round windows with ankylosis of the stapes but where there is bony ankylosis he prefers preparations of phosphorus. He gives it hypodermically. Hirschland[2] claims to have had good results from the administration of this remedy. In some cases the hearing has been improved while in others the hearing has not improved but the tinnitus has become less and the giddiness has disappeared. He says that the drug produces a redness of the drumhead and after a short time the membrane appears sodden and by catheterization one gets the impression of fluid. The improvement is not marked for the watch but for speech. He has used this remedy in a great number of cases of deafness due to chronic catarrh and in cases of suppurative otitis media where the discharge has ceased and there is a dry perforation left. He gives it hypodermically in a Pravaz syringe half filled with a 10 per cent. glycerin and water solution, and generally increases the dose. His results show that it is a useful remedy in some cases, is harmless and in a small number of cases it has rendered good service.

[1]Archiv. f. Ohrenheil, 62 Bd., 3u 4.Heft.
[2]Archiv. f. Ohrenheil, 64 Bd., 2 u 3. Heft.

McCullagh[1] says that thiosinamin exerts a markedly beneficial action in aural cases where there is a formation of new connective tissue. Its administration should always be combined with the usual forms of local treatment, catheter, etc. Equally good results, he claims, may be obtained from its administration by the mouth as hypodermically.

The author has been using this remedy in a number of cases but sufficient time has not elapsed as yet for him to express any definite opinion as to its value in otology.

Schwabach has used ergot in 10 drop doses three times a day for tinnitus with some good results. Denker[2] considers that phosphorus given internally as suggested by Siebenmann has a favorable effect on the course of the disease. It must be continued for two or three years.

Siegle's otoscope and the "rarefacteur" or "masseur" of Delstanche are useful instruments in many cases, and especially where there are recent adhesions and retraction of the drumhead. It is possible to rupture the drumhead as well as to cause ecchymoses and inflammation, if too much force is employed when the air is exhausted, so that the surgeon should always inspect the membrana while using such an instrument. In cases in which there are atrophic changes, as shown by bulging outward of certain portions of the drumhead when the ear is inflated, one should be very cautious when trying this method of treatment, for the membrane may simply be rendered more flaccid without even improving the hearing.

[1] Archives of Otology, June, 1905.

[2] Die Otosclerose. Erlangen, 1905.

A massage apparatus driven by an electric motor is used by some surgeons, the object being "to exert a gymnastic influence by rhythmical oscillatory movements of the drumhead and ossicles, in order to obviate the progressive stiffness and ultimate anchylosis of those parts."[1] Of the therapeutic value of vibratory massage of the drumhead, Schwabach states that it does but little good in cases of sclerosis of the sound-conducting apparatus. He recommends its use in cases of troublesome tinnitus, in which the patients are unable to attend to their business, in cases of simple chronic hypertrophic middle-ear catarrh with dulness and retraction of the drumhead, likewise in subacute middle-ear catarrh, also in cases of acute otitis media due to influenza, and in the sequelæ of chronic purulent otitis media, in which by the ordinary treatment no improvement in the subjective noises and hearing-distance could be obtained. In these cases the author asserts that "massage shows itself to be of essential value."

Urbantschitsch believes in the efficacy of massage of the side and back of the neck in cases of tinnitus aurium. In patients suffering from labyrinthine deafness, according to the same author, much benefit follows massage of the regions supplied by the auriculo-temporal and supra-orbital nerves. He further recommends massage of the Eustachian tube, which is effected by passing a bougie beyond the isthmus and then moving it alternately backward and forward for several seconds, and in some instances for even a minute or more. He has recently devised an electrical instrument for applying friction massage, which is worked by means of a motor. He uses celluloid bougies.[2]

[1] Schwabach: Archives of Otology, vol. xxx. Nos. 4 and 5, 1901.
[2] Monatschr. f. Ohrenheil., May, 1905.

A method devised by Hommel consists in instructing the patient to press the tragus into the external meatus, and then, after removal of the finger, to allow the tragus to return to its place. In this way the air in the canal is alternately condensed and rarefied. The patient should be advised to repeat these manipulations three or four times a day, and on each occasion he should press inward the tragus about fifty times.

Lucae has devised a pressure-probe, consisting of a probe with a cup-shaped cavity on the end, which is applied directly over the short process. Gentle and intermittent pressure is made on the ossicles by means of this instrument, which moves on a spring. Although considerable increase in the hearing-distance and an improvement in the tinnitus often follow its use, such improvement is not likely to be permanent. With the same object in view, various electrical instruments have been invented, by means of which vibrations are conveyed directly to the ossicular chain through a speculum constructed for the purpose. Although opinions differ as to the benefit derived from such treatment, these instruments should be given a fair trial. Patients are advised often by their physicians to make use of the Valsalvian method of inflation, and, as a result, there is a liability that the drumhead will become very much relaxed.

In certain cases in which there is probably a relaxation of the tensor tympani, as well as of the malleo-incudal capsular ligament, and the corresponding attachments, such as would favor a separation of the malleo-incudal articulation, Blake employs a *rubber spring*, which he introduces into the canal, and applies

directly to the neighborhood of the short process, and which acts as a support to these relaxed parts. In a paper on this subject, he reports a case of a musician whose hearing in the right ear was normal, while in the left ear there had been in childhood a suppurative process and a subsequent perforation in Shrapnell's membrane, which was dry at this time. The hearing of the left ear was about two-thirds of the normal distance. He had had a series of head-colds during the past winter, accompanied by a sense of fulness in the left ear, which had led him to practise the Valsalvian inflation frequently for the sake of the relief it afforded. Three days before his visit he had noticed, while practising on the violin, that when the head was bent forward and to the left, as is often the custom in the rendering of more delicate passages, the tones of the violin sounded dull in the left ear, and there was a definite loss of timbre perception. When the head was raised, this disability disappeared, to recur immediately on lowering the head to the position mentioned. An examination showed the membrana tympani to be intact, except that there was a small dry perforation in Shrapnell's membrane. The testimony of the patient was corroborated by tests made with tuning-forks and high musical instruments. There was considerable movement of the membrana tympani outward when the ear was inflated, such as would favor a separation of the malleo-incudal articulation "within such limits as would permit the passage of the larger sound-waves of the medium and lower tones, but would result in the loss in transit through the articulation of the motion incident to the shorter sound-waves of the qualitative

overtones." Blake says "that any mechanical arrangement which would bring about a more natural adjustment of the malleo-incudal articulating surfaces, and would maintain them in that position long enough to allow the relaxed tissues to normally contract, would not only relieve the patient of his recent embarrassment, but would probably have more permanent beneficial effects." Such pressure on the short process was effected by means of a strip of thin, elastic rubber tissue 3 mm. in width and 12 mm. in length, which, being seized at the two ends with angular forceps, was carried into the ear under good illumination, with the convex portion of the resultant ellipse in contact with the short process, when, the two ends being released from the grasp of the forceps and allowed to spring outward against the walls of the canal, pressure was exerted upon the short process. With such support to the relaxed parts, it was found that the head could be placed in any position without impairment in hearing for overtones, and the patient was enabled to resume his professional work. Blake has made a change in the preparation of the rubber strip, owing to the fact that the freshly cut ends of the rubber were found to adhere under the pressure of the forceps, thus preventing the springing apart of the ends when released. The ends should, therefore, be cut, one at an angle and the other on a curve, the curved end being placed toward the posterior wall of the canal and the convex surface of the rubber strip; secondly, that portion which was to be brought into contact with the malleus was lightly smeared with vaseline before introduction. It was found that absorption of the vaseline produced a partial

vacuum beneath the rubber, and resulted in so close an adherence of the latter to the skin that the beneficial effects of the spring were of much longer duration. In some cases, where there is a considerable relaxation of the membrana, Blake substitutes the *paper dressing* (see page 283).

The use of the rubber strip is of benefit in cases of circulatory tinnitus marked by relaxation of the drum-head and the malleo-incudal joint, as shown by the report of such a case. The treatment must be long continued in many instances in order to get any permanent results, while the adjustment of such mechanical appliances must always be a matter of more or less experiment. A slight change in adjustment may bring about a very decided improvement.

NARROWING AND STRICTURE OF THE EUSTACHIAN TUBE.

The Eustachian tube may be constricted through swelling of the mucous membrane, or by the formation of connective tissue, which leads to contraction and stricture. The constriction of the canal is usually the result of chronic catarrh, but may be due to the pressure of a growth in the nasopharynx, and is generally situated in the cartilaginous canal, although osseous growths in the bony canal may lead to stricture of this portion. The presence of a stricture is diagnosed usually by means of the auscultation tube when Politzer's method or catheterization is attempted. Under such conditions no sound is likely to be heard, or else there is a sharp, hissing or whistling sound. There is usually considerable deafness with a distressing tinnitus, a disagreeable

sensation of fulness or heaviness in the head, and occasionally dizziness.

In cases of constriction of the tube due to swelling of the mucous membrane, an attempt should be made to relieve this condition by treatment directed to the naso-pharynx, and by inflation of the middle ear through the catheter. When the stricture is not overcome by such measures, bougies should be tried, and for this purpose those made of silkworm-gut, covered over with lacquer and conical in shape, are the best; or, if they have not sufficient resisting power, whalebone bougies, well polished, rounded, and conical shaped, should be substituted. The bougie should be dipped in liquid vaseline and inserted through a good-sized silver catheter as far as the bony portion of the Eustachian tube. A mark should be made on the bougie so that the surgeon may know the distance that it has been introduced. Care should be taken that the point of the bougie does not injure the drumhead. The instrument should be left in position five or ten minutes. In some cases Politzer uses catgut bougies that have been soaked in a solution of nitrate of silver, while in others he prefers a thin rubber capsule (that is, when the constriction is due to swelling of the mucous membrane), which is attached to the top of the catheter by a thread. By forcing air through the catheter the capsule is inflated. It is claimed by some that by means of electrolysis the calibre of the tube is frequently restored. As a general rule, the results obtained by the measures just described are not very promising, so that they have been discarded by many otologists. In some cases the use of the bougie increases the deafness and the subjective noise, while in other cases, even after successful dilatation

of the Eustachian tube, no improvement follows. Great care should be exercised not to lacerate the tissues when passing the bougie, for fear of inducing surgical emphysema, especially when subsequently inflating the ear. In rare instances the point of the bougie has been broken off in the Eustachian tube, so that the possibility of this accident should be borne in mind.

Operative Treatment. Ever since the time of Sir Astley Cooper, various attempts have been made by surgeons to maintain a permanent opening in the drumhead, or else to excise one or more of the ossicles for the cure of the deafness and subjective noises. *It may be stated in general terms that such operations, performed for the cure of chronic catarrhal otitis media, have been unsuccessful,* although in a very few instances considerable improvement has followed. Many cases have been made *very much worse* by operative interference, and the statement made by some otologists, that the operation of excision of the ossicles will not make matters worse, even if it is not followed by some improvement, *is incorrect.* In a case which seemed to be an ideal one for the removal of the malleus and incus, as these ossicles were bound down by adhesions, the operation was not followed by any improvement to speak of, but, instead, a distressing tinnitus developed. A very nervous patient who submitted to an operation of excision of the entire drumhead, together with the malleus and incus, was assured that the deafness would not be more marked, even if no improvement followed. On recovering from the effects of the ether, the hearing was found to be much worse, and at the present time she cannot even hear with the aid of a conversation tube. The hearing-distance in the other ear also has decreased very much as

a result of the operation, a phenomenon that is often observed in such cases. Numerous complications have followed the performance of excision of the drumhead and ossicles, such as purulent otitis media, mastoid empyema, burrowing abscess of the neck, etc., so that these accidents should always be borne in mind when a patient seeks advice as to the propriety of such an operation.

Cooper and others obtained brilliant results at first by making an artificial opening in the drumhead, but were obliged to abandon the operation, because the aperture soon became closed by newly formed connective tissue, and the deafness was not improved. At first, as suggested by Cooper, a portion of the drumhead was excised. This was followed by an operation devised by Wreden, in which he excised a portion of the long handle of the malleus. Later, an attempt was made by means of nitrate of silver, sulphuric acid, and the galvano-cautery, to produce a permanent opening. Such openings are apt to close very soon, and although the hearing-distance may be increased and the subjective noises lessened for the time being while there is an opening in the drumhead, nevertheless, when the latter closes, both noises and deafness may be very much worse. The object of making an artificial opening is for the relief of deafness and distressing tinnitus in cases in which the ossicles are bound down by adhesions; when the membrana is very much thickened or else contains calcareous deposits; when there is a stricture of the Eustachian tube which cannot otherwise be overcome; also in exploratory operations (to be described more fully), as well as for the purpose of mobilizing the ossicles, especially the stapes. As the object of such an

operation is to allow the sound-waves to strike against the foot-plate of the stapes and reach the labyrinth immediately, it stands to reason that such an operative procedure will be unsuccessful when there is anchylosis of the stapes, or thickening and calcification of the membrane of the foramen rotundum, or implication of the labyrinth. The operation should not be considered for a moment unless there is excellent bone-conduction. Politzer has devised an eyelet of gutta-percha for insertion in the artificial opening, but it has been found impossible to keep this in place.

When the drumhead is very much retracted, the long handle foreshortened, and the posterior fold stands out

Fig. 79.

Handle of malleus foreshortened; anterior and posterior folds of the drumhead very prominent.

prominently, and there is also considerable deafness, with loud subjective noises, which it is impossible to relieve by other treatment, Politzer recommends *section of the posterior fold of the membrana tympani* (see Fig. 79). The incision is usually made just behind the short process, by a small knife, rounded at its point and especially adapted for the purpose.

DIVISION OF THE TENSOR TYMPANI MUSCLE.

Tenotomy of this muscle, the object being to allow the drumhead to regain its normal position, and thus relieve the increased pressure within the labyrinth, was performed first by Weber-Liel, although Hyrtl previously had suggested such an operation. In cases in which retraction of the drumhead is due to this cause alone, such an operative procedure is indicated, but, unfortunately, these instances are infrequent. One is much more likely to find not only shortening of this tendon, but also adhesions in the tympanic cavity, with a shrinking of the folds of the mucous membrane and of the ligaments.

In order to divide this tendon, an incision is first made in the drumhead either in front of or else behind

FIG. 80.

Gruber's tenotome.

the handle of the malleus, preferably in the latter position. An instrument (see Fig. 80) curved on the flat is then introduced through this opening for a distance of at least 3 mm., the tenotome being directed below the tendon of the tensor tympani. By depressing the handle so that the point of the instrument is pushed upward, the tendon is generally divided when the blade is withdrawn. The results obtained by this operation have not been successful, as a general rule, so that it has been abandoned practically, as cases have been reported where considerable increase of the deafness has followed its performance.

Kessel, in 1871, advocated removal of the drumhead and excision of the malleus and incus, and since that time surgeons in this country and Europe have recommended different operations of a similar character, for the relief of extreme deafness, vertigo, and tinnitus aurium in cases of chronic catarrhal otitis media. Some otologists excise the incus alone, while the latest operation to attract attention has been the removal of the stapes, or *stapedectomy*. In cases of suppurative otitis media of a chronic nature, in which the malleus and incus are carious, surgeons are agreed as to the propriety of removing these bones on general surgical principles, and the benefit derived from such an operation has clearly been proved. In cases, however, of chronic non-suppurative inflammation of the middle ear, in which there is obstruction to the sound-transmission through the chain of ossicles, owing to thickening of the mucous membrane of the middle ear, adhesions, and anchylosis, *such an operation as removal of the drumhead with the malleus and incus is uncalled for if there is bony anchylosis of the stapes.* As Blake has expressed it, "the stapes holds the key to the situation," and it is undoubtedly true that in many cases which have been followed by improvement after excision of the malleus and incus, this change for the better has been due to mobilization of the stapes. Kessel further removed the columella in birds without causing disturbance of co-ordination, and demonstrated the fact that escape of the perilymph did not cause permanent impairment of hearing. Operations for the purpose of mobilizing the stapes have been advocated by different surgeons ever since Kessel, in 1875, reported a case in which he practised "circumcision" of the stapes, after dividing

the tendon of the stapedius muscle. He expressed an opinion that still further improvement would have followed had the stapes been extracted and a membrane been allowed to form in the foramen ovale, which would have vibrated more readily than the stapes that had been mobilized. Blake has reported a series of cases in which he removed the stapes, the various steps in the operation being as follows: "The ear to be operated upon should be first carefully tested as to its hearing-power, both by aërial and bone-conduction, and with due reference to the possibility of sound-transmission, either by aërial or bone-conduction, to the other ear. The external auditory canal should be cleansed of cerumen and loose skin, and then washed with a weak bichloride solution on a cotton-tipped probe and stopped with antiseptic cotton until the time of the operation." The instruments used are a paracentesis needle, a small angular knife for the division of the articulation of the incus and stapes, a small blunt hook, and a pair of light but strong angular forceps, to which may be added a small rectangular knife, to be passed between the incus and the stapes, after division of the joint, and a pair of very fine, straight-tipped forceps for the removal of a portion of the bone if fractured in the attempt at extraction. The surgeon should be provided with a number of cotton-tipped probes, a bowl containing a solution of bicarbonate of soda; also vessels containing a saturated solution of boric acid in alcohol, and a 4 per cent. solution of cocaine. The instruments should be sterilized thoroughly, and the hands of the operator scrubbed and rendered aseptic. Before using the instruments they should be dipped in the alcohol and boric acid solution. If possible, the operation should be per-

formed under conditions of local anæsthesia, so as to "permit of the conscious and undisturbed co-operation of the patient." Just before the operation a few drops of a 4 per cent. solution of cocaine are injected into the middle ear through the Eustachian catheter. The patient is placed in a high-backed chair, his head being firmly held by an assistant. An incision is then made see Fig. 81) "beginning at a point midway between

Fig. 81.

the short process and the tip of the long process of the malleus and close to the manubrium, then extending upward along the posterior ligament of the short process and following the periphery to a point posteriorly on a line with the tip of the manubrium." A large flap falls downward and outward, leaving an opening which gives free access to the subsequent field of operation. Any bleeding from the edges of the cut is controlled by the application of the sterilized cocaine solution. The hearing should then be tested and the operation continued, by dividing the tendon of the stapedius muscle with the short curved knife or an angular one, "a large angular knife being subsequently passed behind the incus to insure separation." At times it is advisable to remove the incus at this step in the operation.

By means of a probe and blunt hook, after testing the hearing, one can then decide whether the stapes shall be mobilized mechanically by dividing the folds in the

niche or by circumcision, or whether it is better to extract it. If the former has been decided upon, the region of the fenestra ovalis should be painted with cocaine, and an instrument similar to the one devised by Kessel or Politzer should then be introduced between the crura and the lower wall of the niche, and an incision made from right to left. If this is not followed by decided improvement, an attempt should be made to stretch the adhesions between the anterior and posterior crura and the walls of the niche.

Before extracting the stapes, it should be mobilized. A blunt hook, curved very slightly backward, is then passed behind the head and between the crura, either from below upward or in the opposite direction. The resistance offered when traction is made varies in different cases. At the moment of removal, however, the *pulse* often becomes *slower* in frequency, but occasionally is accelerated, and at times there is *vertigo*. Hemorrhage sometimes follows division of the stapedius muscle, but this can usually be controlled by a cotton-tipped probe. Sterilized cotton or gauze should then be inserted in the meatus and the patient kept quiet for a few days.

The results following extraction of the stapes have been decidedly *unsatisfactory*, and *the operation of stapedectomy* has generally been *condemned*. Of twenty-one cases reported by Blake, in only three was there any appreciable improvement following the operation, and in but two of them was this gain sufficient to be of material advantage. On the other hand, cases have been reported by other observers in which *alarming symptoms* followed this operation, such as *intense vertigo, facial paralysis,* and *deafness*. Before resort-

ing to mobilization or excision of the malleus and incus, or removal of the drumhead, every other method of treatment should first be tried, while operative measures should never be considered if there is labyrinthine disease. Moreover, the surgeon should explain to the patient the possible *dangers* of any operative procedure, and he should never make promises as to any permanent improvement in either the deafness or subjective noises. If the patient clearly understands the situation, and is anxious and willing, under these circumstances, that an operation on the membrana tympani should be performed, the surgeon should first of all do an "exploratory tympanotomy," by making an incision in the drumhead exactly similar to the one already described (see Fig. 81) in the operation of stapedectomy. Thus the area of the tympanum is exposed at a point where surgical access to it is most desirable, while an incision made at this point affords a stiff flap, the edges of which can be brought together and held in position by a paper dressing until the healing, which is usually complete in three days, has been effected. The formation of a cicatrix opposite the niche in the posterior-superior segment is also avoided. Another method of exposing the articulation of the incus and stapes is shown in Fig. 82.

In some persons it is necessary to administer an anæsthetic; but, if possible, the operation should be done by injecting a few drops of cocaine into the middle ear through the Eustachian tube or else by the method described by Blake, as follows: "In a series of experiments on the progressive growth of the dermoid coat of the membrana tympani, made several years since, it was found that the movement outward from

the centre of the membrane at the tip of the malleus toward the superior and posterior periphery was slowest in the inferior third of the posterior segment (A) (see Fig. 83), more rapid in the middle third (B), and most rapid in the upper third (C); and that the rate of move-

FIG. 82.



Triangular incision in the posterior-superior quadrant for exposing the articulation of the incus and stapes. (POLITZER.)

ment, as shown by small pieces of paper pasted upon the membrane, bore a proportionate relation to the corresponding degree of vascularity of the three sections mentioned; while a recent series of tactile examinations shows that the sensitiveness of the posterior segment of the membrana tympani has the same relative correspond-

FIG. 83.

ence, and therefore that an incision begun opposite the round window causes increasing pain as it is extended upward and forward along the periphery." First a preliminary incision 2 mm. in length is made close to the posterior periphery, opposite the round window, and to

this a 2 per cent. solution of cocaine should be applied. After waiting a minute, the cut should be extended upward in the direction already described, and cocaine applied as before. By this method the incision can gradually be extended, without much if any pain, to a point opposite the centre of the posterior ligament of the short process. Tests of hearing should now be made, so that the surgeon can determine as to the advisability of further operative procedure. He may decide to divide the tendon of the tensor tympani and mobilize the malleus, or he may divide adhesions which bind the malleus and incus to the walls of the tympanum, or he may mobilize the incus or stapes, or he may conclude to excise the malleus or incus, or both of these. *Operations performed under the most favorable circumstances,* although followed by temporary improvement at times, *do not offer much hope to the patient,* if he is a sufferer from chronic non-suppurative otitis media. Surgeons generally are opposed to any radical operation on the membrana tympani or ossicula in this disease.

OTALGIA.

In this affection, an examination made with the head-mirror and speculum will show an absence of every sign of inflammation of the auditory canal or drumhead. The pain usually is referred to the ear; but it is reflex, and most frequently due to the irritation caused by the presence of decayed teeth. It may also arise from some laryngeal irritation.

The pain may be localized in the external or middle ear, being due to an affection of the sensory nerve or nerves supplying the ear, the most frequent causes of such neuralgia being hysteria, cold, inflammation of

the nerve-trunks, and pressure on them from tumors of the brain or from caries or necrosis of the temporal bone. At times otalgia is connected with neuralgia of the fifth nerve. Neurasthenic patients who have chronic catarrhal otitis media occasionally complain of pain over the mastoid process which is neuralgic in character.

CLICKING TINNITUS.

This is an objective form of noise which is often audible to the surgeon as well as to the patient. In some instances it can be heard a few feet from the patient, while in other cases it can be detected only with the aid of the diagnostic tube. This clicking noise, which has sometimes a metallic sound, can at times be produced at will by the patient, but it is not generally under his control. It is an infrequent form of tinnitus which may cease at times, only to return. The opinion is held by some otologists that the noise is due to spasmodic action of the tensor tympani or stapedius muscle. It is more probable that it is caused by a spasmodic contraction of one or more of the palate muscles, whereby the mouth of the Eustachian tube is opened and shut, particularly if the mucous membrane is sticky from the presence of mucus. The treatment consists in relieving the catarrhal condition and the administration of tonic remedies, especially strychnine.

CHAPTER IX.

OTITIS MEDIA PURULENTA CHRONICA, OR CHRONIC PURULENT OTITIS MEDIA, OR CHRONIC SUPPURATIVE INFLAMMATION OF THE MIDDLE EAR.

This disease is one of the most important ear affections that claims the attention of the otologist, and is met with very frequently among the poorer classes, owing to neglect of treatment, for there is on the part of a great many persons a strong superstition, as it were, against "checking any discharge from the ear," for fear that some serious disease will attack another part of the body. Beside, the advice is often given to patients who are suffering from this affection that they will "outgrow" the discharge, so that, as a result, the surgeon often finds great destruction of the drumhead, a carious condition of the ossicles and temporal bone, and occasionally some intracranial complication. Of two hundred and thirty-four patients suffering from chronic purulent otitis media, large granulations or polypi were found in the middle ear or external meatus in twenty-four cases, the discharge being of long duration, offensive in character, and associated with more or less caries of the ossicula and walls of the tympanum. Among the most frequent causes, beside cold and influenza, were scarlet fever, diphtheria, whooping-cough, measles, teething, meningitis, tuberculosis, nasopharyngeal catarrh, and adenoid growths.

Since scarlet fever, measles, diphtheria, and the influenza are responsible for most of the cases of acute as well as of chronic purulent otitis media, it is of the greatest importance that the ears should frequently be examined during the course of one of these diseases. Prompt and early treatment will often effect a cure. Neglect of treatment, or "letting the ears take care of themselves," results, as a rule, in chronic purulent otitis media, with caries of the ossicula and other complications.

If a free incision is not made and proper drainage from the middle ear established in a case in which muco-pus is contained within the tympanum, there is then great danger, especially in the exanthematous diseases, that the acute inflammation will become chronic. The pus, not having a free exit, forms caseous masses in the attic and antrum, leading frequently to inflammation of the mastoid cells.

Anatomical Appearances. The mucous membrane of the middle ear becomes much thickened, due to infiltration with round cells, and to enlargement of the bloodvessels and the formation of new ones. The epithelium frequently changes its character, so that, instead of being ciliated, it becomes epidermal, the epidermis extending from the external meatus into the tympanum. The mucous membrane is usually of a deep-red color, and granulations are often found on its surface, while in the deeper layers the lymphatic vessels are usually enlarged.

As a result of chronic suppuration, beside granulations and polypi that frequently develop from the mucous membrane, one notices connective-tissue bands that bind the drumhead and ossicula to the walls of

the middle ear. These new formations are permanent, as a rule. In some cases there are ulcerations in the middle ear, as well as a carious condition of the ossicles and tympanic walls.

There may be a small perforation of the drumhead, or the loss of substance may be so extensive, that a mere rim of the membrana tympani remains. The drumhead may or may not be swollen or thickened, with hypertrophy of the dermal layer. There may also be calcareous deposits between the layers or an excessive development of connective tissue. After cessation of the discharge, the perforation either remains open or else is closed by thin cicatricial tissue, which frequently adheres to the wall of the middle ear. When a rim of membrane remains in the upper part of the middle-ear cavity, the long handle of the malleus usually rests against the inner tympanic wall. (See Plate V., Fig. K.) The tympanic end of the Eustachian tube is generally swollen, and the glands are enlarged, so that frequently the calibre of the tube is diminished. Occasionally, however, but very infrequently, the tube is dilated.

Symptoms. If there is a free exit for the pus, the patient does not usually complain of any pain. If, on the other hand, there is an obstruction to the escape of pus, which is due to the presence of granulations or polypi, and if there is caries of the temporal bone, not only does severe pain occur, but unpleasant symptoms also are complained of, such as oppression and a feeling of fulness in the head and ear. The pus may be prevented from escaping on account of a small perforation, or there may be inspissated masses in the tympanum. Obstruction to the escape of pus may also give rise to

vertigo, nausea, vomiting, a staggering gait, great tinnitus, facial paralysis, and, in extreme cases, choked disk and retinitis. In such cases, when free drainage has been established and all granulations have been removed, these symptoms will disappear, unless the labyrinth has become involved or some intracranial complication has developed. Tinnitus is not usually so troublesome in chronic otorrhœa as in chronic catarrhal otitis media.

The deafness varies very much in the same individual, depending largely on the amount of secretion, the extent of the swelling of the mucous membrane, and also as to whether the Eustachian tube is patent or not. In some cases, especially of scarlet fever, diphtheria, typhoid fever, and syphilis, the deafness is very great. This is occasionally due to implication of the labyrinth, or to the ossicles being bound down by adhesions, or to anchylosis of the ossicula. As the mucous membrane is apt to be more swollen during damp and cold weather, the hearing is usually better when the atmosphere is warm and dry. The deafness is sometimes more marked after cessation of the discharge, a phenomenon which is probably due to the fact that the newly formed connective-tissue bands are more tense after the suppuration has ceased, since the hearing often is markedly improved when the discharge reappears and the connective tissue becomes relaxed.

Bone-conduction, as a rule, is increased in chronic otorrhœa, as shown by Weber's and Rinné's tests, except when the labyrinth is implicated.

Character and Amount of the Discharge. The secretion from the ear varies very much in its character and amount, being thin and watery in some cases and thick

and stringy in others. At first, when the inflammation has become chronic, and before granulations or polypi have formed or the bone has become necrotic, the discharge is apt to be yellowish, laudable pus. It is usually more copious in children than in adults. In some cases in which there is a nasopharyngeal catarrh, there are long strings of mucus, which occasionally have to be removed with forceps from the meatus. These cases are often obstinate to treatment. The discharge may also be of an acrid nature, so that the external meatus and auricle are apt to be considerably excoriated.

When blood-corpuscles are mixed with the discharge, it generally denotes the presence of granulations or polypi. The discharge varies in color from a light yellow to a greenish hue. When, however, it is brownish and has a fetid odor, the chances are strongly in favor of the existence of caries or necrosis of the middle ear or antrum, or possibly of the mastoid cells. A profuse discharge is significant of mastoid inflammation.

Appearances of the External Auditory Canal and Membrana Tympani.—When the ear is examined with a head-mirror and speculum, more or less secretion will be found in the external meatus or middle ear, or crusts, due to dried secretion, will be adherent to the inner end of the canal or drumhead. When the secretion is profuse, the dermal layer of the drumhead and external auditory canal will appear macerated, on removal of which the parts beneath will have a red and glistening appearance. One notices exostoses at times on the walls of the bony auditory canal, and the calibre of the latter may be considerably diminished if granulations on its wall coalesce and form bands. In order to secure a good view of the external meatus and drumhead, it is abso-

lutely necessary, first of all, to syringe the ear and remove all secretion and moisture by means of absorbent cotton applied on a cotton-carrier. Boric acid or bichloride of mercury should be used in the solution. The next step in the examination, after inspecting the external meatus, is to note carefully the condition of the drumhead, as well as the site and extent of the perforation. The latter is found most frequently in the anterior inferior quadrant, although it is observed often in the posterior superior quadrant. The perforation may be extremely small, or the entire drumhead may be destroyed, the latter condition being met with frequently in severe cases of scarlet fever and diphtheria. The hole is usually round or oval, but may be semilunar, kidney- or heart-shaped. There may be two or even three openings in the same membrana tympani, but one perforation is the rule. The kidney- or heart-shaped orifice is found generally in the lower part of the drumhead, with the long handle of the malleus dividing it in the centre. When there is considerable loss of substance the handle of the malleus will be in contact with the inner wall of the tympanum. In cases in which the perforation is extremely small, a diagnosis can be made by having the patient try the Valsalvian method, when a small drop of pus will be seen exuding through the aperture, or, by means of the diagnostic tube, the surgeon will be able to hear a perforation-whistle when the patient inflates the ear by the Valsalvian method, or the surgeon forces air through by Politzer's method or by catheterization.

When the perforation is a large one, the middle ear appears of a deep-red color, except in tubercular cases, when it is pale. As the Eustachian tube frequently is

dilated in these cases, a solution syringed into the ear may pass quite readily into the nasopharynx, hence bichloride of mercury should be used cautiously especially in children. When there is great loss of substance in the posterior half of the membrane, the incus and the incudo-malleal articulation come into view, and occasionally the foramen rotundum and the stapes in the foramen ovale. After cessation of the discharge the inner wall of the tympanum often becomes of a whitish-gray color. The drumhead, or whatever remains of it, varies in color from grayish to a dull red. When there is an unhealthy discharge the membrana appears thickened and the edges of the perforation are red and granular. After cicatrization has taken place, the margins of the orifice gradually become thinner and assume a normal appearance, and frequently become attached to the inner wall of the middle ear, in which case the hearing-power usually is decreased.

PERFORATION IN SHRAPNELL'S MEMBRANE.

A perforation in the membrana flaccida is occasionally overlooked, owing to the fact that there may be another perforation in the membrana vibrans. It is most important to remember in this connection that in about 70 per cent. of all cases there are certain folds or reduplications of mucous membrane, horizontal in position, in the upper part of the normal cavity, which interfere with the proper drainage of the tympanic attic. Beside these folds in the neighborhood of the attic, there are others irregularly placed about the round window and stapes, and still others, but principally striæ, very near the mastoid antrum. There are cases

in which great congestion and swelling of these folds take place, in which all communication with the mastoid cells is cut off. An examination of such a case will reveal an otherwise normal membrana vibrans, except that there is great congestion in Shrapnell's membrane and along the handle of the malleus. Such inflammation is observed occasionally in patients who are suffering from nervous exhaustion. The pain is apt to be very severe, and follows a feeling of fulness in the ear. The proper treatment consists in a free incision along the upper border of the drumhead from the short process carried backward and sufficiently deep to divide the layer of vessels that assist in forming the inner manubrial plexus. When the patient performs the Valsalvian experiment, or the surgeon inflates the ear by Politzer's method, the air does not, as a rule, pass through a perforation in the membrana flaccida.

The danger of inflammation in this portion of the drumhead consists in the fact that there is great difficulty in draining the attic, and, as the pus is likely to decompose rapidly, the result is that caries of the ossicles is liable to follow, with more or less loss of hearing. Unless *a free incision* is made at an early date in the drum membrane, the inflammation may extend to the antrum and mastoid cells, or a carious opening may occur in the thin plate of bone forming the roof, and give rise to epidural abscess or to a temporo-sphenoidal abscess, as happened in a case of mine, to which reference is made in the chapter on Brain Abscess. In a certain number of cases, perforation of Shrapnell's membrane is the result of a general purulent otitis media, the inflammation having ceased in the tympanic cavity,

but remaining obstinate in and about the attic. The perforation may be a small one or the entire membrana flaccida may be destroyed. As a result, loss of bony substance takes place, so that a considerable portion of the attic is disclosed. Granulations or small polypi are frequently found in this region. If the discharge is checked, the movement of the ossicula is much hampered, frequently by thickening of the folds of mucous membrane and by adhesions.

Diagnosis. In making a diagnosis of perforation, it is always important to touch with a probe the different parts of the drumhead and middle ear, not only to see if a carious spot can be found, but also, when cicatrices of the drum membrane have taken place, to determine whether there is adherence of the cicatrix to the bony wall. In this connection, Siegle's otoscope will also aid in determining this point. With the probe the surgeon can frequently decide whether there is anchylosis of the ossicles, and whether the small bones are bound down by connective-tissue bands. The disturbances of hearing are likely to be very marked in such cases, and especially when connective tissue is found about the fenestræ rotunda and ovalis. Connective-tissue bands sometimes become calcareous or ossified, in which case the degree of deafness becomes extreme.

Prognosis. The deafness, after cessation of the discharge, frequently remains stationary. In other cases the degree of deafness progresses steadily, which is due to increased rigidity of the newly developed connective tissue, or to anchylosis of the ossicles, or possibly to involvement of the labyrinth. As long as there is a discharge from the ear the patient should be impressed with the importance of having the ear properly cared

for, since neglect of treatment is likely to lead to some dangerous complication. The state of the general health is especially to be considered in regard to the prognosis. The latter is much more favorable when the disease occurs in a person of good health than in one suffering from tuberculosis, syphilis, the exanthematous diseases, diphtheria, typhus and typhoid fevers.

Unfavorable symptoms are a profuse discharge of long standing, especially if it has a fetid odor, constant pain, narrowing of the external meatus, drooping of the posterior upper wall of the external auditory canal, a brownish discharge from the admixture of blood-corpuscles, the presence of cheesy masses in the discharge, and granulations and polypi.

That tubercular disease is at times a cause of chronic suppurative otitis media is clearly shown in the report of the following case (Fig. 84): The patient, a salesman, thirty-five years of age, has had tubercular adenitis, for eleven years. The left ear has been discharging for about four years. Two years ago he awoke one morning to find the left side of the face paralyzed. There has been at times a slight improvement, and at present the paralysis is not so marked as when it first appeared. He has a large number of suppurative sinuses which appear to be connected with bone. He has never had any joint disease. He does not suffer from vertigo. An examination showed that the external meatus was considerably constricted, and that there was a profuse discharge. The drumhead was almost entirely destroyed. The facial nerve has evidently been involved within the tympanic cavity.

Treatment. To treat the ear properly, it is of the first importance (1) to remove thoroughly all secretion

from the tympanum and external meatus; (2) to establish as perfect drainage as possible; (3) to check the

Fig. 84.

After photograph of a patient suffering from chronic suppurative otitis media incidental to general tubercular adeno-osteitis.

discharge and bring about cicatrization of the perforation. One should, first of all, syringe the ear with

water that has previously been boiled or that contains boric acid in solution (10 grs. to ℥j), or carbolic acid (2 per cent.), permanganate of potash (5 per cent.), or aluminium acetat (2 to 3 per cent.), or bichloride of mercury (1 : 5000). The latter is indicated in all cases in which micro-organisms are present in the discharge, especially in cases of scarlatina, diphtheria, or influenza. There is one precaution that should be taken in regard to the use of bichloride of mercury in chronic otorrhœa, which is, that great care should be used in the case of children and also in all individuals in whom the Eustachian tube is dilated, for in such instances the solution is apt to enter the nasopharynx. It is well, therefore, in order to avoid this accident, to use but little force, and also to incline the patient's head toward the affected side. The ear should never be syringed forcibly, for fear of causing vertigo. After all secretion has been syringed out, the external meatus and middle ear should be dried as thoroughly as possible by means of absorbent cotton on a cotton-holder. The cotton should gently be pushed down to the membrana tympani and rotated, in order to absorb the secretion. When the perforation is large, the purulent discharge can be frequently forced out of the middle ear by means of Politzer's method of inflation, or by the use of the catheter, or it can be removed through exhaustion of air by means of Siegle's pneumatic speculum. Another method consists in forcing a stream of water, containing common salt or boric acid (grs. x to ℥j), through the Eustachian catheter and middle ear, especially if there are accumulations of caseous or cholesteatomatous masses in the tympanic cavity. The syringe should be attached to the catheter by means of soft-rubber tubing.

When the discharge is profuse, the ear should be syringed or douched at least three or four times a day; otherwise, less often. It is difficult for a patient to syringe his own ear satisfactorily. When entrusted with a syringe, the best one is that made of hard rubber, with a large blunt point. A soft-rubber bulb-syringe is also a perfectly safe one.

After thorough removal of all secretion, the next step in the treatment consists in the instillation of astringent solutions containing either zinc sulphate, copper sulphate, or lead acetate, three to eight grains to the ounce of water, with the addition of a drachm of glycerin, or Burow's solution may be used. The solution should always be warmed first, as cold is apt to cause pain or dizziness. When the perforation is small, the solution will frequently not enter the tympanum unless the head is held to one side and the ear is inflated by the Valsalvian or Politzer method, thus allowing the fluid to pass in when the air-bubbles escape. As thorough drainage is essential, it is much better, when the perforation is small and the discharge profuse, to enlarge the orifice by incising it. When this has been done, solutions can more thoroughly be applied to the middle ear. A solution that is especially beneficial in uncomplicated cases is one containing—

℞. Acid. boric.,	gr. xv;
Zinc. sulph.,	gr. viij;
Glycerin.,	ʒj;
Aquæ.,	ʒj.—M.

The solution should be instilled into the ear several times a day. When zinc sulphate does not check the discharge, then copper or lead should be substi-

tuted. If the perforation is large, and the surgeon can see the patient frequently, the so-called dry treatment is generally much more satisfactory than the use of drops. Frequent syringing of the ear often keeps up the discharge in many cases, so that it becomes necessary to resort to the so-called dry method, which consists in the insufflation of boric acid or some other powder into the middle ear, after the secretion has first been most carefully removed by syringing and the use of absorbent cotton on a probe. Finely powdered boric acid should be blown through a speculum into the middle ear by means of a powder-blower. The powder should never be packed in, for fear of preventing the escape of the pus, and thus inducing septicæmia. Beside boric acid, other powders are frequently used, such as iodoform, especially in cases of caries or syphilitic disease, or boric acid and iodoform mixed in equal parts, or equal parts of boric acid and zinc oxide, or xeroform. Iodol is sometimes substituted for iodoform, but it is not as efficient in its action. The ear should be cleansed and the powder insufflated once a day, and less frequently as the discharge becomes diminished in quantity. Occasionally, after several insufflations of powder, the discharge ceases. The dry treatment is not indicated, as a rule, when there is considerable mucus or when the perforation is small. After the canal has been filled with powder a piece of aseptic absorbent cotton should be placed in the meatus.

When granulations have formed and the mucous membrane is considerably swollen, great benefit will follow the *instillation of alcohol* into the canal. Being a stimulant as well as an antiseptic, its use is indicated particularly in cases in which the discharge contains

cheesy masses. Soft granulations shrivel and frequently disappear, while the alcohol forces its way into the various bony cavities. Pure alcohol should not be used where there is acute congestion or symptoms of mastoid tenderness; as it frequently causes severe pain, it should always be diluted with water at first, and gradually increased to its full strength. The alcohol drops should be used at first two or three times a day, and less often as the discharge becomes less. It is well to combine boric acid with the alcohol in the proportion of grs. xv to ℥j. When the secretion has a bad odor, iodoform or iodol should be substituted for the boric acid. It is generally considered best not to carry out this method of treatment when caries exists.

In cases in which there is a swollen condition of the mucous membrane, nitrate of silver (grs. x–xx to water ℥j) is sometimes used to advantage. The secretion in the ear should be removed first by syringing, and then application made to the middle ear by means of cotton on a cotton-carrier. The contraindications to its use are the presence of large granulations, a small perforation of the drumhead, and caries of the temporal bone. To prevent the discoloration of the skin of the meatus which is likely to follow its use, the surgeon should make an application of a solution of iodide of potash to the walls of the canal.

Urbantschitsch[1] has had excellent results from the use of thigenol in the proportion of thigenol 2 to 4 parts, glycerin and alcohol 10 parts each. He claims to have cured cases of chronic otorrhœa where other methods of treatment have failed. He also combines thigenol with boric acid in the proportion of 1–10.

[1] Monatschr., f. Ohrenheil., No. 4, 1903.

It is impossible, however, to tell at a first examination just what plan of treatment should be followed. After trying the dry method, a change to the instillation of astringent solutions or of alcohol will frequently bring about excellent results, and *vice versa.*

THE WICK TREATMENT.

This method consists in the introduction of a strip of boric acid or iodoform gauze, or a long, wick-shaped piece of absorbent cotton, into the external auditory canal down to the perforation in the drumhead. The gauze or cotton-wick should be changed several times during the day, if saturated with discharge. It is obviously necessary to syringe away all secretion and to dry the canal and middle ear, if possible, before introducing the gauze or cotton-wick.

When granulations spring from the drum membrane or the cavity of the middle ear, which do not shrivel up under the alcohol or dry method of treatment, and especially in cases in which the mucous membrane of the tympanum is studded with numerous excrescences which give it a granular appearance, it then becomes advisable to destroy these granulations by means of caustics or by the galvano-cautery, or by scraping them away with a sharp spoon or curette.

Caustics should not be applied except in those cases in which the granulations can easily be seen, while great care should be exercised not to touch the external meatus with the caustic solution. The application of liq. ferri sesquichloridi is sufficient in some instances to cause the granulations to disappear. The solution should not be used again until the eschar has come away.

Nitrate of silver or chromic acid, fused on the end of a probe, is indicated in some cases, and particularly the acid, which penetrates more deeply into the tissues, but which causes more pain. Before touching a granulation with chromic acid the mucous membrane should be swabbed with a 4 per cent. solution of cocaine.

Some surgeons prefer the galvano-cautery. It gives rise to more pain at the time of its application, but the granulations are destroyed more thoroughly, while there is usually no reaction. Scraping the granulations away with a sharp spoon (see Fig. 85) is one of the best

FIG. 85.



Politzer's sharp spoons.

methods, and instruments of different sizes and shapes have been devised for removing not only granulations on the drumhead or tympanic walls, but also those on the ossicles and in the attic. The walls, when carious, may also be scraped at the same time, a procedure which often hastens the cure.

When the perforation is in Shrapnell's membrane, the first indication for treatment is to drain the attic; and if the perforation is a small one, it should be enlarged so as to enable the surgeon to wash out the attic and Prussak's space with a canula and middle-ear syringe (Fig. 86). For this purpose, a 1 : 5000 bichloride solu-

tion is the best, or a solution containing carbolic acid. After thorough removal of all secretion or caseous masses, astringent solutions containing zinc, copper, or lead, should be injected into the attic by means of the middle-ear syringe, or alcohol and boric acid, or alcohol and iodoform, should be substituted according to the indications. Finely powdered boric acid is sometimes insufflated through a slender canula.

Fig. 86.

Blake's middle-ear syringe.

Haug[1] of Munich claims to have cured sixty-four out of ninety-eight cases of suppuration in the attic. He first enlarges the site of the perforation if necessary, so that a tympanic canula of large calibre can be introduced. The irrigations are made with a solution of permanganate of potash or boric acid. The cavity is then dried, and a solution of perhydrol 10 parts, in alcohol and glycerin 10 parts each, is slowly injected into the attic. The solution is retained for a quarter of an hour, with the head turned to the side. The

[1] German Med. Society, Archives of Otology, vol. xxxiv., No. 6, 1905.

canal is then dried and a small pledget of cotton, soaked in a strong solution of iodin-potassium iodid-glycerin inserted. The canal is then packed with gauze. The procedure is sometimes very painful. If necessary it is repeated in ten days. Haug has never repeated this procedure more than three times, because it is without avail if the case has not healed up in that time.

But little force should be used in syringing, as the patient is likely to complain of considerable vertigo. The apparatus (Fig. 87) devised by the late Dr. J.

FIG. 87.

Hewitt's apparatus for washing out the attic.

Hewitt is an admirable one for washing out the attic, and especially so after removal of the ossicles. The tubes are of glass and can readily be kept clean. By having a long piece of tubing attached to the bulb, the latter can be placed on the floor and compressed by the foot, so that both hands of the surgeon remain free.

If after several weeks' treatment by means of instillation of drops and the use of powders the discharge remains

unchanged and the odor is fetid, the surgeon should then remove the carious ossicles and scrape the attic. It is in the surgical treatment of these cases, in which the suppuration is kept up by a carious condition in the upper portion of the middle ear and the antro-attic and walls of the middle ear, that one is apt to obtain very brilliant results. When the suppuration is limited to the upper portion of the tympanum, a cure can undoubtedly be effected in a certain number of cases by excision of the carious ossicles and by scraping and curetting the attic. In most instances an improvement in the hearing follows.

There are, however, a large number of cases of long-standing suppuration where removal of the ossicles does not go far enough, and where it is much better to cut down on the antrum from behind the auricle and perforate the bone with chisels and mallet. (See Stacke operation.) Beside excising carious ossicles, one can by this means reach the antrum much more readily, and thus remove any cholesteatomatous matter, which so often collects in this cavity.

After such an operation, viz., excision of the ossicles, the discharge does not cease at once in the majority of cases. It is usually necessary to continue the treatment previously described—namely, washing out the attic with the middle-ear syringe and using solutions and powders—before a complete cure is established.

The operation of removal of the hammer and incus is indicated not only in cases of perforation of Shrapnell's membrane, in which the secretion has persisted in spite of long-continued antiseptic treatment, but also in those cases of chronic otorrhœa in which there is but a rim of membrane left in the upper part of the tympanic

cavity, and in which there is evidently some obstruction to the escape of the pus, which, becoming stagnant in the attic, causes caries of the ossicula and attic walls. Where cholesteatomatous masses form in the attic the operation is likewise indicated, but the surgeon will not be able to cure such cases unless he performs the Schwartze-Stacke operation.

Operation. The external auditory canal should first be syringed with a bichloride solution 1 : 5000, and afterward carefully dried with absorbent cotton applied on a cotton-holder. All instruments used should be thoroughly sterilized. As the operation is a painful one, it is usually best to anæsthetize the patient, although in some adults who are not of a nervous temperament cocaine may be applied locally. An incision should then be made in the drumhead along its upper border, close to the bony wall, and the attachments of the malleus, the ligaments, and the tensor tympani muscle carefully divided with a knife, as represented in Fig. 88. There is usually considerable hemorrhage, which impedes the progress of the operation. Syringing with hot water will frequently control this. If not, the canal should be plugged with strips of iodoform gauze and the operation suspended for a few moments. The blood should be wiped away frequently with absorbent cotton on cotton-holders, so that the surgeon can see the different steps of the operation. The head of the malleus should be disarticulated from the incus and then seized with a pair of forceps (see Fig. 89) and removed. When there are adhesions, these should be divided before attempting to extract the hammer. To remove the incus is a more difficult matter. If the long process can be seen, it should be seized with a pair of forceps, after the connec-

Fig. 88.



Author's knife for incising the membrana tympani and for dividing adhesions.

Fig. 89.



Politzer's forceps.

tion between the incus and stapes has been divided by a knife. When this is impossible, an incus-hook should carefully be introduced in the upper part of the tympanic cavity behind the incus, and the latter pulled downward and forward. An instrument that is well adapted for this purpose is the incus-hook devised by the late Dr. J. Hewitt, shown in Fig. 90. The hook is passed upward behind the incus, and then brought directly outward and downward. The incus is more apt to be carious than the malleus, and frequently the incus cannot be found. After excision of the ossicles,

the attic should be curetted to remove all granulation-tissue, great care being taken not to injure the facial canal and nerve. The external meatus should then be syringed with bichloride solution, a strip of iodoform gauze pushed down to the middle ear, and a piece of absorbent cotton inserted in the meatus.

Beside removal of the carious ossicles and scraping of the tympanic cavity, it is advisable in some cases in which there is caries of the bony wall just above and

FIG. 90.



Hewitt's incus hook.

bordering on the perforation in Shrapnell's membrane, to remove the diseased bone with a sharp spoon or by means of a pair of forceps especially designed for this purpose.

ATTENTION TO THE GENERAL HEALTH.

When the discharge from the middle ear has been checked (in cases of chronic suppurative otitis media), a child previously pale and anæmic often assumes a healthy appearance, so that it is important first of all to arrest the discharge, and particularly to remove all caseous masses, which, if absorbed, may lead to tuberculosis. I have seen the health of a tuberculous patient greatly improved after the removal of softened bone

and cheesy masses from the mastoid cavity. It becomes the duty of the aural surgeon not only to arrest the discharge and to improve the hearing, but also to inquire most carefully about the patient's general health. Errors of digestion should be corrected. Strumous children should be given cod-liver oil, syrup of the hypophosphites, or syrup of the iodide of iron. Many patients suffering from chronic otorrhœa are broken down in health or are sufferers from tuberculosis or syphilis. Change of climate is often desirable, especially for those afflicted with nasal catarrh.

The surgeon should never fail to examine carefully the nasopharynx, especially in children, to see if there are adenoid growths or enlarged tonsils.

Proper treatment of the nares and nasopharynx is indicated in all cases whenever required, by means of sprays, by the removal of spurs on the septum, etc. (See chapter on Nasal Diseases.)

ARTIFICIAL MEMBRANA TYMPANI.

Yearsley, in 1848, recommended the application of a ball of cotton to a drumhead that had been partially destroyed, for the improvement of the hearing-distance, while in 1852, Toynbee gave to the profession the artificial drum membrane which bears his name (see Fig. 91). It consists of a round disk of thin rubber about 6 mm. in diameter, attached at the centre to a piece of wire, by means of which the artificial drum membrane can be introduced into the canal and properly applied.

Politzer has devised another form of membrane, consisting of a piece of rubber 1 cm. long, which is cut from the side of an india-rubber tube 2-3 mm. in thickness. The lower end, being perforated, is fixed to

a moderately strong wire (see Fig. 92). This appliance, being inexpensive, is especially adapted for use among the poor.

A very simple device for improving the hearing consists of a piece of rubber tubing, which should be cut off to correspond to the length of the external auditory canal, and so rounded at the inner extremity as to rest against the remnant of the drumhead.

Hassenstein has invented a pair of forceps for grasping a finely rolled, wick-shaped piece of wadding.

Fig. 91.

Toynbee's artificial drum membrane.

Fig. 92.

Artificial membrana tympani for use in practice among the poor. (Politzer.)

Ch. Delstanche uses a ball or roll of cotton which has been twisted about a thin wire, and which the patient can be taught to make for himself. Toynbee's membrane is seldom used at the present day, as its application causes irritation of the external meatus and of the drumhead, and, besides, patients often complain of a disagreeable noise in the ear while talking or eating. The cotton ball or wick, on the other hand, can be used even when there is slight moisture, as this will be absorbed by the wadding. Borated cotton, or cotton

soaked in astringent solutions, or smeared over with vaseline, may be used according to the condition of the middle ear.

The patient should be instructed by the surgeon to introduce the artificial membrane by means of Hassenstein's forceps, or other instruments adapted for the purpose, and he will soon become very proficient in applying the membrane to the exact spot at which the hearing is improved.

The first application of an artificial membrane may not be followed by improvement, so that the surgeon should make several attempts before deciding as to whether it is indicated or not.

Different views have been expressed by surgeons as to the mode of action of the artificial membrane, some holding that it is simply necessary to close the perforation, while others believe the improvement in hearing to be due to the pressure exerted against the remnant of the drumhead and ossicles. The latter view is probably the correct one, as improvement in the hearing often follows when pressure is made just over the stapes. The artificial membrane should be tried in all cases in which local treatment has failed to improve the hearing-distance. It should never be used in the case of children or when there is much if any discharge. It is contra-indicated when it causes irritation or inflammation or dizziness.

As the artificial drum membrane is liable to cause a certain amount of irritation, it should be applied and left *in situ* for an hour or so at first, and the time gradually increased each day. The cotton should be changed as often as it becomes saturated with discharge; in such cases the ear should be syringed and dried

before it is reapplied. After the patient has learned the mode of application of the membrane he will be able to change it frequently himself. It is often advisable for him to do this each day.

In a case of perforation of the drumhead, cicatrization of the same may at times be brought about by touching the edges with nitrate of silver fused on the end of a probe, or by scarification of the drum membrane, while incisions made near the opening occasionally produce a similar result. Another procedure (Blake) consists in the application of a disk of paper a little larger than the perforation to be covered. Ordinary writing-paper is used for this purpose, because (1) any desired thickness can easily be procured, and (2) the sizing used in finishing the paper when moistened is sticky enough to make the paper adhere to the membrana tympani. Such a disk is best introduced by means of a probe tipped with cotton and dipped in water, so that the paper adheres to the moistened cotton until it comes in contact with the membrana tympani. The disk of paper should then be pressed firmly into position by a cotton-holder tipped with dry cotton. The paper dressing is useful in only a limited number of cases, viz., those in which the perforation is small, and the inflammation has subsided, and there has been cessation of the discharge, the external surface of the drumhead having resumed a normal condition. It may also be applied to advantage in cases of rupture or mechanical injury, the paper dressing keeping the edges of the wound in apposition. Such a disk of paper will remain *in situ* until either the perforation (a small one) becomes closed or until slight discharge from an inflammatory process washes it away.

In cases in which the defect in hearing seems to be dependent on a relaxed cicatrix, as shown by increased impairment of hearing when the cicatrix bulges out, and by an improvement when it returns to its position, an attempt is sometimes made to bring about contraction of this relaxed tissue by means of multiple incisions made in the cicatrix with a fine needle, or by painting it with collodion.

There is usually more or less disturbance of hearing not only during the course of the suppuration, but also after cicatrization has taken place. Improvement of hearing follows in some cases from the use of Siegle's otoscope, or, better still, Delstanche's masseur, especially when the deafness is dependent on adhesions that can be broken up by the rarefaction or condensation of air in the external meatus. The pressure probe (Lucae) is useful in other cases, although the improvement in hearing, that at times is very marked, does not, as a rule, continue. The probe consists of a cup-shaped disk, which is attached to a long handle, and which works on a spring. The disk is placed usually on the short process, and intermittent pressure is made by means of the spring.

DIVISION OF ADHESIONS. EXCISION OF OSSICLES.

As the result of adhesions due to chronic purulent otitis media, permanent disturbances of hearing remain; also, occasionally, distressing tinnitus or vertigo unless relieved by an operation.

The ossicles may be so bound together and to each other that nothing but a radical operation will afford any relief. No definite promise should be made to the patient as to any permanent improvement in the hearing or the tinnitus or vertigo, but it may be stated,

as a rule, that when the bone-conduction is better than the aërial conduction, and the ossicles are bound down by adhesions as a result of chronic suppurative otitis media, the hearing is generally improved by an operation. The nature of the adhesions should first be determined by means of Siegle's otoscope and a probe. With a suitable knife, band-like formations between the handle of the malleus and the incus and stapes should be divided, as well as adhesive bands between the ossicles and tympanic walls. In some cases the hearing shows a permanent improvement, while in others it is but temporary. If division of adhesions and mobilization of the stapes are followed only by a temporarily improved condition, and the patient is willing to undergo a further operation, then the malleus and incus should be excised. The different steps in this operation have already been described (see pages 277 and 278).

CHOLESTEATOMA AND EPITHELIAL MASSES IN THE TYMPANIC CAVITY.

During the course of chronic suppurative otitis media there are found at times in the secretion masses of epithelial cells, not only in the external meatus, but also in the middle ear. The patient may not suffer any particular discomfort, and these epithelial scales may be found in the discharge for years. If, however, there is any hinderance to escape of the pus, these masses often form a large plug and give rise to considerable pain and pressure symptoms, such as vertigo and vomiting. If not removed, they lead to ulceration of the bone or to an intracranial complication. When the disease is limited to the attic, there being a small perforation in Shrapnell's membrane, caries of the ossicula

is apt to occur, or the disease is likely to extend to the brain through the tympanic roof.

The term cholesteatoma is applied generally to a collection of epithelial cells, round or polygonal, arranged in concentric layers, and mixed with fat-globules and crystals of cholesterin. Sometimes the whole mass is covered by a pearly, shining membrane.

There is a great difference of opinion among pathologists as to the origin of these masses, some maintaining that a cholesteatoma in the temporal bone is of heteroplastic formation, while others consider it to be a retentive tumor. Still others believe these scales to be formed in the external meatus and to extend through a perforation into the middle ear and antrum. As opposed to this latter view, it may be stated that a case of cholesteatoma of the tympanum has been reported without any evidence of suppuration of the middle ear or perforation of the drum membrane. Such cases are extremely infrequent, however, as cholesteatoma is observed usually in cases of chronic suppurative otitis media.

A cholesteatoma may be found in the middle ear, antrum, or mastoid cells, the latter being the most frequent seat of the disease. In some cases the cholesteatoma is of small size, while in others it is very large, leading to more or less destruction of bone from pressure and absorption, or there may be extensive caries. The middle ear, antrum, and mastoid are sometimes transformed into one large cavity, or there may be considerable destruction of the margo tympani or of the posterior and superior bony walls of the external meatus.

Symptoms. The symptoms vary according to the size

and nature of the collection of scales. There is generally a feeling of fulness or heaviness, or there may be pain, headache, and vertigo. The cholesteatomatous masses may either escape through the perforation, or they may extend to the antrum and mastoid cells, and lead to mastoid disease or some intracranial complication, so that an operation becomes necessary. The pus not being able to escape, may give rise, through absorption, to septicæmia or pyæmia, or meningitis may follow, or cerebral abscess, or sinus phlebitis, or severe hemorrhage from erosion of the lateral sinus or carotid artery.

Diagnosis. It is very important to make an early diagnosis of cholesteatoma, as the disease is a destructive one, and radical treatment is required at an early date if the surgeon expects to cure the discharge. The diagnosis can only be made with certainty when the epithelial cells either in the external meatus or protruding from a perforation in the drumhead, are examined under the microscope. Fistulous openings occur in the bony meatus in some cases. When there is a considerable discharge of epithelial scales from the middle ear, associated with pain in the mostoid process, it is probable that the latter is involved, and an immediate operation is indicated.

Treatment. If there is a large opening in the drum membrane and the cholesteatomatous masses are not excessive, an attempt should be made to wash out the tympanum with a bichloride solution by means of the middle-ear syringe (see Figs. 86 and 87). When the perforation is small and is in Shrapnell's membrane, it should be enlarged and the attic irrigated in the same manner. This plan of treatment should not be continued for a long time unless decided improvement

ensues, on account of the complications likely to develop. If the patient complains of pain or dizziness, and the cholesteatoma is in the antrum or mastoid process, the operation of opening the antrum and cells should be performed at once (see chapter on Mastoid Diseases), so that the cholesteatomatous masses can be reached and scraped away with curettes. If the ossicles are found carious, they should be excised and the attic curetted, great care being taken not to injure the facial nerve. In certain cases, when the disease is limited to the middle ear, attic, and antrum, Stacke's operation is indicated. This operation is described in the chapter on Mastoid Diseases.

CHAPTER X.

GRANULATIONS AND POLYPI—CARIES AND NECROSIS OF THE TEMPORAL BONE.

GRANULATIONS AND POLYPI.

Granulations should be considered under the same classification as polypi, for they are observed generally in connection with chronic otorrhœa, and, beside, a granulation may develop into a polypus. Occasionally, granulations as well as polypi appear during the course of an acute purulent otitis media; but, as a rule, they are the result of long-continued suppuration from the middle ear, with or without caries. Large flabby granulations frequently conceal a carious spot either in the bony auditory canal or in the tympanic cavity, or a carious ossicle may be embedded in granulation-tissue. Polypi are attached generally to the mucous membrane of the tympanum, less frequently to the wall of the external meatus and to the membrana tympani. Cases have been reported in which polypi were found in the tympanic cavity without perforation, but such instances are extremely infrequent. When attached to the external meatus, the most frequent site is the posterior superior wall of the bony canal close to the membrana tympani. The importance of examining microscopically every growth removed from the external auditory canal or middle ear is demonstrated in the report of the follow-

ing case:[1] A woman, fifty years old, came to my clinic, complaining of slight tinnitus and deafness. An examination showed a tumor apparently filling the external meatus and attached by a pedicle to the posterior cartilaginous canal wall. There was no involvement of the middle ear. It was removed with a Blake snare, and perfect healing followed. The microscopical examination showed it to be a sarcoma.

There may be a perforation in Shrapnell's membrane with a polypus protruding through the opening. Polypi are found simultaneously in the middle ear, on the drumhead, and in the external auditory canal. They vary greatly in size, being at times extremely small, or sufficiently large to fill the external auditory canal and to protrude from it. The external meatus is liable to be enlarged by the pressure of the growth. A polypus may be attached by a slender pedicle or by a broad base. There may be but a single one, or there may be several. They vary in shape, being either smooth and club-shaped, or they resemble a raspberry, and have a knobbed and glandular appearance. They are bright red frequently, but when exposed to the air they generally lose their color and become pale and pinkish.

Four varieties of polypi usually are described, viz.:

1. Mucous polypi.
2. Fibromata.
3. Myxomata.
4. Angiomata.

The most frequent varieties are the mucous polypi and fibromata. Myxomata are very rare.

1. *Mucous polypi* are more frequently observed than

[1] Report of New York Otological Society, Archives of Otology, vol. xxix. No. 1, 1900.

any of the others, and histologically are identical with polypi found in the nares and nasopharynx. They are formed of a delicate and loose stroma of areolar connective tissue, in which are enclosed round-cells and occasionally a few spindle-shaped cells. These polypi contain numerous bloodvessels, and generally have an

Fig. 93.



Section of mucous polypus.

irregular surface, due to their papillary and glandular structure. Cysts, considered by some authors as retention-cysts, are found in these polypi. The epithelium on the inner surface is usually of the cylindrical or ciliated cylindrical variety, while that covering the surface exposed to the outer air is pavement epithelium.

A soft or mucous polypus may gradually become transformed into a fibroma by the bloodvessels becoming obliterated and by the cellular elements becoming changed into spindle-cells and connective-tissue fibres.

2. *Fibromata.* A fibroma is covered with pavement epithelium, and is composed of connective tissue with spindle-cells interspersed in its meshes. It has fewer bloodvessels than a mucous polypus, and in consequence is paler in color and does not bleed so readily when touched with a probe. It is of a firm consistency, but does not contain cysts and tubular glands. It is usually attached to the periosteal lining of the tympanum or external meatus, and its growth is much slower than that of a mucous polypus.

3. *Myxomata.* Myxomata are observed very infrequently. Histologically, the tumor consists of a multiple layer of pavement epithelium for its external covering, into which flat papillæ project. The stroma is of a homogeneous gelatinous tissue, and contains spindle and star-shaped cells. There are also round glandular cells with a round nucleus similar to lymph-corpuscles.

4. *Angiomata.* These polypi are very vascular, and consist of a network of blood-spaces which appear to take the place of capillaries. The blood enters these spaces directly from the arteries, and thence goes to the veins and thence into the general circulation. When examined under a microscope, the growth is found to consist of bloodvessels connected with a large and irregular central cavity containing fibrous connective tissue.

Although a polypus in the external meatus, as a rule, has its attachment to the osseous wall, it occasionally happens that a large polypus springs from the cartilaginous

canal, especially if there is a sinus at that point leading to carious bone.

A granulation or polypus often develops in the auditory canal from a point where a furuncle has existed. A polypus sometimes contains osseous deposits.

Symptoms. When granulations or polypi do not interfere with drainage from the middle ear, there may be no other symptoms except a little bloody discharge. On the other hand, if a polypus prevents the pus from escaping, the patient is apt to complain of a sensation of pressure and heaviness in the head, dizziness and tinnitus, nausea, vomiting, and even epileptiform symptoms. In extreme cases, paralysis of one side of the body and anæsthesia of the extremities are due to the presence of polypi. The secretion, being forced back into the antrum, may become stagnant, so that cheesy masses are formed, the result being caries and necrosis, or mastoid disease, and possibly an intracranial complication. In the report[1] of a case of chronic purulent otitis media occurring in a man, twenty-four years of age, the symptoms complained of were nausea, vomiting, unsteadiness in gait, and marked deafness. There was facial paralysis of one side, and both auditory canals were filled with granulations. After removal of all granulations and polypi, so that there was free drainage from each ear, there was a marked improvement in the patient's general health, appetite, etc., a decided change for the better in the hearing, and a disappearance of the vertigo and unsteadiness in gait. The facial paralysis eventually disappeared. Occasionally, but very infrequently, a polypus disappears through spontaneous shrivelling. Polypi sometimes come away in the dis-

[1] Bacon: Archives of Otology, March, 1884, vol. xiii. No. 1.

charge spontaneously as a result of the growth becoming turned on its axis and the blood-supply being interfered with. Sometimes granulations coalesce, leading to constriction of the meatus and retention of secretion.

Diagnosis. In order to make a diagnosis of a granulation or polypus, it is first of all necessary to remove the secretion by syringing. Then the canal should be dried by means of absorbent cotton on a cotton-carrier. The surgeon should be careful to distinguish a granulation or polypus from a bulging drumhead from which the dermal layer has come away, or from an exostosis or a malignant growth (epithelioma or sarcoma.) With good illumination and a probe, one can distinguish an exostosis by its being very hard and immovable, and frequently very sensitive to the touch.

An atheroma or sebaceous tumor of the external meatus, although of very infrequent occurrence, might be mistaken for a polypus. In order to decide the point of attachment of a polypus, the probe should be passed around the growth. A large granulation or polypus projecting from the external auditory canal has at times the appearance of a furuncle, but a probe introduced into the canal and passed around the growth will aid one in making a diagnosis.

Where the growth is malignant a diagnosis can only be made with certainty when a section of the tumor has been examined under the microscope. One of the most marked symptoms, however, is pain, which is usually very severe.

A case of aneurism of the carotid artery, appearing as a polypus in the middle ear and followed by profuse hemorrhage when incised, has been reported by Schulte[1].

[1] Monatschr. f. Ohrenheil., March, 1904.

It occurred in a patient, twenty-eight years of age, who had suffered from a chronic otorrhœa for some time. Recently a similar case came to my attention at the Vanderbilt clinic.

Prognosis. The prognosis depends very much as to whether the polypus is attached to the external meatus or tympanic cavity, and also as to whether there are caries of the ossicles and bony walls of the external meatus and middle ear. It is frequently difficult, even with a probe, to determine the point of attachment of a polypus, and in some cases this cannot be ascertained until at least a portion of the growth has been removed. Cases are much more favorable for treatment where the polypus is attached to the external meatus, unless associated with caries of the bony wall.

It is stated by some authors that the appearance of the tumor will aid one in determining the point of its origin, viz., that a red raspberry-like tumor with an irregular surface springs from the middle ear, but a smooth and pale growth has its point of attachment to the external auditory canal.

Treatment. The treatment of granulations is best accomplished by means of (1) curettes or sharp spoons; (2) caustic applications; (3) galvano-cautery; (4) astringent solutions and alcoholic drops.

The speediest and most satisfactory method of removing granulations from the middle ear or external meatus is by means of curettes, sharp spoons, or the so-called ring-knife. The canal should first be cleansed of all secretion and dried, and then a few drops of a 4 per cent. solution of cocaine instilled warm into the ear. After a few moments the canal should again be dried and the

granulations carefully removed. Great caution should always be taken when using cocaine for cases of collapse occur every now and then from instilling a few drops of cocaine into the middle-ear. Some patients are particularly susceptible to its action. In the case of children, it is usully advisable to administer ether or chloroform, as it is next to impossible to keep the head still during the required manipulation. Sharp spoons (see Fig. 85) of different sizes and shapes can all be fitted in the handle for the purpose of removing granulations on the tympanic walls, the external meatus, or the drumhead. Sharp curettes (see Fig. 94), of which there are three sizes, as well as the ring-knife of Politzer, are better adapted to certain cases. The latter instrument can be slipped over a granulation, especially when it is attached to the wall of the external meatus. The knife is then withdrawn quickly, by which means the growth is excised and comes away on the concave surface of the ring. There are two sizes of the ring-knife, the larger one being 3 to $3\frac{1}{2}$ mm. in diameter, while the smaller one is from $1\frac{1}{2}$ to 2 mm.

After removal of the granulations, it is further necessary to cauterize the root or base with nitrate of silver or chromic acid, or to instil into the ear alcohol drops, the latter, however, being contraindicated when there is caries or necrosis. In patients of a nervous temperament, it is frequently impossible to scrape away the

FIG. 94.



Sharp curettes.

growth by means of instruments. In such cases the application of caustics should be tried. For this pur-

pose various remedies have been proposed. For small granulations, a drop of tincture of chloride of iron should be applied to the growth, or the latter should be touched with nitrate of silver fused on the end of a probe, the instrument being first heated red-hot in an alcohol lamp, and then dipped into powdered nitrate of silver. This plan of treatment is especially advisable when small granulations are attached to the middle-ear cavity or the drum membrane. Chromic acid is a much better caustic, as it penetrates more deeply into the tissues than nitrate of silver. A probe should be heated and plunged into powdered chromic acid. It is important, however, that this acid should only be applied to the polypus, and that care be exercised not to touch the drumhead or wall of the external meatus, otherwise severe inflammation is apt to follow. The polypus should not be cauterized a second time until the slough has come away. If the acid occasions much pain, the polypus should be swabbed with cotton soaked in boric acid solution, and afterward dried. It is often advisable to blow into the ear a small amount of boric acid powder after the cauterization.

Galvano-cautery. The galvano-cautery is particularly adapted to the removal of large granulations.

The use of alcoholic drops has been described in Chapter IX. These may be employed with excellent results in the case of small granulations, especially when they are soft and flabby. In some cases great benefit follows the insufflation of equal parts of zinc oxide and boric acid powder, or of xeroform. The same powders are often used after granulations have been excised with instruments.

Treatment of Polypi. Whenever a polypus is large, and particularly if it has a pedicle, the best method of

FIG. 95.

Blake's polypus snare.

removing it is by means of Blake's snare. The instrument (Fig. 95) consists of a metal canula through which a loop of wire is drawn. Instead of having both ends of the wire fastened to a slide-ring, as formerly, there is substituted a cleat, around which one end of the wire should be fastened securely; on the side of the shaft is a similar cleat to secure the other end of the wire. When fastened in this manner, it is evident that by making traction on the slide-ring the wire loop in process of withdrawal through the canal will, instead of merely constricting the growth, have a distinct cutting motion which will greatly facilitate removal of the more resisting polypi. Both ends of the wire may, however, be fastened to the cleat on the slide-ring if the surgeon so desires. The finest malleable iron wire or a very fine silver or brass wire should be used. The instrument, moreover, is so constructed that the handle, slide-ring, and canula can be removed, and the shaft made available as a handle for a paracentesis knife or for curettes or sharp spoons. There is also a small canula which is particularly adapted to excising small polypi in the tympanum. Before attempting to snare

a polypus, the canal should be syringed with a bichloride solution, 1 : 5000, to remove all secretion, and then thoroughly dried. Cocaine solution, 4 per cent., should then be instilled warm into the ear, and after a few moments the surgeon should make out with the probe the point of attachment of the polypus. It requires practice and skill to apply the loop of wire about the growth. The removal of the latter may be followed by considerable hemorrhage, which can be controlled by tamponing the canal with styptic cotton. After the larger portion of the growth has been excised in this way the surgeon will be better able to make out the point of its attachment, the condition of the tympanic cavity, and also as to whether there are other polypi.

When a polypus is found protruding from a small perforation in the drumhead, it is often advisable to enlarge the opening before making an attempt to excise the tumor.

A method formerly in vogue, but now generally discarded as being unscientific, is avulsion by means of forceps. The operation consists in seizing the polypus with a strong pair of forceps, and by twisting and pulling the growth is removed. If attached to the auditory canal, this procedure is not attended with danger, but if arising from the tympanic cavity or drum membrane, great damage may be done by forcible extraction. Curved scissors will be found useful for snipping off a polypus growing from the external meatus. After snaring off the polypus, the root should be cauterized with chromic acid or nitrate of silver, and a strip of iodoform or boric acid gauze should be pushed down to the drumhead. Repeated cauterization should be

performed until the growth is entirely destroyed, and in the meantime any discharge from the ear should be treated as described in Chapter IX.

Beside astringents and powders, which may be used in some cases after removal of the growth, the *alcohol treatment* is often indicated. This method is useful not only for the purpose of shrivelling up the roots of the polypi and granulations on the membrana tympani and in the external auditory canal, but also in those cases in which the polypi are situated in the tympanic cavity, and in cases of exostosis of the external meatus where the growth cannot easily be reached.

In many persons of a nervous temperament, as well as in children, it is impossible to use the snare or other instruments unless an anæsthetic is administered. In many individuals it is impossible to lay down rules for any particular plan of treatment. After excising a polypus with the snare, it may be necessary to complete the operation with a ring-knife or sharp spoon, especially if the bony walls of the external meatus or tympanic cavity are carious. The galvano-cautery is recommended highly by some surgeons where the polypus is large and is of the hard or fibrous variety, and when the snare or other instruments cannot be used. It has the advantage over caustic remedies in that its action is more rapid and thorough, while the pain is but momentary, ceasing immediately after the cauterization.

CARIES AND NECROSIS OF THE TEMPORAL BONE.

In acute purulent otitis media, where pain is a prominent symptom, there is probably more or less inflammation of the periosteal lining of the bony meatus and middle ear. This is generally the case when the disease

is due to scarlatina, measles, diphtheria, tuberculosis, syphilis, and influenza, and unless vigorous antiphlogistic measures are adopted during the early stage (see Chapter VI.) caries and necrosis of the temporal bone are apt to follow, When acute otitis media passes into the chronic form, and proper drainage has not been established for escape of the pus, polypi and strictures of the external meatus may develop, the secretion is apt to become stagnant, and, as a result, decomposition takes place, so that ulceration of the tissues follows and the bone becomes carious, especially if the patient has scrofula, tuberculosis, or some other cachexia. The extent of the caries may be limited, or there may be extensive destruction of the temporal bone. The microorganisms are not so apt to attack the compact bony tissue as the pneumatic and diploëtic varieties, and the disease is more likely to involve the mastoid cells and the posterior superior wall of the external meatus than the other portions of the temporal bone, although it should not be forgotten that the disease may affect the tympanic roof and give rise to meningitis, or to an extradural or temporosphenoidal abscess, or, starting in the mastoid cells, the destructive process may lay bare the lateral sinus and cause thrombosis of the sinus or an abscess in the cerebellum. Cases have occurred in which the disease has extended to the labyrinth and given rise to symptoms suggestive of an abscess in the cerebellum, viz., nausea, vomiting, vertigo, tinnitus, facial paralysis, great pain, and a staggering gait. In such a case, where the pulse was slow and the patient complained of vertigo, vomiting, severe headache, and deafness, on opening the mastoid antrum a carious condition of the semicircular canals was found, which

accounted for all these symptoms. In still another case which came under my care, and in which a diagnosis of "probable cerebellar abscess" was made from the following symptoms, viz., vomiting, a staggering gait with a tendency to fall toward the right side, double choked disk and retinitis, all the alarming symptoms disappeared when the foul-smelling pus and granulations were removed from the external auditory canals and free drainage was established. There was considerable caries of the tympanic cavities as a result of the chronic suppuration.

Still another case of extensive caries of the temporal bone due to chronic otorrhœa occurred in a man, thirty years of age, who was under my care, and who eventually died of a temporosphenoidal abscess. The drumhead was destroyed almost entirely, and there was extensive caries of the inner wall of the middle ear and anterior bony wall of the external meatus (Fig. 96). The case is referred to more fully in the chapter on Intracranial Complications. At the post-mortem examination a large carious opening in the middle cranial fossa was found involving the glenoid fossa.

As a result of caries and necrosis large cavities may be found in the temporal bone. When chronic otorrhœa has existed for some time, and especially when there has been caries of the tympanic walls and ossicula, one finds very frequently a sclerosed condition of the mastoid process, which becomes as hard as ivory. Unhealthy granulations often conceal carious bone, the mucous membrane generally being ulcerated and in an unhealthy condition, while the ossicula are also carious. Caries of the ossicles may occur independently, the disease being limited, as a rule, to one bone, generally the incus.

Frequently the body of the incus and head of the malleus are carious. The malleus and incus frequently are found embedded in a mass of granulations. Caries of the handle of the malleus is not often observed, and it is seldom that the malleus alone is involved. The foot-plate of the stapes generally remains *in situ*

FIG. 96.

Large carious opening in the middle cranial fossa.

although the head and crura may be destroyed. If the foot-plate comes away and the labyrinthine fluid escapes through the foramen ovale, profound deafness is apt to follow. A patient may have fairly good hearing, even when the drumhead and the malleus and incus have been destroyed, provided there is no fixation of the foot-plate of the stapes.

Symptoms. Pain is one of the most prominent symptoms, and it is apt to be very severe and continuous, although in tubercular subjects the disease may occur without pain. The pain is sometimes due to retained secretion, but more frequently it is caused by inflammation of the periosteum or of the bone itself. Other symptoms are tinnitus, vomiting, vertigo, insomnia, elevation of temperature with occasional chills, and increased pulse-rate. The discharge is characteristic, being generally increased in quantity, foul-smelling, and containing blood-corpuscles and occasionally gritty particles of necrosed bone. The external meatus generally is filled with granulations which bleed easily. The remaining rim of the drumhead is swollen and red, while, if the antrum is invaded, there may be a bulging of the posterior upper wall of the external meatus. After syringing the canal and drying it with absorbent cotton, the probe is the most valuable instrument for determining the existence of caries. The bony walls of the meatus are sometimes carious, while abscesses form in the soft tissues about the ear, or over the mastoid, or below the auricle. Occasionally, there is an abscess above the pinna, but rarely in front of it. The pus may force its way into the external auditory canal through one of the fissures of Santorini, or it may break through the soft tissues about the auricle, or it may force a passage to the nasopharynx and give rise to a retropharyngeal abscess. As a result of the formation of abscesses, various fistulæ are found, either in the region of the mastoid, or above, in front of, or below the auricle.

Facial paralysis is another symptom of caries of the temporal bone, and may occur as a result of an inflammatory exudation in the sheath of the nerve. There

may or may not be caries of the Fallopian canal. When caries occurs, it does not necessarily follow that there will be facial paresis. The paralysis may be permanent, depending largely on the extent of the inflammation. As a result of paralysis, there may be conjunctivitis of the same side with ulceration of the cornea.

Prognosis. When there is free drainage for the secretion and the patient is otherwise in good health, the prognosis is much more favorable than when caries occurs in individuals suffering from some exhaustive disease. The destructive process may be limited to the attic and tympanic roof, or it may extend to the dura and brain; while, on the other hand, there may be extensive disease of the temporal bone leading to the formation of a large sequestrum without fatal result.

Caries and necrosis may terminate in recovery with or without much loss of bone substance, or various complications may ensue, viz., meningitis, brain abscess, sinus thrombosis, septicæmia, pyæmia, hemorrhage from the internal carotid artery or internal jugular vein.

As a result of deep-seated caries large sequestra may be formed, including a portion or even the whole of the labyrinth. The outer wall of the mastoid process is occasionally exfoliated (see Figs. 97 and 98).

The auditory canal should first of all be syringed with antiseptic solutions, and as perfect drainage as possible established by means of curettes and sharp spoons. In the case of fibrous strictures of the meatus, very small rubber tubing, through which is passed a quill toothpick, may be used, the quill keeping the tube pervious for escape of the pus, while the rubber exerts elastic pressure on the walls of the meatus. Sometimes caseous masses in the middle ear are forced out

and the pain is relieved by syringing with a warm boric acid solution through the Eustachian tube. When the caries is superficial, excellent results follow in many cases from the injection of Villate's solution.[1]

Dilute sulphuric acid, 1 in 3, is instilled into the ear by some surgeons as well as dilute nitric acid ($\frac{1}{2}$ per cent.) for the purpose of dissolving the carious particles.

FIG. 97.

Sequestrum consisting of the outer wall of the mastoid process.

FIG. 98.



Sequestrum consisting of a portion of the mastoid process.

Such treatment gives relief of only a temporary character. Whenever possible, the carious bone should be scraped and any loose sequestra removed. Scraping of the bone is indicated only when the disease is superficial, and even then care should be exercised not to penetrate the labyrinth if the inner tympanic wall is in a carious condition. The surgeon should likewise be careful not to injure the Fallopian canal and cause facial paralysis, nor to penetrate the middle cranial fossa through the

[1] Villate's solution: sulphate of copper, sulphate of zinc, of each 15 parts; solution of subacetate of lead, 30 parts; vinegar, 200 parts (Bartholow).

tympanic roof. In this connection it is well to bear in mind that the nerve as it passes over the fernestra ovalis is frequently covered inferiorly by mucous membrane and periosteum only. After such an operation on the bone, the auditory canal and middle ear should be syringed with bichloride solution, and powdered iodoform or iodoform and boric acid insufflated. Boric acid gauze and absorbent cotton should then be inserted in the meatus.

When a sequestrum has formed, it should be grasped with suitable forceps, and extracted through the external auditory canal, if possible ; very little force should be employed. In some cases when the sequestrum is large it may be crushed or broken by means of forceps. In other cases, especially when the disease is extensive, it is much better to cut down on the mastoid, making the incision close to the auricle and down to the bone. The auricle should then be displaced forward, and the cartilaginous canal separated from its attachment, in order to allow more space for the operator. If the disease is extensive, it may be necessary to enter the antrum and cut away the posterior superior bony wall, and make one large cavity of this portion of the temporal bone (see Stacke operation).

General Treatment. While local treatment is of especial importance, attention to the patient's general health should never be neglected. As the disease is apt to occur in those who are debilitated, it is highly important to prescribe a nourishing diet, if possible, and plenty of out-of-door life. Cod-liver oil, the hypophosphites, and syrup of the iodide of iron are particularly valuable in tuberculous or strumous children. Iron, quinine, and strychnine are indicated in anæmia, while

iodide of potash should be given if the cause can be traced to syphilis.

Treatment of Facial Paralysis. Large doses of iodide of potash are recommended by some in the early stage when the bone is first inflamed. Counter-irritation over the mastoid by means of iodine ointment is advised by others. The discharge from the ear should be properly treated, and any carious bone removed. After the inflammatory symptoms have subsided, electricity should be tried. If the muscles do not respond to the faradic current, galvanism should be substituted. Frequently no improvement follows until weeks have elapsed, and in some cases even many months.

When the patient has had the paralysis for a year or more, and there seems to be no evidences of improvement, the surgeon should consider the advisability of uniting the facial with one of the neighboring motor nerves. Of twenty-five cases already reported, the results obtained have been very satisfactory in the great majority of cases according to F. Furet[1]. In seven of these cases the facial was joined with the hypoglossal nerve, while in the others the facial and spinal accessory were united. Ballance and others recommend the hypoglossal-facial operation. In cases of anastomosis between the spinal accessory and facial, the movements of the arm and elevation of the shoulder of the side operated on, at once cause a contraction on the corresponding side of the face. The results seem to show disappearance of degenerative action, diminution of facial asymmetry and among most of the cases reëstablishment of the loss of mobility. According to Furet

[1] Traitement chirurgicale de quelques paralysies faciales d'origine otique. Soc. française d'Oto. Larngol., Mai, 1905.

one notices at the sixth month after operation an improvement in the reestablishment of symmetry of the face in repose, while later, in favorable cases, there is a return of certain voluntary movements such as closure of the eye-lid and deviation of the labial commissure, with improvement in phonation and mastication. The cure can be further hastened by gentle massage and electricity.

CHAPTER XI.

DISEASES OF THE MASTOID PROCESS.

Before entering into a consideration of the diseases of the mastoid process and their complications, it is necessary to consider the anatomy of the temporal bone, as well as its position in relation to important structures, such as the large bloodvessels, the dura mater, and the brain.

The temporal bone consists of a squamous, a petrous, and a mastoid portion. Ossification of the temporal is developed by ten centres, viz., six for the petrous and mastoid portions, one for the squamous and zygoma, one for the tympanic plate, and two for the styloid process. Inflammatory changes of a destructive character are more apt to take place in the petrous and mastoid portions than in the squamous and tympanic. During the first years of life the points of union between these different portions are very vascular, and at times are not firmly united, so that inflammatory products, especially tubercle bacilli, can easily break through these vascular walls and give rise to a collection of pus in the soft tissues behind the auricle, a condition of affairs which is of frequent occurrence in scrofulous children.

In the child the external auditory canal is as large in proportion as in the adult, except that in the former the canal consists almost wholly of cartilage, while in the adult it is bony for two-thirds of its length.

In the infant at birth there is an incomplete ring with two tubercles, one in front and one behind; these tubercles approximate one another in the lower wall of the meatus at the glenoid fossa, and form a foramen which does not become closed until the fifth or sixth year of age. This aperture, however, sometimes remains open

FIG. 99.



Section of bone showing the inner or mucous surface of the drumhead; also the articulation of the head of the malleus with the body of the incus contained within the attic.

permanently, or it may be covered by thin fibrous tissue, and, as it is in close proximity to the parotid gland, it is easy to understand how inflammation may extend from the external meatus to the parotid gland, or *vice versâ*. During an attack of mumps the drumhead frequently is congested, and in some cases the labyrinth is affected as well.

The tympanum communicates with the Eustachian tube in front and the mastoid antrum and cells behind. The drumhead forms the greater part of its outer wall. Beneath the floor of the tympanum, and separated from it by bone, is the jugular fossa posteriorly and the carotid artery anteriorly. In the upper portion of the

Fig. 100.

a. Antrum. *b, b, b, b.* Mastoid cells. *c.* External osseous meatus.

tympanum is the attic (Fig. 99), containing the articulation of the head of the malleus with the incus; the attic opens posteriorly into the mastoid antrum (Fig. 100) and the latter into the mastoid cells. Some writers describe a third cavity, viz., the aditus ad antrum, which is the passage between the attic and the antrum. The tegmen tympani, or roof of the middle ear, consists of a very thin plate of bone, upon which rest the dura

mater and brain. In some cases there are defects in the tegmen tympani, which are covered over with fibrous tissue, so that in acute otitis media the inflammation may extend directly to the dura and give rise to meningitis or brain abscess. In cases of fracture of the temporal bone the tympanic roof may be involved, symptoms indicative of this complication being escape from the meatus of blood and cerebrospinal fluid.

Macewen[1] enumerates the following communications with the tympanic cavity, and states that pathological processes may extend through these apertures, viz.:

1. The external auditory meatus.

2. The Eustachian tube and canal for the tensor tympani muscle.

3. The mastoid antrum and cells.

4. The petro-squamosal fissure running the whole length of the tegmen tympani, open in early years, and marked by numerous channels in after-life.

5. The masto-squamosal fissure, open in childhood, occasionally remaining unossified in after life, and marked by numerous foramina when ossified.

6. The fenestræ ovalis and rotunda communicate with the vestibule and cochlea, respectively, and the former communicates with the internal meatus ; a venous canal from the cochlea communicates with the pyramidal part of the jugular fossa.

7. The Fallopian canal, having communications with the tympanic cavity : in infancy, by a defect in its anterior wall above the fenestra ovalis, and permanently by the aperture for the stapedius muscle and the small nerve twig for the same; by the channel for the chorda tympani, which again pierces the wall of the tympanum

[1] Pyogenic Infective Diseases of the Brain and Spinal Cord, pages 5–7.

at the canal of Huguier in the fissure of Glaser; by an arterial twig, the stylo-mastoid branch of the posterior auricular, which passes into the tympanum.

The Fallopian aqueduct opens above the cribriform plate into the internal meatus, and through the hiatus Fallopii and a smaller hiatus in front of this (transmitting the greater and lesser superficial petrosal nerves, respectively) into the middle cranial fossa, so inflammation may extend along the perineural sheaths and set up lepto-meningitis of both posterior and middle fossæ.

8. The foramen between the carotid canal and the jugular fossa for Jacobson's nerve to the sympathetic plexus and inner wall of the tympanum.

9. The fossa subarcuata—in childhood a wide foramen transmitting a vein to the cells under the superior semicircular canal, and at a still earlier period to the mastoid cells near the outer aspect of the infantile mastoid.

10. The lateral and sigmoid sinus—by means of numerous minute veins through the posterior and lower part of the tegmen tympani and through the mastoid antrum and cells.

11. The superior petrosal sinus—by means of numerous minute venous channels.

12. The inferior petrosal sinus—by means of veins from the labyrinth.

13. The temporo-maxillary vein—by many minute veins from the anterior part of the tympanic cavity.

14. Free communication through the bone with the dura matral system of veins.

15. The carotid canal—pierced by several foramina for nutrient vessels and branches of the sympathetic nerves to the tympanum, also by small veins.

16. The jugular fossa—the roof of which is pierced by a great number of veins communicating with the floor of the tympanum and the labyrinth, and also by the foramen for the auricular branch of the vagus (Arnold's) communicating with the Fallopian canal in its course.

17. The tympanic branch of the internal maxillary artery which enters at the fissure of Glaser, to supply the fore part of the tympanum.

There are many veins, and they have an important bearing, since inflammatory products are occasionally carried by them to the large sinuses and to the brain itself. Of considerable importance is the mastoid vein, which proceeds from the sigmoid sinus directly through the skull externally, and, joining a large vein from the occipital region, passes to the deep cervical vein, which ultimately unites with the vertebral. The course of this vein is not constant, however. The mastoid vein may become involved when a thrombus is found in the sigmoid sinus. The veins from the tympanum empty partly into the middle meningeal, partly into the sigmoid sinus, and partly into the superior petrosal sinus. The mastoid veins, as a rule, enter the sigmoid sinus. Too much stress cannot be laid upon the importance of remembering the position and relation of these veins one to another, as will be seen in a later chapter on intracranial complications.

When the chorda tympani nerve is injured, as may happen from a blow or fall, or from instrumental manipulation, there will occasionally be observed a loss of the sense of taste in the anterior two-thirds of the tongue on the side affected. The facial nerve may be paralyzed in its course through the middle ear from injury

due to an operation, or from disease of the middle ear, or the facial paralysis may be of central origin. As a result of such paralysis conjunctivitis occurs, and occasionally ulcerations on the cornea, the patient being unable to close the eye. The sense of smell is impaired. The patient has difficulty in mastication, the food collecting between the cheeks and gums on the affected side. Taste is affected through the chorda tympani. The patient is unable to whistle. In paralysis of the facial due to a central lesion, the patient generally can close the affected eye, and the muscles of the face are much less affected, while the chorda tympani is not involved. In children, the wall of the canal for the facial is extremely thin, and occasionally there is a small opening in this wall, so that facial paralysis is of more frequent occurrence in children than in adults. It is important to remember the course of the Fallopian canal, for the surgeon is likely to injure the nerve during operations on the mastoid as well as when excising carious ossicles. The facial canal, in its course through the upper parts of the tympanum, lies just above the foramen ovale and directly behind the attic and antrum, so that the surgeon should avoid the inner and anterior wall of the antrum. After removing carious ossicles, great care should be exercised in scraping the attic. The patient's face should constantly be watched, to see that there is no twitching of the facial muscles. Even when this precaution has been taken, facial paralysis has occurred, as the muscles will probably not respond to stimulation when the patient is completely under the influence of an anæsthetic. In early life the mastoid process is but slightly developed, and consists principally of the antrum, which is large and very near the surface

of the skull. The ossicula in the infant are of almost the same size as in the adult.

The mastoid cells in adult life vary very much as to size, and from anatomical investigations it is known that no two processes are exactly alike. The cells in one case may consist of large air-spaces, which communicate with one another, as in Plate I., the so-called pneumatic mastoid process; or they may be arranged as

Fig. 101.

Pneumatic mastoid process.

in Fig. 101; or the mastoid process (Figs. 102 and 103) may be composed of small-celled diploë and osseous tissue filled with fatty substances; or the mastoid process may be sclerosed. The pneumatic mastoid process is generally larger than either the diploëtic or sclerosed varieties. In some cases the mastoid is composed partly of pneumatic cells and partly of diploëtic tissue. According to my experience, in cases in which the external wall of the mastoid process is small, and especially when the

FIG. 102.

Inner surface of a diploëtic mastoid process, with the groove for the sigmoid sinus lying just above the cells.

FIG. 103.

Section through diploëtic mastoid.

1. Lateral sinus. 2. Mastoid cells. 3. Anterior canal wall cut away to show membrana tympani.

lower portion is poorly developed, the lateral sinus is more likely to run closer to the antrum than in those cases in which the mastoid is filled with large air-cells. In such, moreover, great care should be taken not to

PLATE VI.

FIG. 1.

Pneumatic Mastoid, showing Unusual Development of Styloid Process.

1. Squamous portion. 2. Mastoid cells. 3. Styloid process. 4. Stylomastoid foram

FIG. 2.

Arrested Development of Mastoid Process.

1. Mastoid process.
2. Digastric groove.
3. Styloid process.
4. Glenoid fossa.
5. Zygomatic process.
6. Auditory process.
7. Suprameatal spine.
8. Opening for mastoid vein.

FIG. 1.

rior bony canal wall cut away to show membrana tympani.

Membrana tympani. 2. Umbo. 3. Long ss of incus. 4. Suprameatal spine. 5. oid process. 6. Styloid process. 7. Squa- portion. 8. Middle root of zygoma.

FIG. 2.

Mixed variety showing diploëtic and pneumatic mastoid cells.

1. Diploëtic tissue in tip. 2. Pneumatic cells. 3. Antrum. 4. Attic. 5. Incus. 6. Membrana tympani. 7. Umbo. 8. Digastric groove. 9. Aquæductus Fallopii. 10. Tegmen tympani. 11. Squama.

FIG. 3.



xed variety showing diploëtic and sclerosed mastoid.

. Sclerosed tip—diploëtic tissue above. 2. uæductus Fallopii. 3. Horizontal semicircu- canal. 4. Superior semicircular canal. 5. action of hiatus with the aquæductus Fallopii. Promontory.

FIG. 4.



Mixed variety showing pneumatic and diploëtic mastoid cells in an unusually thick bone.

1. Pneumatic cells—diploëtic above. 2. Portion of parietal bone showing continuation of cells. 3. Antrum. 4. Promontory. 5. Portion of carotid canal. 6. Jugular fossa. 7. Groove between condyle of occipital bone and the foramen lacerum posterium. 8. Digastric and occipital grooves. 9. Aquæductus Fallopii.

injure the sinus when opening the mastoid antrum. The bony wall between the mastoid cells and the sigmoid sinus is very thin, and in some cases there are defects in the bone which are filled in with fibrous tissue.

In the child there is an opening found in the petrous bone on its posterior surface, the fossa subarcuata, lying between the external auditory meatus and the groove for the superior petrosal sinus. It terminates in a series of cells just beneath the superior semicircular canal. Soon after birth this canal gradually fills up with more or less bony tissue, so that eventually it is but a small sinus. It is usually of small size in the adult, and as a rule it is closed. It is of importance because, when present, inflammation may extend from the mastoid cells to the brain, as a vein enveloped in dura mater passes through it. The mastoid antrum, in the child just after birth, is covered by a much thinner layer of bone than in the adult, so that pus makes its way much more readily to the external surface of the skull. The mastoid process does not assume the typical shape until the child is about three years of age. The mastoid antrum is the point which the surgeon should always expose when operating for mastoid disease, especially when some intracranial complication has occurred, for it is there that most pathogenic processes have their origin.

MASTOIDITIS EXTERNA.

Primary Acute Inflammation of the Mastoid Periosteum.

Primary acute inflammation of the periosteum covering the mastoid process, is rare, although such cases do

occur and have been reported, the congestion first appearing in the periosteum covering the mastoid. The causation of *mastoid periostitis* is frequently obscure, but it may be due to cold, injury, or extension of the inflammation from a furuncular condition of the external meatus. The hearing in such cases is generally normal, as no changes are apt to take place in the drumhead. There may be a slight congestion of the posterior bony wall of the external meatus. The auricle generally stands out prominently, while the tissues over the mastoid are painful to the touch, red, and œdematous. The pain is apt to be severe in some cases, and there may or may not be considerable fever. When one or more furuncles have formed in the external meatus, the inflammation may extend to the soft tissues about the ear, and especially in the mastoid region, so that the parts are very tender and œdematous, with the auricle standing out prominently. The beginner is very likely to mistake this condition for disease of the mastoid process, but the presence of a furuncle in the canal will give the surgeon a clue to the causation. It is possible, however, in severe cases of acute otitis media, for the periosteum over the mastoid process to be affected, especially in strumous subjects. Unless a free incision is made down to the bone, there is a liability that the latter will become involved and the inflammatory process extend to the mastoid cells. The disease may run its course in a few days and subside rapidly, or suppuration may take place. The abscess may open into the external meatus through one of the incisuræ Santorini. As a result of periostitis, the superficial layers of bone underneath may become necrosed unless an

PLATE VIII.

FIG. 1.

Upper portion of temporal bone removed, showing tympanum, with ossicles in position, mastoid, antrum and cells, semicircular canals, cochlea and internal auditory canal.

FIG. 2.

Section through temporal bone, showing pneumatic mastoid cells and an unusually long stylomastoid process

incision is made through the periosteum down to the bone. In the early stage of this affection the artificial leech should be applied, followed by the Leiter cold coil. If these measures do not afford relief, a free incision should be made through the soft tissues over the mastoid down to the bone. If the bone beneath has become carious, this should be removed with chisels and gouges.

MASTOIDITIS INTERNA.

Primary Acute Inflammation of the Mastoid Cells.

Cases have been reported from time to time, and undoubtedly occur, in which the mastoid cells are primarily involved. The disease may be due to cold, trauma, syphilis, tuberculosis, or the influenza, or may develop spontaneously. The microörganisms of influenza, or the diplococcus pneumoniæ may be found in this disease, as well as other microbes.

Henrici[1] of Rostock has reported eight cases of tuberculosis in the mastoid in children and recently has added the histories of three more cases under seven and two in the eighth year. His combined statistics show that primary tuberculosis of the mastoid process occurs in somewhat less than one sixth of the cases of mastoiditis of children. According to this author, this form has especial interest on account of its character as primary bone tuberculosis, and not as a disease transmitted from bone tuberculosis of the tympanum. The usual mas-

[1] Further experiences in primary bone tuberculosis of the mastoid process in children. German Med. Society. June 9-10, 1905. Archives of Otology. Vol. xxxiv., No. 6.

toid operation generally leads to recovery, although the period of recovery is somewhat prolonged. The prognosis is more unfavorable the younger the child. This writer believes that the primary form is much more frequent than tuberculous mastoiditis extending from the tympanum.

The symptoms are pain in the region of the mastoid, which, though slight at first, usually becomes very severe and throbbing. This is followed by tenderness on pressure over the mastoid bone, and if the pus makes its way through the bony wall fluctuation will be detected in the soft parts and the auricle will stand out prominently. In this affection the disease may reach its height in a week or ten days, and the inflammation subside without suppuration, or an abscess may form, which may perforate the outer cortex after several weeks of suffering. There is always danger that the inflammation may extend to the sigmoid sinus if pus forms and is not evacuated, or that the dura may become involved through the thin tympanic roof.

As in periostitis of the mastoid, so in this disease, the hearing usually is unaffected. It is difficult to make a differential diagnosis between primary mastoiditis and mastoid periostitis, except that in the former the pain is deeper seated. If a subperiosteal abscess has already developed, it will be impossible to tell whether the mastoid cells are implicated until a free incision has been made down to the bone and the latter has been examined for a sinus. It is always well in such cases to open the cells to see if pus has formed.

Prognosis. The prognosis is usually favorable in simple, uncomplicated cases. The surgeon should be guarded in expressing an opinion, however, when the disease occurs in a syphilitic or cachectic individual.

Treatment. The treatment of this disease is similar to that for acute mastoid inflammation occurring secondarily to acute purulent otitis media.

Acute Inflammation of the Mastoid Cells occurring in connection with Acute Suppurative Otitis Media.

Inflammation of the mastoid cells occurring secondarily to acute otitis media is the form most frequently observed. The disease starts generally in the nasopharynx, and extends by the Eustachian tube to the tympanum, and thence through the antrum to the mastoid cells. The inflammation of the cells may follow an acute catarrhal or acute suppurative otitis media, or occur as the result of chronic purulent otitis media, frequently of long standing. Mastoid disease developing in connection with chronic purulent inflammation of the middle ear will be described separately.

Acute inflammation of the mastoid cells may follow prolonged exposure to wet or cold. Other causes are the exanthematous diseases, diphtheria, tonsillitis, bronchitis, typhus and typhoid fevers, tuberculosis, syphilis, influenza, the use of the nasal douche, and "sniffing up" salt and water for the relief of nasal catarrh. It is probable that in almost all cases of acute otitis media the mucous lining of the mastoid cells is simultaneously involved. The diploëtic mastoid cells are not affected as often as the pneumatic variety. In some pneumatic processes the opening from the antrum into the cells is large. If, however, the opening is small, or it becomes occluded through swelling of the mucous membrane, the escape of pus is prevented, and the discharge, being forced back, forms an abscess.

Since the influenza made its appearance in this country there have been, according to my experience, many more cases of mastoid disease than ever before. It is only a few years ago that but twelve, or at most twenty, cases of this disease were recorded in the annual report of the New York Eye and Ear Infirmary. In the report for 1896 there were 135 mastoid operations, while for 1897 the number had increased to 161, and in 1905 to 552. Formerly, it was the exception for both mastoid processes to be involved simultaneously in a case of acute otitis media. During the past few years such a condition of affairs has been very frequent. A patient will have a sudden severe earache, which is followed in a few hours by a bulging of the drumhead and symptoms of mastoid inflammation. Of twenty cases that came under my observation recently in private practice, all had acute otitis media. In seventeen there was acute inflammation of the mastoid cells, and in seven it was necessary to open the mastoid process. In ten recovery took place without operation. Of these, the drumhead was incised in six cases. In one of them, viz., that of an infant, several months old, in whom alarming symptoms developed suddenly, suggesting a meningeal complication, puncture of the drumhead gave speedy and marked relief.

Symptoms. The most prominent symptom is pain in the mastoid process, which radiates usually in different directions—upward in the temporal region and downward and forward toward the teeth. Great pain is felt on percussion of the bone or when firm pressure is made. The pain on pressure usually is limited to the region just over the mastoid antrum or to the middle and lower thirds, especially at the tip. Sensitive-

ness to pressure does not necessarily signify that the bone is diseased at that particular point, so that it is always important to make a careful examination of the entire mastoid process during an operation in this region. In some cases pus forms in the mastoid cavity with very little, if any, pain. This frequently happens when the disease occurs in connection with tuberculosis, and occasionally when due to influenza. As mastoid disease is observed frequently during the course of scarlatina, diphtheria, and other infectious diseases, it is most imperative for the surgeon to make a *free incision in the drumhead* at an early date. An abscess in the mastoid cells, occurring secondarily to acute purulent otitis media, is more likely to be located in the middle and lower thirds than in the upper third. Other symptoms of mastoid inflammation are: increase of surface temperature of the affected side; bulging of the posterior and upper portions of the drumhead, with drooping of the adjacent soft parts of the external meatus; after perforation has occurred, the appearance of a nipple-like process, at the extremity of which is found a minute opening, and profuse discharge from the ear. The temperature may vary from 99.5° to 104° or 105° F. As a rule, in simple, uncomplicated cases it is lower in adults than in children. A patient with extensive caries of the mastoid cells and with a carious opening into the cranial cavity may have a temperature but slightly above normal, so that a comparatively low temperature does not necessarily signify that the case is not one of considerable gravity. Not long ago a patient was admitted to my service at the Infirmary with an acute purulent otitis media. There was absolutely no pain on pressure over any portion of the mastoid process. As

the discharge continued to be very profuse in spite of all treatment and the patient had a very septic appearance with a badly coated tongue, a mastoid operation was performed. The mastoid cells were filled with pus and the bone was very much diseased. That the patient did not complain of pain when pressure was made on the bone was due to the fact, probably, that the outer cortex was very thick. The profuse discharge in this instance was the principal symptom. Young infants frequently have a temperature of 105°, while in adults one is much more apt to find the temperature below 100° F., even when extensive mastoid disease occurs. Great surprise is often expressed by the general practitioner that there can be any serious disease with such a low temperature. I have never seen a case of acute inflammation of the mastoid cells without slight elevation of temperature, except in one instance, and I look upon this symptom as having a very important bearing on the diagnosis. If a patient is under observation who has had an acute purulent otitis media for a week or ten days, and an examination reveals bulging of the upper wall of the auditory canal and drumhead,and there is pain on pressure over the mastoid antrum or tip ; and, further, if there is some rise in temperature, frequently very slight, and especially if there is a profuse discharge, these symptoms taken together are very characteristic of mastoid inflammation. The condition of both mastoid processes should always be compared, because in some persons pressure made over the normal mastoid causes pain, especially in hysterical subjects. Bulging of the upper and posterior portion of the drumhead and drooping of the adjacent cutaneous lining of the external meatus are, to my mind,

absolute symptoms of mastoid involvement, and I believe that in such cases it is always necessary to perforate the mastoid cells. Formerly it was taught that œdema and redness of the tissues behind the ear, with prominence of the auricle, were evidences of mastoid disease. Surgeons have frequently waited for these symptoms, as well as for the appearance of fluctuation. An operation should never be deferred till such a late stage, as valuable time may be lost by so doing. The fact should not be overlooked that redness and œdema over the mastoid, and tenderness on pressure, occur in cases of furunculosis of the external meatus. An examination of the auditory canal will exclude this.

A case was lately brought to my attention by the family physician on account of certain symptoms suggestive of mastoid disease. The patient, a young lady of seventeen, had an attack of chickenpox, and at the same time complained of severe pain in the ear and over the mastoid process. She had a temperature of 101½° F. The external auditory canal was swollen, so that it was impossible to obtain a good view of the drumhead. The tissues over the mastoid were very tender to the touch and œdematous. The hearing-distance, however, was normal, and a careful inspection of the inflamed tissues showed the cause to be due to the presence of papules beneath the skin.

Acute inflammation of the mastoid cells may terminate in recovery without the formation of an abscess, but more frequently suppuration occurs and carious destruction of the bone results, so that after weeks of suffering a large sequestrum may be found. In influenza cases rapid destruction of bone is the rule, so that

it is imperative in such to open the cells very early. The pus may force its way through the tympanic roof, and give rise to meningitis or brain-abscess, or may cause destruction of the thin plate of bone covering the sigmoid sinus and produce thrombosis, or perforation may take place at the mastoid tip (Bezold), thus allowing the septic matter to burrow beneath the deep fascia, and give rise to septicæmia and pyæmia, or to a postpharyngeal abscess.

In the pneumatic mastoid process, where the cells are large and the outer cortex thin, the pus is more likely to force a passage externally and form a subperiosteal abscess than in the diploëtic and sclerosed processes. In the latter, according to my experience, the tendency is for the pus to escape through a carious opening in the tympanic roof, and to give rise to an extradural abscess or an abscess in the brain-substance itself.

Facial paralysis is observed in some instances as a result of disease of the mastoid process itself as well as of injury inflicted by the surgeon during an operation. In a child, one and one-half years of age, in whom facial paralysis made its appearance before operation, the mastoid cells were found extensively diseased, so that it was necessary to remove bone in all directions. Although the prognosis as to recovery from the paralysis in such cases usually is considered unfavorable, steady improvement followed the application of electricity, although the facial muscles did not respond for some time to the current. In this case an abscess had formed over the mastoid process. This was incised and a sinus found, with the probe, leading down to the antrum and tympanum. Such abscesses behind the auricle are observed frequently in

children who are strumous or tuberculous, and the disease in some cases is more marked in the antrum than in the middle ear itself. In young children, as previously mentioned, the different portions of the temporal bone are not united except by fibrous tissue, so that purulent secretion easily may force its way through, especially as the outer wall of the antrum is particularly thin. The pus escapes through a carious opening in the external wall of the antrum or else through the mastosquamosal suture, and as the periosteum is not so firmly adherent in a direction upward over the squamous portion as downward toward the tip, the subperiosteal abscess generally is situated above and behind the auricle, the latter standing out prominently. As such an abscess is seen more frequently in cachectic subjects than in those who are healthy, there is likely to be extensive caries, not only of the mastoid antrum, but also of the squamous portion of the temporal bone. In an infant operated upon but a short time since, there was a large subperiosteal abscess with extensive caries extending to the middle cranial fossa. The pus had undoubtedly burst through the antrum while the drumhead was not even perforated and but slightly congested, showing that the disease was confined to the antrum. The surgeon should always in such cases, after making a free incision in the soft parts, make a thorough examination with the probe for carious bone, and in most cases it will be found.

Some portions of the temporal bone are attacked more frequently by caries and necrosis than others. The different portions are involved usually in the following order: 1. The mastoid process. 2. The antrum and tympanic roof. 3. The bony wall of the tympanum.

covering the sigmoid sinus. 4. The posterior bony wall of the external meatus. 5. The floor of the tympanum, including the posterior wall of the canal for the carotid artery. Lastly, the petrous portion, including the bony labyrinth.

The various micro-organisms found in the mastoid antrum reach this cavity, as a rule, by means of the Eustachian tube. It is possible, however, for a purulent discharge in the ear to escape into the nasopharynx and reaching the lungs cause a pneumonia; or, if swallowed, the patient may have a foul-smelling diarrhœa. The streptococcus pyogenes and the staphylococcus aureus are frequently found, not only in the purulent secretion from the middle ear, but also in cases of brain-abscess, suppurative leptomeningitis, and thrombosis of the sigmoid sinus, especially when the thrombus has become disintegrated. Occasionally one discovers the presence of the staphylococcus pyogenes albus and citreus, the bacillus pyogenes fœtidus, and the diplococcus pneumoniæ of Fränkel. According to my experience, when streptococci are present the inflammation is apt to be of a severe type and to resist the ordinary treatment, while complications are likely to follow. In a case recently seen in consultation the acute otitis media followed an attack of tonsillitis. Both mastoid processes were involved, and the sigmoid sinus was thrombosed on the right side. Streptococci were found in the discharge. The case terminated fatally.

It is advisable to examine microscopically the pus from the tympanum, and to have cultures made, since much information can be obtained in this way as to the probable severity of the disease and as to the prognosis. A short time ago a patient was under my care who had

an acute purulent otitis media in consequence of an acute nasopharyngeal catarrh. As there was considerable prostration, which was out of all proportion to the symptoms, the temperature being but slightly above 99° F., a culture was made and the Klebs-Löffler bacillus was found. On the following day a fibrinous cast was discharged from the tympanum through the perforation.

During the past six years the total number of ear secretions examined at the N. Y. Eye and Ear Infirmary amounted to 2037. Of these 3 showed coli communis, 5 tubercle bacilli, 17 spirillum of Vincent, 52 Friedlander's bacillus, 62 staphylococcus, 260 pneumococcus, 445 streptococcus, 594 mixed, and 599 miscellaneous, negative, etc.

In this report Dixon states that the pus taken for examination from the external auditory canal usually presents a mixed infection—that is, various micro-organisms; but generally the predominant germ is found in the mastoid cells, the sigmoid sinus, or brain itself, when these regions are secondarily involved. For instance, the streptococcus is found frequently in pure culture in cases of infective sinus thrombosis, while the micro-organisms from the auditory canal may be of the mixed variety. If the streptococcus and the pneumococcus are present in the external auditory canal, one generally finds these organisms associated in the more deeply seated suppurations, the other micro-organisms which may have been present in the discharge from the canal having disappeared.

Prognosis. The prognosis is generally favorable in a simple, uncomplicated case if the disease is of recent origin. When occurring in connection with influenza, typhus fever, scarlatina, measles, tuberculosis, and

diphtheria, on account of the rapid destruction of bone which frequently takes place, the prognosis becomes more grave, particularly if the disease has existed for a long time. In ten days' time, particularly if the patient has diphtheria, scarlet fever, or influenza, the entire mastoid process may be filled with pus, and present a carious opening into the posterior or middle fossa. Frequently in cases of mastoiditis where streptococci are present the surgeon will find the sigmoid sinus embedded in pus and granulations even when the disease has existed for a short time and the symptoms have not been indicative of gravity. It is not always possible to tell whether the brain or the sigmoid sinus is implicated before the operation is performed, so that until the condition of the mastoid cells has been investigated the surgeon should be guarded as to the prognosis.

Treatment. If the surgeon is fortunate enough to see a case of acute inflammation of the mastoid cells in the first stage before pus has formed, the patient should be put immediately to bed and kept very quiet, and should be given a very light diet, consisting principally of milk. Calomel should be administered in $\frac{1}{10}$ to $\frac{1}{4}$ grain tablets, one every hour for six doses, or until a laxative effect has been produced. Tincture of aconite, in one drop doses, is also an excellent remedy in the early stage. The artificial leech should be applied at once in two or more places on the mastoid bone, and especially just over the antrum and over the tip. It is advisable at the same time to make a *free incision in the drumhead* (see Fig. 60), along its posterior border from a point nearly opposite the stapes down to the lowest point of the membrane, especially if there is fluid within the tympanic cavity. In some cases the surgeon should

prolong the incision up to the posterior fold of the drumhead and through the tissues into the attic, and then carry the incision along the posterior and upper wall of the external auditory canal. For this purpose a large knife should be used (see Fig. 59), and at the location just mentioned, in order to drain the attic and antrum if possible. The day has gone by for using a small paracentesis needle. After the patient has been put to bed the Leiter coil (see Fig. 104) should be adjusted and kept in place for *from thirty-six to forty-eight hours, but never longer than this.* The cold generally relieves the pain at once, but if left on for too long a time it may mask other important symptoms. Cold should under no circumstances be applied *except in the first or hyperæmic stage, and should never be reapplied* after having been removed. The coil consists of a malleable plate which can be so moulded as to fit any mastoid process. Ice-water, by the siphon principle, passes through this coil by means of rubber tubing attached to the coil. The upper tube connects with a basin containing water and a large piece of ice, while the lower basin catches the water which has passed through the coil and the other tube. The best manner of arranging these basins is to place one on a level with the patient's head and the other about six inches lower. In this position the water is prevented from flowing too quickly through the tubes.

I cannot too strongly protest against the frequent use of blisters for just these conditions, a practice which is not only pernicious, but which also tends to aggravate the disease. Many a case of mastoid disease that has come under my care has been due to the application of a blister behind the ear for the relief of an acute otitis

media. After the soft parts behind the ear have been inflamed by such measures it is almost impossible to make a diagnosis of mastoid disease. If, instead of recommending blisters or poultices, the physician would apply a leech, and if the pain were not relieved by it and the drumhead were bulging, would make a free incision, there would be fewer cases of mastoid disease under the care of the otologist.

Fig. 104.



Leiter coil.

By means of the treatment here outlined, it is possible to relieve the pain in many cases, and to abort the disease if the case is a simple, uncomplicated one. This plan of treatment is, however, less successful in cases of mastoiditis occurring in connection with scarlatina,

influenza, syphilis, and tuberculosis. When the mastoid cells are involved during the course of one of the exanthematous diseases, influenza, or diphtheria, the surgeon should first of all make a free incision in the drumhead (see Chapter VI.), and if the inflammatory symptoms do not disappear quite rapidly under the above-mentioned treatment, the mastoid cells should be opened early, otherwise there is likely to be a rapid destruction of bone. After the drumhead has been incised, it is well to douche the ear at first with a normal saline solution or a 4 per cent. boric acid solution every hour or so, according to the amount of the discharge and the degree of pain present. In cases due to scarlatina, diphtheria, influenza, or streptococcus and pneumococcus infection, in addition to the boric acid douching, the ear should be syringed at least twice a day with a solution of bichloride of mercury (1 : 5000). The patient should be kept in bed for some time, even after the acute symptoms have passed. Chloroform or nitrous oxide gas should always be administered when incising the membrana tympani. In the case of young children chloroform is to be preferred to nitrous oxide gas. Lately I have used ethyl chloride by means of an inhaler especially adapted for the purpose. Chloroform should be used most cautiously, however, and only in such quantity that the patient will not feel pain when the incision is made. When an anæsthetic cannot be given, the pain from incising the drumhead can be lessened very much by painting the membrane with a five per cent. solution of cocaine in equal parts of aniline oil and 95 per cent. alcohol as recommended by Gray.

In a case under my care in which acute otitis media occurred during an attack of measles, the tempera-

ture suddenly rose to 104° F. There were inflammation and bulging of both drumheads, together with tenderness on pressure over the antrum and lower third of the mastoid process on each side. Incision of each drumhead and application of the Leiter coil, with frequent syringing of the auditory canals, was the treatment employed. After incising the membranæ tympani the temperature fell to 100° F. The patient recovered without operation.

In a child of eleven, who had an attack of grippe and tonsillitis, followed by earache on the left side and inflammation of the mastoid process, the temperature was 103° F. There was a sticky discharge from the canal two days later, which gave some relief to the patient. This was but temporary, however, for on the following day she vomited and had a return of the earache. She was seen by the author in consultation on the next day, and at that time there was tenderness on pressure over the middle and lower thirds of the mastoid. There was some sticky sanious discharge from the left auditory canal, with bulging of Shrapnell's membrane. There seemed to be a small perforation in the membrana flaccida, and also one in the anterior inferior quadrant of the drumhead. In spite of the treatment, which consisted in the application of leeches, incision of the drumhead, and the use of the ice-bag (see Fig. 106), the mastoid symptoms became more marked, and twenty-four hours later the temperature rose to 106.2° F. It was necessary to open immediately the mastoid cells and remove granulations and softened bone. The temperature varied from 104° to 105.8° F. At this time evidences of pneumonia on the left side were first discovered, in spite of the fact that a careful examina-

tion of the chest had been made each day. Two days later the temperature began to decline gradually, and perfect recovery followed (see Fig. 105).

Other cases, similar to the preceding one, have come under the care of the writer, in which the first symptom

Fig. 105.



Temperature chart of a case of acute otitis media followed by acute mastoiditis, complicated with pneumonia.

was a severe earache, usually following an attack of influenza. An important diagnostic point is the presence usually of the pneumococcus in the discharge from the ear, and the fact that the temperature is very much elevated and does not show the sudden variations which are characteristic of infective sinus thrombosis (see Fig. 126). It is extremely difficult to detect the pneumonia at first, on account of the consolidation being probably deep-seated. When the physical signs are well marked, the discharge from the ear is apt to become scanty, but later, when convalescence has been established, the otorrhœa may again be profuse. It is especially important in all cases in which pneumonia is suspected to avoid giving an anæsthetic, and particularly ether. When the temperature remains elevated after

the mastoid cells have been opened, one can be fairly confident that some complication exists, possibly a pneumonia, or thrombosis of the sigmoid sinus or other intracranial disease. There are cases of mastoiditis, however, in which the temperature remains high after operation, and in which the cause is due undoubtedly to septic absorption. This is especially the case where pus

Fig. 106.



Author's ice-bag for application over the mastoid process.

has been found in the mastoid cells, and where the temperature has been considerably elevated some days before the operation. I have several times been inclined to perform some further operation in certain cases in which the temperature did not come down soon after opening the mastoid cells. It is always advisable to abstain from administering antipyretics in cases of mastoid disease or other complications, because it is most important

for the surgeon to make a diagnosis at a very early stage, and to operate immediately in case of a brain complication, if he wishes to meet with success. If it is desirable to reduce the temperature, it is better, especially in the case of children, to resort to sponging the body with alcohol and water. If the rise in temperature is due to the presence of pus, it is important for the surgeon to find this out as soon as possible.

During dentition children are very apt to have attacks of acute suppurative otitis media, and occasionally the mastoid process becomes secondarily involved. In a case[1] of acute suppurative otitis media, which was characterized by high temperature and symptoms of brain implication, and in which it was necessary to remove carious bone from the mastoid antrum, the patient recovered, but the causation of the disease was most obscure until four molar teeth made their appearance.

Inflammation of the mastoid cells may follow an unskilful attempt at removing adenoid vegetations. Such a case came under my observation, and it is surprising that similar instances are not more frequent, when one considers how roughly the operation is occasionally performed, and how often the orifices of the Eustachian tubes must be lacerated. The case reported[2] was that of a woman, twenty-four years of age, who complained of a sore throat. She was told by her physician that she had adenoids, and on the following day he excised the growths. She soon afterward had an attack of earache, followed by mastoid inflammation. This case illustrates two most important points: first,

[1] Bacon: Annals of Ophthalmology and Otology, Jan., 1894, vol. iii. No. 1.

[2] Bacon: Transactions of the American Otological Society, 1892.

that all operations should be done with as much gentleness and skill as possible; and, secondly, that an operation for the removal of adenoids should never be undertaken until all acute symptoms have subsided.

Of late years otologists have reported cases of *Osteomyelitis* as if the disease were entirely distinct from an ordinary suppurative inflammation of the mastoid cells. According to Dr. E. H. Nichols of the Harvard Medical school, "infective osteomyelitis is an acute suppurative inflammation of the bone and may be produced by any one of a variety of pathogenic microörganisms or by a mixed infection". Osteomyelitis should simply be considered a severe type of bone inflammation and due generally to the presence of the staphylococcus or streptococcus, particularly the latter. Knapp[1] says "that osteomyelitis exists only in the diploic variety of the mastoid process," and he reports a case where the disease extended to the occipital bone with disseminate abscesses in the bone and under the deep muscles of the neck. Two operations were performed, resulting in total resection of the mastoid and removal of a carious piece of the occipital bone with exposure of the cerebellar dura. The case did not recover for months. The disease is found more frequently among the young and it is possible that the changes in the bone of young children may have something to do with the development of the disease. It is well-known that, in infants and young children, myelin cells are present in the soft parts of the bone in considerable quantity, only to become very much less numerous as the child grows until finally in the adult there are but few myelin cells except in the long bones, the sternum and the vertebræ.

[1] Archives of Otology, Vol. xxxii, No. 6.

The patient seems to suffer from profound intoxication with marked general depression and usually has a high temperature. The area of tenderness over the mastoid bone is extensive and œdema is apt to appear at an early stage. When an operation is performed the suppurative inflammation will be found to extend over a greater area than is usual, involving in some instances the occipital and parietal bones. Even after all diseased bone has been removed at the time, the surgeon may, at a subsequent operation, find that the disease of the bone has extended in various directions. The author has operated on a number of these cases and and when occurring as a complication of measles, osteomyelitis is quite likely to be followed by meningitis.

It is important in this connection to refer to a complication that is an extremely infrequent one, viz., the so-called *white erysipelas* affecting the scalp. In the case[1] of a young woman who had an acute suppurative otitis media followed by mastoid disease, the cells were opened and the entire bony contents removed with granulations. The temperature, instead of falling, gradually became higher, reaching 104.4° F. Later the patient had a chill lasting twenty minutes, and vomited. It was noticed that the scalp near the wound was œdematous, tender to pressure, but not pinkish in color. The tenderness and œdema gradually extended over the scalp, while the discharge from the ear and wound was much diminished in quantity. The temperature in a few days became normal, and the patient made an excellent recovery. Another point of interest in the case was the fact that before the operation the urine was found to contain a large amount of sugar, which disappeared,

[1] Revue de Laryngologie, d'Otologie, et de Rhinologie, 1895, No. 8.

however, before the patient left the hospital. Several urinary examinations were made afterward, but no sugar was found. The case before the erysipelas developed presented symptoms characteristic of septic absorption. The so-called white erysipelas has been known to affect the eyelids, the scrotum, the prepuce, the vulva, and the inferior extremities of persons enfeebled by disease. The glycosuria seemed to be of nervous origin and due to reflex action.

Indications for Operation. It is undoubtedly true that some cases of simple, uncomplicated inflammation of the mastoid cells recover spontaneously if left to nature, while, on the other hand, it should not be forgotten that lives have been sacrificed because the operation of opening the cells had been deferred too long. As the operation is a comparatively safe procedure if performed under aseptic conditions, I think that where the surgeon is at all in doubt he should always err on the safe side and open the cells at once. I would not, however, be understood to make light of the operation of perforating the mastoid, because it should never be undertaken by one who does not thoroughly understand the anatomy of the temporal bone, and who has not had considerable experience in treating aural diseases.

If, under the use of antiphlogistic measures and the Leiter coil, the pain rapidly subsides, while the discharge becomes less and the symptoms generally improve in a few days, the temperature showing no evidence of septic absorption, the chances are that an operation will not be necessary. On the other hand, if the pain persists and the temperature remains high after the appearance of the discharge, and especially if the latter is profuse, and inspection of the drumhead shows

it to be bulging in its upper and posterior quadrant, with drooping of the adjacent soft tissues of the external meatus, and, moreover, if there is considerable deafness with pain on pressure over the mastoid process, these symptoms taken together should cause the surgeon to decide upon an immediate operation.

It is not safe to delay operation in cases of mastoid disease caused by the influenza, scarlatina, measles, diphtheria, or where there are streptococci and pneumococci (either Sternberg's or Friedländer's) present in large numbers in the discharge from the ear. Experience has shown that complete destruction of the bony cells and perforation of the inner wall may take place in such cases in an incredibly short time, without the usual well-marked symptoms of mastoid disease. In a patient, fifty years of age, who was under observation a short time ago, and who had been treated for two weeks for an acute purulent otitis media, with tenderness over the mastoid cells, it was deemed best to open the cells, owing to the fact that there was a profuse discharge, which did not yield to treatment, as well as a temperature slightly above normal, which persisted, and tenderness on pressure just over the apex of the mastoid. In this case the cells were entirely destroyed, and there was a large carious opening over the lateral sinus. The patient was up and about, and did not complain of pain in the mastoid except when firm pressure was made over the tip. This latter symptom is a most valuable one, and should never be lost sight of in arriving at a diagnosis. Undoubtedly, in the near future, otologists will be more ready than ever to make cultures from the pus in purulent otitis media, for considerable information can be obtained in this way as to the prognosis and treat-

ment. Examination of the blood is also of great value, especially when the surgeon is in doubt about operating. An increased leucocytosis with the polynuclear elements considerably above the normal will often decide one to operate at once. An increase in the polynuclear elements is especially significant.

In a report of forty cases[1] of acute purulent otitis media, complicated by acute inflammation of the mastoid cells occurring in my private practice, it was necessary to perform a radical mastoid operation in ten instances. In two of these operative cases a subsequent operation for sinus thrombosis was performed, one terminating fatally. This was the only case of death among these forty cases. *In thirty cases, or 75 per cent., the mastoid cells were not opened.* The ages of the forty patients varied from six to sixty-five years. There were twenty children under fifteen years, and of this number sixteen were males and four females, while of the remaining twenty adults, ten were males and ten females. The cases which recovered without operation were treated in the manner already described, viz., by means of the artificial leech, the Leiter coil, and by making a free incision in the drumhead. This conservative line of treatment should never be carried out for any length of time except by one who has had considerable experience in treating aural diseases, and who can recognize very early the more serious symptoms so characteristic of mastoid disease. The aural surgeon should be in close touch with the patient, and if he does not see him each day, he should be in a position to know his exact condition, so that he may operate at a moment's notice in case of necessity. The surgeon should make

[1] Bacon: "Non-operative Cases of Acute Inflammation of the Mastoid Cells," Archives of Otology, 1901, vol. xxx., No. 1.

every effort to cure his patient without opening the mastoid cells. If, however, the latter operation becomes necessary, he should not hesitate for a moment, not only to remove all diseased tissues from the mastoid antrum down to the tip, but he should be fully competent and prepared to operate at all times for any intracranial complication.

Operation. Before undertaking an operation either for mastoid disease or one of the complications arising from it, the surgeon should first of all find out about

FIG. 107.



Sterilizer.

the patient's general condition. A thorough urinary analysis should be made and the patient's heart and lungs should be examined. On the day of the operation the patient should have nothing but liquid food, and even this should be forbidden for at least three hours before taking the anæsthetic. If possible, a laxative should be administered the evening before, followed by some mineral water in the morning.

Ether is to be preferred, on account of its being

a safer anæsthetic than chloroform. In short operations, however, chloroform may be used if carefully administered. The ether should be given by a thoroughly competent assistant, who should attend to this duty alone, and thus relieve the surgeon of this anxiety. On account of the disagreeable sensations caused by the inhalation of ether, it is well to give nitrous oxide gas at first, and then change to ether. An excellent apparatus (Bennett's inhaler) has been devised for this purpose, and may be obtained of the instrument-makers.

All instruments should he sterilized thoroughly, and for this purpose the sterilizer shown in Fig.107 is a very convenient one. The blades of the scalpels should be dipped in a boiling 5 per cent. carbolic solution or in absolute alcohol. The hands of the operator and of his assistants should be scrubbed thoroughly with soap and water by means of a nail-brush, and afterward washed with a bichloride solution, 1 : 3000. Of late years I have used sterile gloves for all operations.

The instruments used in a mastoid operation are the following: one narrow and one broad scalpel, artery forceps, periosteal elevator, dressing-forceps, gouges and chisels, a mallet with lead filling, rongeur forceps, sharp spoons (seven sizes), retractors, grooved directors, probes, catgut ligatures, needles, and silk sutures.

The hair of the patient should be shaved within a radius of about two and a half inches (Fig. 108) of the auricle. The parts thus shaved should be scrubbed with soap and water, and afterward washed with a solution of bichloride of mercury, 1 : 3000. The external meatus should be syringed with a warm bichloride solution, then dried and filled with a narrow strip of iodoform gauze. The head should be wrapped with a sterile towel

PLATE IX.

Fig. 1.



1, 2. Scalpels.
3. Halsted's artery forceps.
4. Small French artery forceps.
5. Straight scissors.
6. Curved scissors.
7, 8. Dressing forceps.
9, 10. Probes.
11, 12. Grooved directors.

FIG. 2.



1. Ballance's hand gouges.
2, 5. Large gouge and chisel.
3. Ballance's long gouges.
4, 6. Schwartze's gouges and chisels.
7. Mallet with lead filling.

PLATE X.

FIG. 1.



1. McBurney's needle holder.
2. Aneurism needle for ligating internal jugular vein.

3 to 9. Sharp spoons, different sizes.
10, 11. Retractors.

FIG. 2.

1. Periosteum elevator.
2. Stacke's protector.
3. Jansen's rongeur forceps.
4, 6. Rongeur forceps.
5. Author's rongeur forceps.

and fastened with a safety-pin. Similar towels should be laid over the patient's face, neck, and shoulders and only the region to be operated on should be exposed. After these precautions have been taken, the auricle being gently pulled forward by an assistant, an incision

FIG. 108.



Lines of incision made in operations for mastoiditis, sinus thrombosis, and brain abscess. From point 1 to point 2, incision for mastoid operation. Incision extended from 2 to 4 and from 2 to 5 in operation for brain abscess. Incision from original line for mastoid operation to point 3 in operation for sinus thrombosis.

should be made by the surgeon, just behind and close to the auricle (from point 1 to point 2, Fig. 108), from the extreme tip of the mastoid to a point just above the upper border of the auricle. The soft parts and perios-

teum should be divided, and the latter pushed aside by means of the periosteal elevator. Hemorrhage should be controlled by means of the artery clamps and by pressure. Dry sterile gauze cut in strips makes the best and most satisfactory sponges. The entire surface of the bone should then be examined carefully for a carious spot or sinus, and, if found, an opening at that

FIG. 109.



Integument and periosteum removed from mastoid process. (Gerber.) Temporal muscle. *T.*, *L.t.* Linea temporalis. *Sp.* Spina supra meatum. *Fo* Temporal fossa. *G.* External auditory canal. *Ste.* Tendon of Sterno-cleido-mastoid muscle.

point should be made into the cells by means of a gouge and mallet. A large gouge should be used at first, and all the diseased bone carefully removed with sharp spoons. The surgeon should endeavor to reach the antrum, knowing that the latter is just above and behind the external meatus. Great care should be taken in scraping the antrum, for fear of injuring the facial nerve.

Macewen uses burrs (Fig. 110), which can be rotated by means of an electric motor (Fig. 18). He prefers them to chisels and gouges. They are excellent instruments when carefully handled; but when considerable bone is to be removed and the operation is to be performed as quickly as possible, the surgeon can work more rapidly with chisels, gouges, and rongeur forceps.

Fig. 110.

Macewen's burrs.

If there is no sinus evident, nor a soft spot of bone, the surgeon should first chisel an opening into the antrum and establish communication with the tympanum. A guide for this operation is the spina suprameatum, a point about one-quarter of an inch below the linea temporalis. The bone should be removed at a point about one-eighth inch behind the spine and the instrument directed downward, forward, and inward, in order to reach the antrum. The entire external wall of the antrum should be removed. Small pieces of bone should be cut away at a time, and the parts carefully inspected by means of good illumination. A probe or director should be used constantly to see if an opening has been

made into the antrum. The sharp spoon will also be found very useful in this step of the operation. As a rule, the bone should not be entered at a depth greater than five-eighths or three-quarters of an inch, for fear of injuring the facial nerve or external semicircular canal. There is also danger of wounding the lateral sinus and dura mater in either the posterior or middle cranial fossa, unless one

FIG. 111.



Opening of the antrum in simple empyema (Gerber). *A.* Mastoid antrum. *Sp.* Suprameatal spine. *G.* External auditory canal.

uses the chisels carefully and in the direction already mentioned. The opening in the bone should be cone-shaped from without inward. After establishing a communication with the antrum and attic, the surgeon should then turn his attention to the other portions of the mastoid, remembering that in some acute cases the pus is limited mostly to the middle portion, while in others it is found in the tip with granulations and softened bone. In

some instances the sigmoid sinus will be found exposed and bathed in pus and covered with granulations. Cases have been reported of sinus thrombosis which probably developed from inadvertent exposure of the sigmoid sinus during a mastoid operation—Kœrner explains this pyæmia from the irritation of the sinus by the introduction of gauze so that a parietal thrombus is formed, and he warns against exposing a normal sinus unneces-

FIG. 112.



Removal of the mastoid cells completed (Gerber). *A*. Antrum. *Su*. Sigmoid groove. *C*. Few remaining mastoid cells.

sarily. In scraping away the granulations it is an easy matter to wound the sinus. A thorough examination should be made of the tip, to see if it is perforated, for in some cases pus may burrow down the neck beneath the deep fascia. If an abscess has already formed in the tissues just about and below the tip, it should be opened freely and the pus evacuated. The tympanic roof should

be examined to see that a carious opening does not exist. All softened bone, granulations, etc., should be removed thoroughly by means of sharp spoons, and the entire mastoid cavity thoroughly irrigated with a 1 : 5000 bichloride solution. If possible, the surgeon should try to force some of the solution through the antrum and the external meatus. A small perforation in the drumhead should be enlarged. The cavity in the bone should then be packed with narrow strips of moistened iodoform gauze, and afterward several sutures may be taken in the upper and lower angles of the wound, to diminish the size of the opening. Larger strips of iodoform gauze should then be placed about the ear, so as to cover the wound and the auricle, and outside of this sterile gauze, while the whole should be covered with a layer of absorbent cotton and bandaged. It is frequently advisable to change the outer dressings on the following day, if they have become saturated and stiffened, so as to be uncomfortable to the patient. Unless the patient has more or less fever, with pain, the packing in the mastoid should be left in place for at least two or three days. If the iodoform gauze is left *in situ* too long, it may become adherent to the bone cavity, and cause bleeding when removed. To prevent this and to make the first dressing as painless as possible, some surgeons use a perforated piece of sterile rubber tissue which is inserted into the mastoid cavity and then filled with narrow strips of gauze so as to make equable pressure on all parts of the bony cavity. After the first dressing, boric acid or plain sterile gauze should be substituted for the iodoform gauze, and should be applied loosely, so as to allow the cavity to fill up as rapidly as possible with granulation tissue.

In some cases it is better to continue the use of iodoform gauze especially if the granulations show a tendency to be pale and unhealthy. In the latter case the balsam of Peru should be poured into the wound before repacking it with gauze. Several days should elapse between the dressings, if possible, for frequent syringing with antiseptic solutions has a tendency to prevent the granulations from growing. After the operation the patient usually experiences great relief, and the temperature, if elevated, becomes lower. The discharge from the ear is usually diminished in quantity or ceases altogether. The time before cicatrization takes place varies from two to six weeks. It is longer frequently in cachectic or scrofulous individuals. If all of the carious bone has not been removed, a second operation becomes necessary. When granulations are exuberant they should be cauterized with a solid stick of nitrate of silver. Silver drainage-tubes, iron nails, rubber tubing, etc., should never be used for purposes of drainage, for gauze is to be preferred.

In some cases, if the antrum has not been opened and there is no communication with the middle ear, and the disease is limited to the middle or lower third of the mastoid process, and especially if the inflammation occurs in a pneumatic mastoid process and is near the external surface, the surgeon may allow a blood-clot to form, and he may close with sutures the entire wound, provided, however, that every trace of disease has been removed with sharp spoons. If the patient complains of pain, or there is a rise of temperature, the wound should immediately be reopened and treated by packing it with gauze in the *usual way*.

In the treatment of an abscess behind the auricle, frequently observed in the case of scrofulous children,

it is not sufficient to make an incision and evacuate the pus, for it should be remembered that in all probability the purulent matter has forced its way either through the outer wall of the antrum or through the petroso-squamosal suture, and that in such a condition of affairs considerable carious bone undoubtedly exists. The child should be etherized and a long incision made behind the auricle close to the bone, and the condition of the antrum and cells carefully investigated.

Diet. After the operation it is desirable to keep the patient quiet in bed for a few days at least, until the temperature becomes normal and the patient feels inclined to sit up. While there is any elevation of temperature the diet should be a light one, consisting for the most part of milk and beef-tea. After a week's time, if the temperature is normal and the case progresses favorably, I frequently allow a hospital patient to go home and be treated as an out-patient, impressing upon him the importance of remaining in-doors during inclement weather.

Attention to the General Health. Particular attention should be paid to the general health, especially when the patient is cachectic or scrofulous. In the latter case cod-liver oil, iron, strychnine, or the hypophosphites are indicated, while syphilitic or tubercular subjects should have appropriate constitutional treatment. This is a subject of the greatest importance, for in many instances local treatment alone avails but little if attention to the general health is neglected.

MASTOID DISEASE OCCURRING IN CONNECTION WITH CHRONIC PURULENT OTITIS MEDIA.

Very often, during the course of chronic otorrhœa the ossicles become carious, while the antrum is apt to be filled with purulent material, and in some instances the entire antrum and mastoid process may contain cholesteatomatous masses, or there may be within the bony cavity cheesy matter, softened bone, granulations, or tubercular material. In other cases the mucous lining may be very much thickened, so that the cells are obliterated, while in still other instances the granulation-tissue contained in the cells may turn into bone, so that the mastoid process becomes wholly or in part changed into dense osseous tissue as hard as ivory (sclerosed mastoid). There may be a circumscribed caries of the mastoid process, or it may be very extensive, leading to various complications. For instance, a patient recently came to my clinic who had had a discharge from the ear for ten months. He complained of pain and tenderness over the mastoid process, especially the tip. A leech was applied, which gave some relief. He returned in a few days, when there were evidences of slight facial paralysis on the same side as the ear inflammation. He was admitted immediately to the hospital, and when placed under ether great destruction of bone was found, with the mastoid cells filled with pus and broken-down tissue. The lateral sinus was exposed, and there was a large carious opening in the tympanic roof. This was enlarged, and considerable pus escaped from the middle fossa. The boy recovered. This case goes to prove the statement previously made that a patient may have,

with very few symptoms, extensive caries and necrosis of the mastoid process.

The most frequent cause of mastoid disease in connection with chronic purulent otitis media is interference with the drainage from the antrum or middle ear, so that pus collects in the mastoid process and forms caseous material. Constriction of the external meatus, the formation of polypi, granulations, a small perforation in the drumhead, suppuration limited to the attic with perforation in Shrapnell's membrane, and cholesteatomata, may cause mastoid disease. Various microorganisms in the discharge from the middle ear, including tubercle bacilli, may give rise to mastoid inflammation. Other causes are cold, injury, and the various diseases mentioned in the etiology of acute mastoid disease.

Symptoms. The symptoms are similar to those mentioned in connection with acute mastoiditis, especially when occurring as a result of exposure to cold or retained secretion; but when eburnation of the bone takes place, this process may be gradual and extend over a period of years, so that there are few if any symptoms.

The external auditory canal frequently is found very much constricted and filled with retained secretion, the latter being foul-smelling and sanious, especially if granulations or polypi are present and there is carious bone. A cholesteatoma may exist for years in the antrum without occasioning serious trouble until some inflammatory symptoms arise.

Although during the course of a chronic purulent otitis media there may at stated intervals be attacks of pain in the mastoid process which do not lead to abscess,

it is well to bear in mind the fact that serious complications are liable to develop at any moment, such as sinus thrombosis, meningitis, brain abscess, perforation of the mastoid tip, septicæmia, pyæmia, etc., unless operative measures are undertaken at an early period. In tubercular cases there may be great destruction of bone without any well-marked symptoms of pain. In such cases formerly it was considered inadvisable to operate. Experience, however, goes to show that removal of carious bone and tubercular material is followed often by marked improvement in the patient's general health, and such an operation should always be undertaken, provided the patient's general condition is satisfactory. As hygienic surroundings as well as good food are so necessary in the care of these unfortunate patients, it is to be hoped that in the near future wards will be so constructed on the roofs of all hospitals that the fresh-air treatment can be successfully carried out in all cases of tuberculosis. Much has already been accomplished on these lines.

If left to nature, perforation of the external bony wall of the mastoid takes place in some cases, an abscess forming in the soft tissues, which ruptures and leaves one or more sinuses leading to carious bone. The external cortex is the most frequent seat of perforation, although the pus may force a passage through in other directions.

In Fig. 97 a sequestrum, three-quarters of an inch by three-quarters of an inch in size, is shown, which was removed by operation. It consists of the greater part of the external wall of the mastoid of a boy, aged two years, who had had a recent attack of measles. The right ear had been discharging for a month, and

an abscess had ruptured behind the ear, which gave great relief to the pain.

Diagnosis. The diagnosis can be made if there is pain on pressure over some portion of the mastoid, with constriction of the external meatus, retention of pus, and granulations present in the canal. In cases of cholesteatoma it is more difficult to decide, because frequently the pain is slight; while, on the other hand, in osteosclerosis the pain is often quite severe.

Prognosis. A guarded prognosis should always be given when the patient is debilitated, tuberculous, or cachectic, or there are extensive caries and necrosis of the temporal bone or some intracranial complication.

Treatment. Whenever symptoms of mastoid disease are well marked in a case of chronic purulent otitis media, the first object of the surgeon should be to establish good drainage from the middle ear. All granulations or polypi should be removed carefully from the canal, so that one can see the drumhead or what remains of it. At times, especially when the pain in the mastoid is due to inflammation, and not to the retention of pus or cholesteatomatous masses, antiphlogistic measures will be sufficient; but, as a rule, it is much more advisable to open the mastoid antrum, for with only slight symptoms a patient may develop some intracranial complication, and it is not prudent to wait long before operating.

If only a rim of the drumhead remains (see Fig. 113), while the ossicles are carious and the disease seems to be limited mostly to the attic and antrum, it is undoubtedly true that in some instances a cure can be established by excision of the drumhead and ossicles, and by curetting and washing out the attic and tympanic cavity with antiseptic solutions through a middle-

ear canula. The same holds true of cases in which the perforation is in Shrapnell's membrane and pus is pent up in the attic.

When the attic contains cholesteatomatous masses it is always advisable to open the antrum, for the surgeon can never be certain of the extent of the disease until

FIG. 113.

Drumhead almost entirely destroyed. Rim of membrane left in the upper portion. Ossicles bound down by adhesions. Long handle of malleus attached to inner wall of middle ear.

he has explored the mastoid cells. On perforating the outer cortex of the mastoid he may find complete destruction of the entire mastoid cells.

The surgeon should always operate whenever there are sinuses in the mastoid region leading to carious bone, or when the posterior bony wall of the meatus is diseased.

In certain cases in which the mastoid becomes eburnated and the cells have disappeared, great relief will follow the chiselling away of portions of the mastoid process, and especially is it important to make an opening through to the antrum.

In chronic mastoid inflammation it is always advisable to enter the antrum, in order to enable the surgeon to wash it out, together with the middle ear, through the opening behind the auricle. In a pneumatic or diploëtic mastoid process one should investigate the entire bone, from the antrum to the tip, being careful not to injure the sigmoid sinus, which occasionally runs very close to the antrum in a sclerosed or diploëtic mastoid process. If the operation is done carefully under aseptic conditions, and the patient's general health is fairly satisfactory, the operation is generally successful. In cases of diabetes, when the disease is in an advanced stage, the operation is one of more or less gravity, but should be undertaken when the patient's life is in danger from extension of the inflammation to the brain.

The same instruments are used as in the operation for acute mastoid disease, and the same solutions. The incision is identical with the one recommended previously, and the entire mastoid process should be inspected.

In cases of *osteosclerosis* it is generally advisable to chisel away sufficient bone to reach the antrum and establish drainage with the middle ear. Such a procedure usually relieves the pain, which is so distressing to the patient.

One should, of course, be careful not to injure the sigmoid sinus, which may be exposed and covered with granulations, or the facial nerve, or the dura itself.

Instead of the disease being limited to the mastoid process, the surgeon may find that the posterior bony wall of the meatus or the walls of the tympanum are carious. With the object of removing the outer wall of the attic and the posterior superior bony wall of the meatus, surgeons have devised different methods of operation, the idea being to expose entirely the tympanic cavity and remove all carious bone. This procedure, viz., removal of the posterior superior bony wall and outer wall of the attic, is indicated in cases in which there is extensive caries involving the attic and antrum, the middle ear, or bony meatus, and especially where there are cholesteatomatous masses or numerous granulations or polypi which impede the drainage from the tympanic cavity.

KÜSTER'S METHOD. First the auricle is detached from its insertion, and displaced forward. An incision is made close to the pinna and behind it, as in the mastoid operation, from the tip of the mastoid to the commencement of the helix. The tissues are divided down to the bone, and the periosteum pushed aside so that the posterior superior portion of the meatus is exposed. The cartilaginous meatus is then, by means of a periosteum elevator, carefully pushed away from its posterior superior attachment, being pulled downward and forward. The further steps in the operation depend very much on the extent of the carious bone. If the disease is limited to the outer wall of the meatus, it may be necessary to remove this alone by means of small chisels or the sharp spoon, or the surgeon may think best to open the antrum in the usual way, and afterward to cut away the posterior superior wall of the external meatus to the tympanum. One should be careful to avoid the facial nerve, which

lies, however, in the deeper portion of the posterior wall. A small gouge or cutting forceps is best adapted for removing the bone. The necrotic malleus and incus should be excised, and, if necessary, almost the entire mastoid process, antrum, middle ear, and external meatus can be transformed into one large cavity. This operation is indicated particularly in cases of cholesteatoma. The wound is then irrigated with bichloride solution and packed with iodoform gauze. A drainage-tube is introduced into the external auditory canal to keep the cartilaginous meatus in position, and the auricle is replaced. When the wound behind the ear is a large one, it is well to pack it with gauze and allow it to granulate from the bottom. If, however, the posterior superior wall alone has been removed, the iodoform gauze may be inserted through the external meatus, and the wound behind the auricle closed by sutures.

Stacke Operation. Stacke divides the operation into three steps:

1. Incision through the soft tissues.

2. Removal of the cortex of the suprameatal triangle and the posterior bony canal, and the extraction of carious ossicles, with curetting of the tympanum, aditus, and antrum.

3. Construction of the flap from the membranous canal and its application to the walls of the bony canal, antrum, etc.

The various steps in the operation are performed as follows: An incision is made through the soft tissues, following closely the attachment of the auricle, and extending from one centimetre (three-eighths of an inch) below the mastoid tip upward to a point a little above and forward of the superior border of the external audi-

tory meatus; the lower two-thirds of the incision is made directly down to the bone through the periosteum; the upper third extends through the skin only down to the fascia of the temporal muscle. This muscle and fascia are then drawn aside and an incision made through the periosteum underlying them and bordering upon the superior margin of the bony meatus. The periosteum of the anterior flap is then elevated and drawn forward

FIG. 114.

Middle ear, attic, antrum and adjoining mastoid cells converted into one large cavity. (Schwartze-Stacke operation.)

1. Mastoid tip. 2. Styloid process. 3. Digastric groove. 4. Opening for mastoid vein. 5. Portion of posterior canal wall containing facial nerve. 6. Glenoid fossa. 7. Zygoma. 8. Squamous portion of temporal bone.

until the suprameatal triangle and posterior wall of the bony canal are plainly exposed, this procedure being simplified at times by cutting away a few fibres of the sterno-cleido-mastoid muscle at the tip, so as to afford greater room. This stage of the operation is completed by separating the membranous canal from its bony attachments and by pulling it outward. A retractor is then introduced and the soft parts are pulled forward, so that the bony structures, including the tympanic ring, are seen readily.

The second step consists in cutting the tensor tympani muscle and extracting the malleus, if it is visible, through the canal; then the remnant of the drum membrane, if any is left, is excised and Stacke's protector is introduced into the attic. The bone forming the external wall of the attic is cut away and the incus removed. The protector is then inserted into the aditus

FIG. 115.



Flap incision in Stacke operation. (Gerber). *a* and *b* flaps made from the cartilaginous canal. *op.* bone cavity the result of operation.

ad antrum and the bony wall external to it chiselled away so that the protector can be pushed into the antrum.

The overlying cortex and remainder of the posterior bony wall are removed with a chisel. The protector acts as a guard to insure the safety of the facial nerve. By such means the antrum, attic, aditus and tympanum are converted into one large continuous space, which must be curetted thoroughly for the removal of granu-

lations and all soft carious spots in the contiguous walls.

The third step in the operation consists in making a horizontal incision in the vertical meridian of the membranous canal above and extending from the tympanic ring, the entire length of the canal, outward to the concha. A second incision is now made, beginning

FIG. 116.



Stacke's flap in position. (Gerber) *a* and *b* flaps.

in the concha, where the first one terminates, and running at right angles to it downward and backward in the canal, thus making a nearly four-cornered flap (Fig. 115), which is crowded backward and downward by tampons which are inserted through the external auditory canal, so as to cover as much as possible of the denuded bone. Antrum, aditus and tympanum are now packed with gauze, without irrigation, and the skin

of the upper half of the flap closed with sutures, the lower half being covered with antiseptic gauze and bandaged. In cases in which the post-auricular opening is not closed immediately Stacke stitches the upper edge of the flap to the inferior angle of the wound. (Fig. 116.)

The object of the subsequent treatment is to secure, if possible, a smooth epidermal covering of the large cavity made in the bone. Granulations form upon the raw surface of the bone, and when they become exuberant should be cauterized with nitrate of silver or the galvano-cautery. The epidermis should advance from the sides and cover the entire surface.

Stacke keeps the patient in bed for seven or eight days. The outer dressing may be removed earlier, but the deeper dressings are not changed until the eighth day. The dressings after this are changed more frequently, and finally are dispensed with when it is apparent to the surgeon that the epidermis has covered more or less of the excavation.

Schwartze-Stacke Operation. Schwartze's operation differs in several important points from the method advocated by Stacke. In the first place, Schwartze does not believe in separating the entire membranous canal from its bony attachments and in pulling it out, provided the calibre of the canal is not abnormally narrow. He leaves the anterior wall attached to the bone, in order to prevent subsequent necrosis and the formation of a sequestrum in the external meatus. He introduces a slender retractor and pulls the soft parts forward, thus gaining a sufficient amount of space. Then he opens the mastoid antrum, starting from the outer surface of the mastoid process in the usual way, and proceeds from this point to complete the operation, instead

of cutting the bone from within outward, as Stacke suggests, viz., by beginning the operation in the attic, and then passing a protector upward and backward into the antrum as a guide for entering the latter cavity.

Schwartze does not believe in the use of the Stacke protector, for the reason that it is likely to injure the tegmen tympani. He recommends, on the other hand, a broad chisel, and considers that with short and careful blows of the hammer there is little danger of injuring the stapes, which usually lies embedded and protected in the swollen mucous membrane. He advises the use of a sublimate solution (1 : 1000), not only during the operation, but later in the changing of the dressings whenever there is any noticeable fetor. He also recommends the syringing of a sterilized salt solution through the Eustachian catheter into the middle ear in cases in which there is caries about the tympanic opening of the Eustachian tube. If there is no fetor, however, he does not irrigate the wound, but simply dusts over it iodoform powder. He does not close the mastoid wound at all immediately after the operation, believing that the packing with gauze cannot be accomplished successfully through the external auditory canal alone. Schwartze does not change the first dressings till the sixth day, unless the patient has fever, or a discharge from the wound, or else complains of pain. If, on the other hand, the surgeon has decided to close the mastoid wound immediately after the operation, he should not allow the dressings to remain unchanged longer than three days, for fear that the retained secretion may force open the sutures. As iodoform gauze tends to irritate the tissues and to favor the growth of exuberant granulations, it should be used only in the first dressings.

Otherwise he prefers simple sterilized gauze, and when the epidermis begins to form he advocates the use of aristol powder, to protect the newly formed skin. He discontinues the tampons and allows the mastoid wound to close through cicatricial contraction, when the epidermis has already extended over the greater portion of the antrum, aditus ad antrum, and middle ear.

In order to prevent the occurrence of stenosis from the growing together of exuberant granulations, the latter should be destroyed with a knife especially constructed for the purpose or by means of the galvanocautery. Great care should be taken to keep the aditus and antrum open in order to secure good drainage.

Schwartze believes that the employment of the lead-nail is preferable to packing the wound with gauze in many cases, because the latter method is extremely painful; and since the treatment must be kept up for weeks, this is an important consideration, especially in the case of children. On account of the severe pain experienced in changing the dressings, it is often advisable to moisten the tissues with a 20 per cent. solution of cocaine.

Various modifications of these different operations have been suggested and practised by different surgeons. The best method of operative procedure, however, is first to cut down on the mastoid process and lay bare the antrum, and then decide, according to the indications, just what operation is best adapted to each individual case, the object being to remove all carious bone without doing injury to the facial nerve.

In the writer's experience it is always necessary to curette thoroughly the tympanic orifice of the Eustachian tube in order to close completely this opening and

thus prevent reinfection of the middle ear from the naso-pharynx.

In all cases in which the disease is limited to the attic and antrum and there is no evidence of cholesteatoma nor exposure of the dura, and especially when the lower two-thirds of the mastoid process is sclerosed, the Stacke operation is indicated. The posterior opening should in such instances be closed and the wound treated through the external auditory canal. If on the other hand, the case is one of cholesteatoma, or the dura is exposed in either the middle or posterior cranial fossa, the Schwartze-Stacke operation should be performed and the wound behind the auricle should be kept open and the cavity allowed to granulate up from the bottom. The flap can then be stitched as in Fig. 116. Thiersch grafts can be used to advantage in some cases but they should never be placed over the dura as fatal cases have been reported from such practice. At a meeting of the Otological Section of the New York Academy of Medicine,[1] Gruening reported a fatal case where a most thorough plastic operation had been performed in connection with the radical operation for chronic purulent otitis media. The canal and also the tympanic wall were lined with skin but behind that skin pus was found and the patient died of meningitis. He called attention to the danger of making large flaps to cover over denuded bone.

[1] Archives of Otology, Vol. xxxiv., No. 6, Dec. 1905.

CHAPTER XII.

INTRACRANIAL COMPLICATIONS.

BRAIN ABSCESS. THROMBOSIS OF THE SIGMOID AND OTHER SINUSES. LEPTOMENINGITIS.

As a result of chronic purulent otitis media and mastoid inflammation, and, occasionally, of acute purulent otitis media, the thin roof of the tympanum or antrum is liable to be perforated through osseous erosion, and in consequence there follows either a collection of pus in the middle cranial fossa between the bone and the dura—an extradural abscess—or else an abscess forms in the temporo-sphenoidal lobe. When erosion takes place in the bony wall of the mastoid process, which lies over the sigmoid sinus, a collection of pus usually forms about the sinus, and the latter becomes surrounded with granulation-tissue; or a thrombus forms in the sigmoid sinus itself, or else an abscess is developed in the cerebellum.

Before erosion takes place the tympanic roof or inner wall of the mastoid process becomes very thin and of a dark color. The minute veins are often thrombosed, and, owing to the ulceration of the bone, granulations are thrown out from the external surface of the dura or from the wall of the sigmoid sinus, as the case may be, which is nature's method of resisting the further advance of the disease. Absorption of the carious bone takes place at the same time. One should be very care-

ful, therefore, in the removal of polypi from the tympanic cavity, especially when the disease has existed for a long time, for fear of wounding the dura, which rests on the tympanic roof. In the same way, the sigmoid sinus may easily be injured in scraping away softened bone and granulations in operations on the mastoid process. After removing granulation-tissue from the dura mater one is apt to find the latter considerably thickened (see Fig. 118). Granulation-cells and a plastic material are often found, which project upward into the brain substance and become adherent to the pia mater, so that the veins and lymphatics of the pia are liable to be affected. Very small thrombi are undoubtedly carried by the smaller branches of the cerebral arteries which supply the temporal lobe, and in this way give rise to cerebral abscess. The morbid products may also reach the brain tissue by means of the perivascular sheaths. Occasionally it is impossible to find any aperture in the bone, although the latter may be very much discolored.

Cases have been reported, especially in children, in which infective processes have reached the dura from the middle ear by means of the petro-squamosal or other sutures, and in such instances the vessels have been thrombosed. In tubercular disease the entire petrous bone may become carious and nothing remain of it except a thin shell, and in consequence of such disease, tubercular meningitis may develop. When erosion of the bone takes place, either through the tympanic roof or through the inner wall of the mastoid, the patient may have a suppurative leptomeningitis, together with an extradural or cerebral abscess. The roof of the tympanum may be eroded at the same time that the sigmoid

sinus is involved, and, as a result, there may be an abscess in the temporo-sphenoidal lobe, together with an infective thrombus in the sigmoid sinus. When all these complications are present in one case, there is great difficulty in making a diagnosis. The plate of bone separating the sigmoid sinus from the mastoid cells is extremely thin, so that erosion of the bone is very likely to be observed at this point. The bone first becomes discolored and softened, and then granulations are thrown out. A collection of pus is found either between the sinus and the bone or behind the sinus, in the direction of the occipital bone. The walls of the sinus become thickened and a thrombus forms, the presence of which may not give rise to immediate mischief; while, on the other hand, in an incredibly short space of time the patient may have most serious and alarming symptoms of sinus thrombosis. If left to nature, the wall of the sinus may become softened, so that a minute perforation takes place. Hemorrhage does not follow, because generally the vessel is occluded both above and below the aperture. By means of the cerebellar veins the infection extends to the cerebellum, or emboli may be carried to the cerebellum through the smaller cerebellar arteries and their perivascular sheaths. As a rule, however, there is direct contact between the sigmoid sinus and the cerebellar abscess, and the infective processes extend from the dura in the same manner as in the case of an abscess in the temporo-sphenoidal lobe. In rare instances the pathogenic processes reach the cerebellum from the tympanic cavity through the internal auditory meatus along the sheath of the auditory or facial nerve. Beside an extradural abscess, or an abscess in the temporo-sphenoidal lobe or cerebellum,

an abscess may form between the dura and pia mater, which may become circumscribed through adhesions; in consequence of which there may be inflammation and possibly ulceration of the brain-tissue, with softening. The pus may escape through a perforation in the membranes of the brain to the outer surface of the dura or through the roof of the tympanum.

As a rule, when an abscess has formed in the temporosphenoidal lobe, there will be found an opening at the base of the brain which communicates with it, so that abscesses occurring in this portion of the brain are caused generally by direct extension from the seat of the disease—a point to be remembered, for in operating the surgeon should always try to reach the seat of the infective process.

CEREBRAL ABSCESS.

Under this heading should be included (1) extradural abscess, (2) intradural or cerebral abscess—(*a*) *cerebral*, and (*b*) *cerebellar*.

Extradural Abscess.

As already mentioned, the pus in an extradural abscess may be found between the dura mater and tympanic roof or about the sigmoid sinus after the infective process has extended from the mastoid cells. Frequently, when operating for mastoid inflammation, the surgeon finds a carious opening just over the sinus, even when the symptoms indicated only mastoid disease.

A boy, ten years of age, who came to my clinic, had a careworn expression, cold hands, a rapid and bounding pulse, and a temperature of 103° F. He had no

distinct chill, but on the preceding day had had chilly sensations. He gave a history of an occasional discharge from both ears for several years, and during the preceding three days had had severe pain in the left mastoid process. He heard the watch only by contact. There was a distinct swelling over the left mastoid, with a point of fluctuation just over the antrum. The mastoid cells were opened, and pus and softened bone were removed from the entire process. The sigmoid sinus was not involved. A probe was introduced into the middle cranial fossa through a carious opening in the tympanic roof. The incision was carried upward, and the bone cut away with rongeur forceps and chisels, so as to expose the dura and evacuate the pus. While the temperature was high in this case, it does not follow that this is the rule, for frequently there is little if any fever. A short time ago a patient came under my care who had a temperature but slightly above normal. He had had a chronic purulent otitis media of some years' duration. He consulted me on account of a sudden pain in the mastoid on the same side. He also had slight facial paralysis. An immediate operation was performed, and not only were the mastoid cells found filled with softened bone and pus, but there was also a large carious opening through the tympanic roof. This opening was enlarged and the pus evacuated. Recovery followed. It should not be forgotten that a leptomeningitis or phlebitis of the sigmoid sinus may complicate a case of extradural abscess.

Symptoms. There are no distinct symptoms by which the surgeon can make a diagnosis of extradural abscess before operation, for, as previously mentioned, osseous erosion in the tympanic roof or in the thin plate of bone

covering the sigmoid sinus frequently is discovered only at the time of opening the cells. One should remember that such an accident is especially liable to occur when streptococci are present in the discharge or the patient has tuberculosis or one of the exanthematous diseases. Most cases of extradural abscess occur in connection with chronic purulent otitis media. It may be stated, however, that cases have been observed in which erosion had taken place during an attack of acute otitis media. One may suspect the existence of an extradural or subdural abscess when the discharge suddenly becomes less or ceases altogether, and the patient shows signs of meningeal irritation. A fatal result may follow from basilar meningitis or abscess of the brain.

Treatment. An incision behind the auricle should be made as in the operation for mastoid disease. If an opening has occurred in the tegmen tympani or tegmen antri, the incision in the soft tissues should be carried upward and curved forward around the upper border of the auricle (from point 2 to point 4, and from point 2 to point 5, Fig. 108), thus displacing the latter forward, so that the surgeon will be enabled to cut away the bone with forceps and chisels, and thus have an opportunity of exploring the middle cranial fossa and of removing all carious bone. A narrow strip of iodoform gauze should be inserted through this opening and the wound allowed to fill up with granulation-tissue. The mastoid cavity should be treated in the manner already described (see chapter on Mastoid Diseases). The surgeon should never inject any solution into the cranial cavity unless he is certain that there is no opening in the dura. The edges of the wound should be washed with a saturated boric acid solution.

Intradural or Cerebral Abscess.

Abscesses of the brain occurring as a result of middle-ear inflammation are found usually in either the temporo-sphenoidal lobe or the cerebellum. Cases of extradural abscess are observed much more frequently than abscesses either in the temporo-sphenoidal lobe or cerebellum. Cerebral abscess almost always occurs in

Fig. 117.

1. Primary abscess cavity connected with sinus in tympanic roof. 2. Secondary abscess cavity in temporo-sphenoidal lobe. Both abscess cavities found free from pus at autopsy.

connection with chronic purulent otitis media, although a case[1] of temporo-sphenoidal abscess has been reported which followed acute otitis media (see Fig. 117). In this case the disease was limited mostly to the attic, the acute otitis media being of only eight weeks' duration. Several incisions were made in Shrapnell's membrane,

[1] Bacon: Archives of Otology, 1896, vol. xxv. No. 3.

which not only gave exit to a few drops of pus, but afforded at the same time great relief to the patient. He finally consented to enter the hospital and receive proper treatment, but only after serious symptoms had manifested themselves. An operation was performed and an abscess in the temporo-sphenoidal lobe evacuated. He died, however, two hours after the operation,

FIG. 118.

1. Carious opening in tympanic roof. 2. Thickened dura turned back, but when in position covers carious opening.

apparently from shock (see Fig. 120). At the post-mortem a carious opening was found in the tympanic roof (see Fig. 118), with thickened dura over it, and a small sinus in the latter leading to a small cerebral abscess cavity, which was empty. Another abscess cavity was found at the inner half of the third temporo-sphenoidal convolution (Fig. 117). This cavity was also free from pus, as it had been evacuated at the operation. There was probably some connection between the

two abscess cavities, although none could be detected with a probe. In three cases [1] of abscess in the temporo-sphenoidal lobe and one in the cerebellum in which an operation was performed, all were due to chronic purulent otitis media.

Fig. 119.

Photograph of brain showing (1) a portion of hernia cerebri and (2) abscess cavity in temporo-sphenoidal lobe.

Symptoms. The first stage of abscess is marked usually by irregular symptoms, such as irritability, pain, nausea, or vomiting, and frequently the discharge from the ear becomes scanty or stops altogether. Change in a patient's disposition—viz., a talkative person becoming morose, or *vice versâ*—is another symptom of importance. The patient frequently complains of chilly sensations. As the disease progresses the cerebration becomes

[1] Bacon: New York Eye and Ear Infirmary Reports, January, 1897.

dull and the patient appears stupid and restless. When pus has formed in the brain, the most important symptoms are a distinct lowering of the pulse-rate and a tem-



Temperature-chart of a typical case of abscess of the brain without complications, following acute purulent otitis media.

FIG. 120

perature either slightly above normal or subnormal (see Fig. 120). There are exceptions to this rule, for in a

case[1] of temporo-sphenoidal abscess reported by me the pulse was rapid and the temperature very high with marked fluctuations, so that thrombosis of the lateral sinus was suspected (see Fig. 121). The patient was a man, thirty-two years of age, who had had a chronic otorrhœa on the left side for at least fifteen years, as a result of measles. He received no treatment to speak of during this time, until his family physician was suddenly sent for. The patient had at that time a temperature of 104° F., an intense headache, and marked pain in the ear. Later he had a severe convulsion lasting twenty minutes, with violent muscular twitching and frothing at the mouth. He also had chills. The symptoms were characteristic of sinus thrombosis. At the operation, however, the sinus was found normal, but there was a large abscess in the temporo-sphenoidal lobe. This case proves that, exceptionally, a patient with a cerebral abscess may have chills and a high temperature, as well as convulsions, so that a differential diagnosis between abscess of the brain and sinus thrombosis becomes extremely difficult.

When aphasia develops in a patient with a chronic purulent otitis media of the left side, it is fairly certain, if the patient is right-handed, that there is a collection of pus in the temporo-sphenoidal lobe. This symptom is of the highest diagnostic value—*optical aphasia*, so called and described by different writers. The patient when shown an object knows it, but is unable to name it; but he is able to recognize and understand its name, and will often describe the object. He can sometimes name the object if allowed to handle or smell it. In the case referred to, where the temperature was

[1] New York Medical Journal, August, 1896.

FIG. 121.



Temperature-chart of an uncomplicated case of abscess in the temporo-sphenoidal lobe, as a result of chronic suppurative otitis media. Course of the temperature very unusual, suggestive of sinus thrombosis.

high, the patient had well-marked aphasic symptoms. Beside the difficulty in naming an object, he could not readily call to mind the appearance of the object. Starr has called this symptom *visual amnesia*, and says "it is due to a suspension of function of an association tract lying in the temporo-occipital lobes and occupying the white matter deep under the cortex. It lies in a position in which it is almost inevitably affected by abscess of the temporal lobe developing after otitis media. In left-handed persons it will occur in abscess of the right side." Severe headache is also an important symptom. Inflammation of the optic nerve may or may not be present. It is observed more frequently in cases of cerebellar than of temporo-sphenoidal abscess. According to the statistics of the Halle Clinic as given by Hansen,[1] optic neuritis in cerebral abscess was bilateral in 87.50 per cent. of the cases while the eye of the affected side alone was involved in 12.5 per cent. The mastoid cells often are found dense and like ivory in cases of cerebral abscess. This seems to me an important point, for in such instances the pus is likely to perforate the tympanic roof, and thus cause an extradural abscess or an abscess in the temporo-sphenoidal lobe. Percussion of the skull on the affected side frequently causes pain, and Macewen has called attention to the fact that there is a difference in the percussion note on the side of the head in which the abscess is located. Other symptoms sometimes noticed are digestive disturbances, increase of reflexes in the limbs of the side opposite to the abscess, a difference in the size of the pupils, and facial paralysis.

In *cerebellar abscess* the most characteristic symptoms,

[1] Archives of Otology, October, 1904.

beside slow pulse and low temperature, are severe headache, nausea, vomiting, vertigo, a staggering gait, and facial paralysis. Facial paralysis may be due to pressure on the nerve in the pons. It should be remembered that the nerve may be affected in its course through the middle ear. I have seen a case in which the pulse was slow and the patient had vertigo, vomiting, deafness, severe headache, etc., so that a diagnosis of cerebellar abscess was made. When the antrum was opened, the semicircular canals were carious, which fact accounted for the symptoms. It is very important to examine thoroughly the external auditory canals in all cases of suspected cerebral abscess; for instance, in the case of a child who was brought to my clinic a short time ago, and who had characteristic symptoms of cerebellar abscess, including choked disks with retinitis, removal of granulations from the middle ears, with the establishment of proper drainage, effected a cure. The choked disks and retinitis disappeared.

I have reported a case of cerebellar abscess[1] in which I not only removed a button of bone at point *G*, Fig. 122, having first laid bare with chisels the sigmoid sinus, but I also trephined the skull at point *C*, to investigate the temporo-sphenoidal lobe and the upper wall of the tympanic cavity. The case was that of a male, aged thirty-one years, who had had a purulent discharge from the right ear for five years. When admitted to the hospital the prominent symptoms were nausea, vomiting, and vertigo. The vertigo was not only present when lying down, but was much more marked when he tried to stand. He also complained of severe pain in the head. He had a staggering gait from the first with,

[1] American Journal of the Medical Sciences, August, 1895.

an inclination to fall toward the left. Other symptoms were facial paralysis (right side), œdema of the right optic nerve, slow pulse, and a temperature but slightly above normal. At the autopsy an abscess was

FIG. 122.



A. External auditory meatus. *B B'*. Reid's base-line running from lower margin of the orbit through the centre of the external auditory meatus. *C*. Trephine opening, ¾ inch in diameter, to expose temporo-sphenoidal abscess, centre-pin of trephine being placed 1¼ inches above the centre of the external bony meatus. *D*. Point at which the mastoid antrum should be opened. *E E'*. Position of sigmoid sinus. *F*. Trephine opening, ¾ inch in diameter, for exposing sigmoid sinus, centre-pin of trephine being placed at a point 1 inch behind and ¼ inch above centre of external bony meatus. *G*. Trephine opening for cerebellar abscess, centre-pin being placed 2 inches behind and ¼ inch below centre of external bony meatus.

found in the right cerebellar hemisphere (see Plate XI.).

Other symptoms of cerebellar abscess sometimes observed are: slowing of the respiration; Cheyne-Stokes respiration; repeated yawnings; slowness of

PLATE XI.

Abscess of cerebellum secondary to chronic suppurative otitis media. The right cerebellar hemisphere has been divided and the lower half removed, exposing the abscess cavity and its wall (*a*) and the area of hemorrhagic softening posterior to the abscess (*b*).

cerebration and general apathy; irritability and intolerance of light; delirium; rigidity of the neck and retraction of the head; clenching of the teeth from rigidity of the muscles; nystagmus; motor or sensory paralysis. If the medulla is affected at a point low enough, the paralysis may be unilateral and on the same side as the abscess. If, however, the pons is involved more on the side of the abscess, the paralysis will be apparent on the opposite side.

When there is a cerebellar abscess together with sinus thrombosis there may be rigors or chills, with high temperature and tenderness along the course of the jugular vein, if that vessel is affected. Suppurative leptomeningitis is another complication which may occur and thus render the diagnosis more difficult. Diffuse leptomeningitis gives rise to high temperature (without remissions, which are characteristic of infective sinus thrombosis), a rapid pulse, general irritability, and with the special senses very acute.

During the past few years considerable attention has been devoted to the subject of *leucocytosis*, a term used to signify an abnormal increase in the number of leucocytes present in the blood of patients suffering from various affections, with the result that it has been shown to exist in suppurative affections of the ears. The subject of leucocytosis in connection with suppurative diseases of the middle ear and their complications is in a somewhat chaotic condition at present, and the exact degree to which it can be relied upon is questionable. Daily examinations of the blood should be made in order to determine whether the leucocytosis is increasing or decreasing, and what relative changes are taking place in the cell percentage. For example, if the

leucocytosis is falling, and the polynuclear percentage is above normal (68) and rising, the prognosis is not so good as when both are falling, or when the leucocytosis is increasing and the polynuclear percentage is falling. Or as tersely put in Sondern's axiom "the increase in

FIG. 123.

Left temporal bone of a patient who died of cerebral abscess.

a, *a*. Openings in the sclerosed mastoid process, made with chisels. *b*. Large carious opening leading into the middle cerebral fossa. *c*. Zygoma.

the relative number of polynuclear cells is a direct indication of the severity of the toxic infection and the degree of leucocytosis an absolute indication of the body resistance toward the infection."

But no hard and fast rules can be laid down at present to aid the aural surgeon. According to a study of

eighty-nine cases of acute and chronic purulent otitis media with and without complications made by Dixon at the New York Eye and Ear Infirmary recently, it was found that the leucocytosis in those cases with mastoiditis as a complication ranged from 5,000 to 17,800, and the average polynuclear percentage was within normal limits. It was not until the more serious complications occurred, such as epidural abscess, sinus thrombosis and intracranial invasion, that the blood count began to have any very marked significance.

The leucocytosis occurring in the course of intracranial complications does not appear to be so high as in pneumonia. For instance, a sudden and sustained rise to about 20,000 or over with a polynuclear count above normal and rising has been found in cases of brain abscess, while 22,000 and over with a rising polynuclear count has been found in meningitis, and above 30,000 is an indication of pneumonia even in the absence of physical signs at the time.

While the above would appear to be true in the main, exceptions occur, and further investigation is imperative before we can rely with much confidence on the result of blood examinations in otitis and its complications, no matter how valuable it may be in general surgery; and it seems that the most important point to start from is a knowledge of the normal average leucocytosis and cell percentage in each individual. Impracticable as this may appear, I feel this to be the only rational basis from which to draw conclusions.

The reader is referred to works dealing with the subject of blood examinations for information as to the various methods of determining the number of cells, the hæmoglobin percentage, etc.

In 1888 an operation for cerebral abscess following disease of the ear was performed by Dr. Robert F. Weir and myself. The patient[1] was under my care at the New York Eye and Ear Infirmary for chronic suppurative otitis media, with extensive necrosis of the temporal bone. He had had a purulent discharge from the ear for the preceding two years. The mastoid process was opened in two places (see Fig. 123), and was found very dense, but it did not contain pus. Subsequently an abscess was located in the temporo-sphenoidal lobe and the pus evacuated. The patient's condition improved for a time, but later he passed into a comatose condition. The immediate cause of death was secondary hemorrhage from a vessel in the brain.

When not interfered with by operative measures, a cerebral abscess terminates, as a rule, in death. The patient may pass gradually into a comatose condition, while the pus escapes on to the surface of the brain or bursts into the ventricles. When the pus reaches the surface of the brain a new inflammatory action is set up, leading possibly to acute leptomeningitis. As a result the temperature becomes high and the pulse rapid. Other symptoms are vomiting, restlessness, rigidity of the limbs, great weakness, and clonic spasms. When the pus bursts into the ventricles the patient's condition changes at once. The temperature rises suddenly to 104° or 105° F.; the pulse becomes very rapid; the respirations are increased in frequency; muscular twitchings occur, followed by convulsions, coma, and death.

Abscesses have become encapsulated and remained quiescent for years without giving rise to serious trouble,

[1] Transactions of the American Otological Society, 1888.

but such instances are extremely infrequent. Cases have been reported also in which abscesses in the temporo-sphenoidal lobe have drained through a carious opening in the tegmen tympani, and the pus has escaped through the external meatus. A cerebellar abscess has also been reported where the pus forced its way through a carious opening in the sigmoid groove, the sinus having been destroyed. Occasionally, but very rarely, an abscess may drain in this way and be followed by recovery. One should never for a moment anticipate such a result, for the time to operate successfully is at an early date, before the patient has passed into a comatose condition.

It is frequently very difficult to make a differential diagnosis between cerebral abscess, sinus thrombosis, and meningitis, as at times one or more of these diseases may occur in the same case.

It is not often that a patient has an abscess in the temporo-sphenoidal lobe and also one in the cerebellum, but such a case[1] came under my observation. The patient was a woman, thirty-two years of age, who had suffered from a chronic discharge from the right ear, and whose hearing in consequence had been defective for some years. For a month before she came under my care she complained of pain in the right ear and severe pains in her head. She also had vertigo and general weakness, and nine days before admission to the hospital she had two distinct chills on the same day, followed by vomiting. The mastoid cells were opened, and were found diseased. Later the temporo-sphenoidal lobe was investigated on the right side for an abscess,

[1] Transactions of the American Otological Society, 1900, vol. xxx. part iii.

on account of the following symptoms, viz., paralysis of left abducens; paralysis of left side of face; slight left hemiparesis; moderate left hemianæsthesia; left homonymous hemianopsia; choked disks on both sides. The bone was cut away from the external auditory canal in a direction upward for a distance of two inches, so that the brain in the middle fossa could thoroughly be exposed. The dura was found thickened and bulging, and a minute opening was discovered about an inch and a half above the external auditory canal, leading in a direction inward, upward, and backward for at least three inches, and disclosing an abscess cavity. The patient seemed to do well immediately after the operation. A few days later, however, she developed muscular twitchings of the face and legs. She also became slightly delirious and tried to remove the bandages. The muscular twitchings persisted, and affected the right side of the face, right arm, and leg. At times the patient for the next few days was rational and quiet; then again she would scream out and try to tear off the bandages. During this time the temperature varied from 98.4° to 102° F., pulse 80–110, respirations 18–24, showing a course exceptional to that usually observed in cases of brain abscess. At the autopsy nothing unusual was observed about the meninges. There was, however, a large abscess cavity in the temporo-sphenoidal lobe, which had been well drained. On opening the right lobe of the cerebellum, an abscess was disclosed, containing about half an ounce of dark foul-smelling pus. It appeared to have had its origin in the dentate body, which it destroyed. It then extended across to the opposite lobe, which it invaded to the extent of half an inch. No communication whatever could be found

between the abscess in the cerebellum and that in the temporo-sphenoidal lobe. This case demonstrated the fact that a chronic purulent otitis media may remain quiescent for many years, when suddenly the patient complains of excruciating pains in the head and ear, and general weakness, followed by vomiting, vertigo, chills, and an intracranial complication.

It is impossible to know the percentage of recoveries of brain abscesses, for, as a rule, all successful cases are reported, but the same cannot be said of unsuccessful ones. According to Kœrner's records recovery has occurred in 50.2 per cent. of cases of cerebral abscess, and in 52.8 per cent. of cerebellar abscess. In ninety cases reported by Kœrner, sixty were males. Of one hundred and thirty-nine cases of sinus thrombosis, including some that have not been recorded as yet, ninety-five terminated in recovery. Of forty-two cases reported to July, 1900, at the New York Eye and Ear Infirmary, there were thirty-three recoveries and but nine deaths, so that the surgeon is fully justified in operating. An exploratory operation whereby the dura is exposed and a needle introduced into the lateral lobe of the cerebellum or temporo-sphenoidal lobe is attended with but little risk to the patient if aseptic precautions are carefully followed out.

Treatment. In all operations on the mastoid antrum and cells, and when opening the cranial cavity, the most thorough antiseptic precautions should be followed. The ear should thoroughly be syringed with bichloride solution, 1 : 5000, and then carefully dried and a strip of iodoform gauze pushed down into the meatus. The hair should be shaved about the ear for a considerable area. The mastoid region and the surrounding parts

should then be scrubbed with soap and water, and afterward with pure alcohol, or sulphuric ether, or bichloride solution, 1 : 3000. Dry sterile towels should then be placed over the patient's face and shoulders, and a sterile towel wrapped around the head and fastened with a safety pin or artery forceps. All instruments should be sterilized. The blades of the scalpels should be dipped into absolute alcohol, and carbolic solution, 1 : 20.

The surgeon should open the mastoid antrum first (see Fig. 100), and make this the starting-point of the operation, for it is in this cavity that the micro-organisms exist, and where they rapidly fructify as long as the soil is fertile. Antiseptics injected into the middle ear do not reach them. Beside scalpels and retractors, it will be necessary to have a mallet, gouges and chisels of different sizes, a periosteal elevator, forceps for controlling hemorrhage, rongeur forceps of different sizes, probes, and exploring needles. After removing all cheesy matter and granulations from the antrum, and excising the ossicles if carious, the condition of the tegmen tympani and tegmen antri should be investigated. If a sinus is found leading to the middle cranial fossa, it should be enlarged, and pus, if present (extradural abscess), evacuated. The surgeon should examine the condition of the dura before injecting any solution into the extradural cavity. The latter cavity may be washed out safely with a saturated boric acid solution, provided no opening exists in the dura which leads into the pia mater or brain itself. It is best to do but little irrigation, for fear of disseminating the morbid products. The cavity should be dusted over with equal parts of boric acid and iodoform, and iodoform gauze

should be inserted into the wound, so that it may fill up with granulation-tissue. When pus is seen to ooze from an opening in the dura, or one has reason to suspect a temporo-sphenoidal abscess, the surgeon should extend the previous incision (for exposing the antrum) in a direction upward and then forward around the upper border of the auricle, thus dividing the temporal artery (see Fig. 108, from point 2 to point 4). The hemorrhage is controlled easily by means of a ligature. Another incision should be made in a direction upward (from point 2 to point 5), and the flaps dissected so as to give the surgeon plenty of room. If the bone just above the external meatus is not too dense, it should be cut away rapidly with large chisels and bone forceps, and the opening enlarged so that the dura can be fully exposed. In some cases, especially where the bone is very dense or the patient is in such a condition that it is necessary to explore the temporo-sphenoidal lobe as rapidly as possible, a disk of bone should be removed with a trephine, the centre-pin of the trephine being placed at the point *C*, Fig. 122, a distance of an inch and a quarter above the centre of the external bony meatus. The trephine opening should be at least three-quarters of an inch in diameter. It is probable that most abscesses in this region are the result of inflammation extending from disease of the antrum. It is, therefore, best to attack the abscess cavity at this point (antrum), and to enlarge the opening in a direction upward, in order to secure the best possible drainage. From this region, also, a large exploring needle may be introduced into the temporo-sphenoidal lobe if a brain abscess is suspected, after a circular flap has been made in the dura. As the pus is often very thick I

have been in the habit lately of introducing a director in different directions in the brain substance. The instrument devised by Horsley is better adapted for this purpose than the exploring needle, as it consists of two blades, resembling an ear speculum. which can be separated after introduction, thus affording the surgeon a view of the abscess cavity. The blades can be pushed gently through brain tissue without the use of a knife, until at least the capsule of the abscess is reached. After the abscess cavity has been detected, a free opening should be made in the dura with a scalpel, and if possible the surgeon should enlarge the incision in the brain tissue with a director, for fear of injuring some bloodvessel. A free incision may, however, be made in the temporal lobe with a scalpel if the bloodvessels are carefully avoided.

Cerebral pulsation may not be evident if the abscess is small and deeply seated. There may, however, be entire absence of pulsation even if the abscess is near the surface and the dura protrudes through the bony opening. After making an opening down to the abscess, great care should be exercised in evacuating the pus and sloughs. If possible, the little finger should be introduced to remove the sloughs, or a Volkmann's spoon may be used if great care is taken and light is thrown with a head-mirror into the abscess cavity. The surgeon should endeavor to remove all the sloughs, otherwise the cavity is likely to refill with pus. After the abscess has been opened the encephaloscope devised by Dr. Fred. Whiting, of New York, can be used to great advantage. It is much like an ordinary ear speculum but larger (Fig. 124), the extremity being closed by an obturator so that it can be pushed into the

abscess cavity without effort, and thus any laceration of the brain substance can be avoided. The instrument is used with a head-mirror. It should be introduced first for a short distance only and then the obturator should be removed with a slight rotary movement, and at the bottom of the encephaloscope can be seen a meniscus which points toward the abscess so that the direction of the abscess path can readily be determined. Accord-

Fig. 124.



Whiting's Encephaloscope.

ing to Whiting, the surgeon can decide whether he is dealing with a single abscess cavity or a large abscess path communicating with several other abscesses—and mechanically the instrument enables the operator to introduce gauze into the abscess cavity without any risk of tearing the abscess wall and carrying infection into healthy brain tissue. The cavity should then be washed out most gently with a boric acid solution or a 1 : 100

carbolic solution by means of a double canula, the tube carrying away the fluid from the abscess cavity being a little larger than the other one. But little force should be employed in syringing, otherwise great damage may be done to the brain tissue. After washing out the abscess cavity, strips of iodoform gauze should be packed in loosely for the purpose of draining it. These may have to be changed every day at first, but later less frequently. Under such frequent dressings, boric acid gauze should be substituted, for fear of the toxic effect of iodoform. In the case of an acute abscess, or when all septic matter has been removed, an absorbable decalcified chicken-bone drainage-tube may be introduced and stitched to the skin, so as to keep it in position. Outside of the abscess the wound, after aseptic irrigation, should be covered first with several layers of iodoform gauze, and then with a pad of bichloride gauze. Outside of this a thick layer of absorbent cotton should be placed, and the whole should then be bandaged.

After several days, in the case of a chronic abscess, when the walls have become firm, a 1 : 5000 bichloride solution may be injected into the abscess cavity to advantage. In a case of mine which recovered, and in which streptococci were found, great improvement ensued after using a bichloride solution.

In the case of a cerebellar abscess, it is always advisable to expose first the sigmoid sinus, as the latter frequently is thrombosed. A horizontal incision should then be made in the soft parts and carried backward (see Fig. 108, from line of original mastoid incision to point 3), so as to give the surgeon plenty of room not only to lay bare the sinus, but also to cut away with

rongeur forceps the bone lying to the outer and external aspect of the sinus, and reach the abscess in that way. Otherwise the treatment is identical with that already described (abscess in the temporo-sphenoidal lobe).

If the bone is very dense, and for any reason it becomes necessary to explore the cerebellum for an abscess without first investigating the sigmoid sinus, the following method of procedure should be followed: A button of bone should be removed with a trephine (see

FIG. 125.



Author's rongeur forceps.

Fig. 122) by placing the centre-pin of the instrument at a point, *G*, a distance of two inches behind and one-quarter of an inch below the centre of the external auditory meatus. As shown in the same figure, Reid's base-line extends from the lower margin of the orbit through the centre of the external auditory meatus. The directions which formerly were given for exposing the sigmoid sinus by means of the trephine were to remove a button of bone by placing the centre-pin of the trephine at point *F*, a distance of one inch behind and one-quarter of an inch above the centre of the external meatus.

Macewen[1] says that the intermediate path in an

[1] Transactions of the Seventh International Otological Congress, Bordeaux, 1904.

abscess of the brain as well as the primary focus must be eradicated—otherwise a secondary abscess will result. Therefore, whenever a delay of twenty-four hours is possible after the mastoid operation and exposure of the dura, he introduces small pieces of gauze soaked in a 5 per cent. solution of carbolic acid between the dura and the bone to induce adhesions. These are formed generally in twenty-four hours. The abscess is then opened.

THROMBOSIS OF THE SIGMOID AND OTHER SINUSES.

Before describing the symptoms of sinus thrombosis, it is important to review in a general way the gross anatomy of the brain, particularly as to its venous supply.

The superior cerebral veins collect the blood from the upper portions of the cerebrum, and after anastomosing with one another and the inferior cerebral veins, empty into the superior longitudinal sinus. The inferior cerebral veins enter the lateral, cavernous, and superior petrosal sinuses. The middle cerebral vein, after being joined by branches from the frontal and temporo-sphenoidal lobes, pours its blood into the cavernous sinus.

The *sigmoid sinus* is a continuation of the lateral sinus, and is that portion which lies in the sigmoid groove. It is joined by the inferior petrosal sinus and forms the *internal jugular vein.* The sigmoid sinus receives blood from the superior petrosal and occipital sinuses, also from the superior longitudinal and straight sinuses by means of the lateral sinus. The lateral and sigmoid sinuses, together with the internal jugular vein, are of especial interest to the aural surgeon, for it is in these

vessels that most frequently a thrombus is found when the middle ear and mastoid cells are involved, although the other sinuses that are in communication with them are liable to be affected, especially the superior and inferior petrosals.

Small veins from the mastoid cells and tympanum also enter the sigmoid sinus, more especially at that point called the knee or the anterior convexity of the sinus. A *mastoid vein* joins the sigmoid sinus at its posterior aspect, and frequently is thrombosed when the former is involved. A thrombus forming in the sigmoid sinus may not only extend to the petrosals, but even to the cavernous sinus as well. Although not constant, there is a communication between the vessels of the external portions of the skull and the sigmoid sinus by means of the mastoid and posterior condylar veins. The latter connect the sigmoid sinus with the vertebral veins and the deep veins in the posterior part of the neck.

Lymphatics. The lymphatics in the scalp enter the mastoid, parotid, and occipital lymphatic glands. Some of the lymphatics of the face are superficial and others deep. The deeper ones terminate in the internal maxillary glands, while the superficial ones join the submaxillary and parotid lymphatic glands.

The lymphatics within the cranial cavity are situated in the pia mater and choroid plexus, taking their exit from the cranial cavity along the internal carotid and vertebral arteries and internal jugular vein, and join the deep cervical glands, so that the latter glands are liable to be enlarged when there is some irritation or inflammation within the cranium. Being deeply seated, they are liable to escape detection.

FIG. 126.



Cerebra veinsl and sinuses. (From MACEWEN.)

From this brief review of the anatomy it will be possible to understand more readily the pathology and symptomatology of sinus thrombosis.

A thrombus may be *primary* or *marasmic*, as well as *secondary* or *infective*. The marasmic is said to occur almost always in the longitudinal sinus, very seldom in either the lateral or cavernous sinus. It is observed in the very young and in the aged, and is developed generally during the course of some wasting disease, such as chronic diarrhœa, pneumonia, tuberculosis, and carcinoma. The heart's action becoming weakened and the blood being thickened, there is a tendency for coagulation to occur.

It is the *infective* or inflammatory thrombus that concerns the aural surgeon, and which is due to the presence of a pathogenic microörganism, the sigmoid sinus being affected most frequently on account of its anatomical relation to the temporal bone, more especially the mastoid cells. Infective thrombosis is observed seldom in children and aged persons. It may be due to fractures of the skull, scalp-wounds, anthrax affecting the lips, mouth, or orbital or nasal cavities, and erysipelas of the face and head. Middle-ear disease and mastoid inflammation are the most frequent causes. The thrombosis is due to the presence of pathogenic microörganisms, such as the staphylococcus, the streptococcus, the pneumococcus, or the bacillus mucosus capsulatus. They may exist alone or together in the same case. Experience teaches that the streptococcus is the most virulent and gives rise to serious symptoms, and the bacillus mucosus capsulatus is the most insidious. The infection may be due to direct extension of the inflammation in the bone to the sinus walls, whereby they become

inflamed and thickened, and a thrombus forms in the interior of the sinus. There may, however, be fissures in the tympanic roof or antrum through which the infection travels, or small veins in the bone, joining with the sigmoid, become thrombosed and convey the morbid products in this manner. The most frequent cause, however, is by direct contact with the diseased bone.

When the sinus first is affected the endothelium becomes swollen and desquamates, and coagulation takes place. The deposit of fibrin may be limited to one side of the wall (parietal clot), or the whole circumference of the vessel may be involved (obstructing thrombus). After this the clot, as a rule, breaks down, so that pus is formed, and, being filled with microörganisms, the latter are likely to be carried into the general circulation. Ulceration may take place on the wall of the sinus, but hemorrhage does not usually follow, because the thrombi formed above and below this point are firm enough to prevent the escape of blood. The infective material may also be carried by the tributary veins in the neighborhood of the thrombus into the general circulation. In some cases a partial re-establishment of the current takes place through the thrombus, by which means the pathogenic material is swept into the general blood-current. Another way in which general infection takes place is by the escape of morbid products through the sinus walls into the surrounding parts, so that another focus of infection is established whereby micro-organisms are carried into the circulation through the perivascular spaces and congested vessels.

It is probable also that the pyogenic micro-organisms may give rise to general infection through absorption by means of the lymphatics. The streptococcus is found

most frequently in the lymph-vessels. As a result of thrombosis of the sigmoid sinus the same phenomena frequently occur in the internal jugular vein, especially its upper portion, as well as in the superior petrosal sinus and the anterior and posterior condylar veins. A large number of internal jugular veins, excised at the New York Eye and Ear Infirmary, have been examined and found to contain microörganisms in the wall, rarely in the clot itself, as shown by sections.

When a thrombus forms within the sinus certain changes take place in its walls, owing to a peripheral exudation of plastic material which would tend to strengthen the weakened wall if the tissue should become changed into connective tissue. The same, if it occurred on the wall next to the brain cavity, would also help to prevent the occurrence of leptomeningitis through the formation of firm adhesions between the dura and membranes. Unfortunately, such is not the rule, as the plastic material is apt to break down and increase the dangers of infection. When, through ulceration of the sinus wall, an opening occurs, the pus may find its way through the mastoid foramen and form an abscess in the tissues in the posterior portion of the mastoid process. As a result of infection of the posterior condylar veins an abscess may develop beneath the deep fascia in the posterior cervical triangle. In consequence of involvement of the visceral layer of the sinus, acute leptomeningitis is the rule, although in some instances the meningitis becomes localized.

When the thrombus breaks down, the bone in its vicinity becomes of a dark-brownish or greenish color, very much softened and eroded. The discoloration or

pigmentation is so marked in some instances that the course of the lateral or sigmoid sinus becomes stamped on the cerebellum itself. When the thrombus disintegrates, the infective particles generally are transmitted first to the small vessels in the lungs, so that infarctions occur, and as a result there may be an infective pneumonia or gangrene. The pleura, pericardium, brain, kidneys, and liver may also be affected in the same way, or an abscess may form in the cerebrum or cerebellum as a result of thrombosis from extension of the infective processes.

Instead of disintegration of a thrombus, the latter, in rare instances, becomes organized; and in such cases the same reparative inflammation takes place as in other organs of the body.

When the infective process has been slight, the obliteration of the sinus may be established at that point by means of fibrous tissue, as the result of the formation of granulations on the walls of the sinus and adjacent dura, which penetrate into the sinus itself as well as the surrounding bony tissue. Such granulations tend to prevent the entrance of infective material into the sinus.

Symptoms. The symptoms should be divided into (1) local and (2) general or systemic.

1. *Local.* A profuse discharge from the ear often becomes diminished in quantity. Pain is generally present, being slight in some instances, while in others it radiates over the side of the head and is excruciating. As the mastoid cells usually are affected at the same time, there is more or less pain when pressure is made on the different portions of the mastoid process, and

especially in the neighborhood of the sinus and over that point where the mastoid vein enters the skull.

Œdema of the superficial tissues about the mastoid tip, extending to the region of the mastoid vein and also to the upper portion of the posterior cervical triangle (Griesenger symptom), with tenderness on pressure over this area, is another symptom of importance, although the same condition of affairs exists in uncomplicated mastoid disease, especially when perforation has occurred at the mastoid tip (Bezold perforation).

When the thrombus has extended to the cavernous sinus and the ophthalmic vein has become engorged, there is likely to be a certain amount of œdema of the eyelids on the corresponding side.

The enlargement of both superficial and deep cervical glands is a symptom that cannot be relied upon. In some cases it is present, but absent in others. When the mastoid vein has not become involved, and the thrombus is located in the lower portion of the sigmoid sinus, there may be an over-distention of the mastoid vein. In some instances it has been noted that the veins on the other side of the head were more dilated than usual, owing to increased pressure. Not only is there tenderness on pressure in the upper third of the posterior cervical triangle, but also very frequently along the course of the internal jugular vein, especially in its upper portion, when the latter has become involved. This is, however, one of the later symptoms. The patient generally has an anxious and pallid appearance, with beads of perspiration standing out on the face and forehead.

As a result of infective thrombosis, the tissues in the

cervical region become œdematous and filled with a plastic exudation, which is liable to break down and form a large abscess, unless the source of the infection is removed. Abscesses occurring in the later stages are observed at times under the deep cervical fascia, along the course of the internal jugular vein, in the deeper portions of the upper third of the posterior cervical triangle.

Neuroretinitis is noticed in some cases, more especially in the later stages, but it is not a constant symptom.

2. The *general* symptoms are more or less pain in the mastoid region radiating about the side of the head, a chill followed by a sudden rise in temperature to 105° or 106° F., nausea, vomiting, vertigo, and constipation. In the late stages there is apt to be diarrhœa, which is due to septic poisoning.

When a patient suffering from a purulent otitis media, and especially if there are symptoms of mastoid disease, has a sudden chill, followed by a rise in temperature (see Fig. 127), the indications point to the formation of a thrombus, provided a careful examination of the patient's general condition has been made and other diseases can be excluded. There are apt to be marked remissions in the temperature, the latter falling frequently to 99° F., but only to rise again. Rigors occur either early in the disease or may be a late manifestation, and are accompanied by profuse perspiration. The pulse and respiration generally are accelerated during the course of the disease, becoming markedly so during the late stage before a fatal termination. The patient generally retains consciousness to the end, provided there is no meningitis or brain abscess. In only 30 per

cent. of uncomplicated cases has loss of consciousness been noted. The tongue usually is coated, and the breath has a disagreeable, fetid odor. When general



Temperature-chart of a case of infective thrombosis of the sigmoid sinus, due to acute purulent otitis media.

FIG. 127.

systemic infection has taken place and the disease progress unfavorably, certain symptoms predominate, namely, pulmonary, abdominal, or meningeal, according to the organs affected.

In the case of a girl, aged nine years (see Fig. 127), who was admitted to the New York Eye and Ear Infirmary, December 13, 1898, a thrombus, which was partly broken down, was found in the sigmoid sinus eleven days after the first symptoms of acute otitis media developed. Eight days before coming to the hospital she complained of severe pain in the right ear, and was unable to sleep at night. When examined, there were marked redness and swelling of the tissues over the mastoid process, with a decided drooping of the soft parts of the posterior superior wall of the canal. There was bulging of the drumhead, but no perforation. On December 13th the drumhead was incised freely, and the mastoid cells were opened and found to contain pus. The sigmoid sinus was uncovered, so that it could be opened subsequently if further symptoms pointed to thrombosis. As the temperature rose to 106.2° F. two days later, and there had been a high temperature with marked remissions since the operation, it was deemed advisable to operate for sinus thrombosis on December 16th. A free incision was made in the sigmoid sinus and a large thrombus removed. The blood current was re-established from the upper end of the sigmoid sinus, but from the lower end, toward the bulb, the blood flowed but slowly even after a curette had been passed well down to the bulb. Although the lower end of the thrombus contained a drop or two of pus, it seemed better not to prolong the operation by ligating the internal jugular vein on account of the patient's condition, but to wait for twenty-four hours. As the patient was decidedly better on the following dry, no further operation was performed. The patient recovered.

Another case was that of a woman, twenty-four years of age, a primipara, who, expecting to be confined in about six weeks' time, came under my care on account of an earache on the left side, resulting from an attack of grippe. For two days she had severe pain and tinnitus, and then the right ear began to ache more severely than the left one. When admitted to the hospital she complained of having been dizzy for the preceding three days, and stated that night before last she had vomited. She had had no chills nor fever. Both drumheads were bulging, with tenderness on pressure over each antrum and mastoid tip. The mastoid symptoms were less marked on the left side than on the right. There was also partial facial paralysis on the right side. The right mastoid process was opened immediately and an examination, under the microscope, of the pus removed from the cells showed marked streptococcus infection. In the left mastoid the disease was even more advanced, having involved the tympanic roof, but when pus from this side was examined microscopically no streptococci were found. Three days later the patient had a severe chill with a sudden rise of temperature to 106.5° F., followed by profuse sweating. There had also been nausea and vomiting before the occurrence of the rigor. A diagnosis was made of sinus thrombosis, and the right sigmoid sinus was opened on account of streptococci having been found in the mastoid cells on this side. A large thrombus was removed, which also showed the presence of streptococci. The blood current was re-established from the upper end, but the blood did not flow from the bulbar extremity. The internal jugular vein was not ligated, as no evidences of the vein having been involved could

be detected, and the patient's condition did not seem to warrant further operation at that time. Four days later the temperature was normal. A few days after, she gave birth to an eight months' child. The patient made an excellent recovery and both mother and child were subsequently discharged from the hospital.

That the temperature does not always show marked oscillation is shown in the report of the following case of scarlet fever seen in consultation with Dr. Gruening. A double suppurative otitis media developed as a complication of the scarlet fever and an incision was made in each drumhead. The temperature remained normal for five days. Mastoid disease, however, followed and it was necessary to operate and remove a sequestrum from each side. The child did well for ten days and granulations began to form over each sigmoid sinus wall which had been uncovered, when the temperature suddenly rose to 105° F. It did not oscillate but remained high. There was no pneumonia. The sinus was opened on the side on which the wall was grayish and not as well covered with healthy granulations as the other. A septic thrombus was removed and the child recovered.

As to ligation of the jugular vein, Jansen[1] says that "numerous recoveries after incision of the sinus alone certainly go to prove that ligature is frequently unnecessary," and he further believes that one should ligate the jugular only when there is positive evidence of its being involved or when there is pyæmic or septic fever.

[1] Archives of Otology, 1901, vol. xxx., Nos. 4 and 5.

Körner[2] gives the following statistics:

Method of Operation	No. of cases.	Result doubtful.	Mortality.	Recovery
Ligation of jugular without opening of sinus.	9	0	4	5
Opening of sinus without ligation of jugular	132	0	55	77 = 58.3%
Ligation of jugular before opening of sinus.	94	2	36	56 = 59.6%
Ligation of jugular after opening of sinus.	68	1	29	38 = 55.9%

When the lungs are implicated, the first symptoms are dyspnœa and cough, with more or less localized pain in the chest. These localized areas of pain are due to infarctions. On examining the chest twenty-four or forty-eight hours later, one will hear moist râles. The sputum is rusty, contains bacteria, and has a very fetid odor. Gangrene of the lung follows, with extensive disintegration. Death in such cases is due to exhaustion. In the abdominal type the disease is likely to be mistaken for typhoid fever. The symptoms are great weakness, loss of appetite, coated and dry tongue, diarrhœa of a fetid character, and delirium. Rigors are infrequent.

The meningeal type is noted less frequently than the pulmonary or abdominal, and when present is usually associated with one of the other two forms. Cephalalgia and vomiting are marked symptoms. The meningitis is due to infective thrombosis, or may occur as a complication. Other symptoms are high temperature; agitation; considerable excitement; tonic and clonic contractions of the face, neck, and upper extremities, or paresis; squint; retraction of the head and neck; delirium in the later stages. Rigors are apt to develop

[1] Die Otitischen Erkränkungen des Hirns, etc., 1902.

when purulent leptomeningitis occurs as a complication. When the leptomeningitis extends to the spine there are, in addition, shooting pains in the spinal region, great prostration, and a sensation of constriction around the waist or girdle pains. The patient gradually passes into a comatose condition, although the mind is clear in the early stages.

Diagnosis. The difficulties in making a diagnosis of sinus thrombosis frequently arise from the complications that are likely to develop, such as meningitis or cerebral abscess, or both. In a simple, uncomplicated case, in which a patient with purulent otitis media has chills, high temperature with remissions, a rapid pulse, vomiting, and in which there is also more or less œdema about the mastoid tip, extending upward and over the exit of the mastoid vein and in the upper third of the posterior cervical triangle, a diagnosis of sinus thrombosis is comparatively certain. The surgeon should not wait for symptoms denoting disintegration of the thrombus and general systemic infection, as the success of the operation depends almost wholly on its early performance. Still, cases do recover in which the operation has been deferred until a late stage, but such instances do not offer much hope.

The Time to Operate. This is a question that has been much discussed by surgeons, and although opinions differ, the sentiment of the majority is certainly in favor of an immediate operation in case of sinus thrombosis as soon as the surgeon is sure of the diagnosis. Except in rare instances, unless an operation is undertaken to remove a thrombus, a fatal result must be expected. The earlier the operation is performed, the better the chances of success.

Another point in the operation about which different views have been entertained is the advisability of ligating the internal jugular vein. Those opposed to ligation contend that many cases have recovered without tying the vein; that infarctions may be carried into the lungs by the other internal jugular vein; and that, as it is extremely difficult in many cases to find the vein, owing to infiltration of the tissues, the procedure adds greatly to the gravity of the operation. While it is undoubtedly true that minute infective particles may be swept into the circulation in spite of ligation of the internal jugular vein, the general opinion is in favor of tying the vein in all cases in which the sinus contains putrid material, pus, or a disintegrated thrombus, as well as in those cases in which the presence of a thrombus in the internal jugular is evident from the cord-like induration felt along the border of the sterno-mastoid muscle. A thrombus may form in the sigmoid sinus in an incredibly short space of time. I have seen a case in which thrombosis occurred as a result of acute otitis media of only ten days' duration, due to the presence of streptococci. I have also had under my care a boy of eight who developed an acute otitis media as a result of tonsillitis. When called to see him a free incision was made in the drumhead. There were symptoms of mastoid disease at that time. At first there was an improvement, but on the third day the temperature gradually rose to 104.2° F. (rectal), pulse 140. At this time the boy did not complain of pain, but was stupid and inclined to sleep. Under ether an incision was then made behind the auricle, the mastoid cells were exposed from the antrum to the tip, and softened bone with granulations

removed. Very little pus was found. The bone was softened, especially in the direction of the jugular bulb, and there was considerable œdema about the mastoid tip and in the direction of the upper third of the posterior cervical triangle. On exposing the sigmoid sinus it appeared to be normal. Temporary improvement followed, but the temperature soon rose again to 104.2° F. (rectal), and two days later, just one week after the beginning of the attack, the boy had a chill lasting about twenty minutes, followed by a temperature of 103.4°, and later 104.4° F. The patient was very drowsy and stupid, and had a rapid pulse. It was decided best to open the sigmoid sinus at once. The sigmoid groove was cut away so that the sinus was exposed for at least an inch and a half. A soft clot was adherent to the walls of the sinus near the bulb, and was removed with a curette. The circulation was re-established at both ends of the sinus, the latter plugged with strips of iodoform gauze, and the wound bandaged. The boy recovered. The case is mentioned to show that, even in patients suffering from acute otitis media, thrombosis of the sigmoid sinus may occur in an incredibly short space of time, and that one should be constantly on the alert for such a complication, especially when the temperature remains high after incision of the drumhead and the presence of streptococci has been demonstrated.

Another case[1] which recently came under observation was a child of four years, who had symptoms of sinus thrombosis, with a temperature of 105° F., due to a mastoid inflammation of both sides. In this patient the infection was caused by pneumococci. The boy had

[1] Transactions of the American Otological Society, 1898.

an attack of earache first on one side and later on the other. A free incision was made in each drumhead, and for a short time improvement followed. Both mastoid processes were involved, and it became necessary to remove softened bone and granulations. The temperature immediately fell, and recovery followed. At each operation the sigmoid sinus was exposed, so that, if necessary, a thrombus could be removed if the patient had further symptoms characteristic of such a complication.

A case seen in consultation with Dr. Gruening had been operated on for double mastoiditis following the grippe—he did well for two weeks, when suddenly he had a temperature of 104° F. following a chill which fell to below normal, so that within twenty-four hours there was a difference of seven degrees in his temperature. The patient had had chills during the previous summer, and an examination of the blood showed that the rise in temperature was due to malaria which was cured by the administration of quinine. Sinus thrombosis was naturally suspected.

Prognosis. If the diagnosis is made at an early stage, and the operation undertaken at once, the chances of a successful result are fairly good, provided the patient's general condition is satisfactory. When septic poisoning has developed, although there is little hope of a favorable outcome, one should nevertheless remove the source of infection at once.

Method of Operation. The first step in the operation consists in making the usual incision behind the auricle for exposing the mastoid cells. The antrum should first be opened freely and its contents thoroughly removed. The tympanic roof should be explored for

a carious opening. To uncover the sinus, the surgeon should make a second incision through the soft parts at a right angle to the incision already made for the mastoid operation (see Fig. 108, from the original mastoid incision to point 3), and this should be carried backward so that the flaps can be dissected away from the skull. After having removed all mastoid cells the sigmoid groove can generally be seen in position. The bridge of bone between the sinus and the antrum should be removed carefully. The sinus in some instances lies very close to the antrum, so that if the latter is opened first and the bone chiselled away layer by layer in a direction directly backward, the anterior knee of the sigmoid sinus will come into view, or if the bone is softened just over the sigmoid groove in the middle portion of the mastoid process, the vessel may be laid bare at that point. (Fig. 130.)

The sinus should be exposed to view for at least one inch, in order to give the surgeon an opportunity to determine the existence of a thrombus, and also to enable him to control the hemorrhage, if necessary. The condition of the sinus wall should be noted, whether thickened or covered with granulations. If the latter are present, they should be carefully scraped away, so that the sinus walls may be thoroughly inspected, while with a probe one should examine to see if a sinus leads to a cerebellar abscess or not. If a thrombus is present, the sinus will appear unusually prominent, and, by palpation, it will seem firm and dense. The sinus may or may not pulsate. Formerly it was thought that if pulsation was noted there could be no thrombus, but such a view is erroneous. If an exploring needle is introduced, its point should be

directed both upward toward the torcular and also downward toward the bulb. The needle may show the presence of serum, pus, a disintegrated clot, or even blood. The withdrawal of blood does not necessarily indicate the non-existence of a thrombus, because the latter may be very soft (parietal), and there may be a stream of blood running through the centre of the vessel.

If, after exposing the sinus, the surgeon is still in doubt as to the presence of a thrombus, it is much wiser in many instances, provided the patient's condition is satisfactory, to postpone further operative interference for at least twenty-four hours, because by inserting a hypodermic needle a perfectly healthy sinus may become infected. The only certain method of determining as to the occurrence of thrombosis is to make *a free incision into the sinus wall.*

If the surgeon determines that there is pus or a disintegrated clot in the sinus, or the thrombus has extended into the upper third of the internal jugular vein, as shown by the presence of a cord-like feeling along the sterno-cleido-mastoid muscle, he should, before incising the sinus, make an incision in the integument along the anterior border of the sterno-mastoid muscle, extending from the mastoid tip to the upper border of the clavicle. The internal jugular vein should then be exposed carefully, and after tying both the facial and superior thyroid veins, two ligatures should be applied to the internal jugular vein, one just above the clavicle and the other as near the jugular bulb as possible. The portion of the vein between these two ligatures should then be excised. The greater portion of

this wound should be closed by sutures, an opening being left for drainage.

If it has been decided not to ligate the internal jugular vein, then an incision should be made in the

FIG. 128.



Ligation of the internal jugular vein (Gerber). *V. f.* Facial vein. *C.* Carotid Artery. *J.* Internal jugular vein. *A. ste.* Sterno-cleido-mastoid artery. *M. ste.* Sterus-cleido-mastoid muscle. *N. v.* Vagus nerve.

long axis of the sinus and all obstructions removed by means of sharp curettes. One should introduce the curette first toward the torcular and re-establish the current

in this direction. By means of a gauze tampon applied directly at this point, and held in position by means of the finger, the walls that have been incised being folded inward, the hemorrhage can be easily controlled. The curette should then be passed downward toward the bulb, and the current re-established as in the upper portion of the sinus.

If the blood cannot be made to flow, and the surgeon has reason to believe that the clot has extended into the internal jugular vein, he should at once tie the vein, provided the patient's general condition is such as to warrant further operative interference. Reference has already been made to two cases of sinus thrombosis, in which the internal jugular vein was not ligated, owing to the condition of the patients. On the day following the operation there was marked improvement in each instance, and in consequence the sinus was not ligated. If, however, the flow has been re-established, a gauze tampon should be applied at once to control the hemorrhage. The wound should be dusted over with iodoform and boric acid, 1 : 4, covered with several layers of gauze, and then a bandage should be applied. The same aseptic precautions should be carried out as in operations for mastoid disease or cerebral abscess. The dressings should not be changed for several days if the patient's general condition shows a decided improvement.

Primary Phlebitis of the Jugular Bulb. Considerable attention has been paid of late to a phlebitis and thrombosis occurring primarily in the jugular bulb and especially in the case of children. It is easy to understand how readily the infection may pass through the floor of the tympanum directly to the jugular bulb which lies

beneath, particularly if the plate of bone is very thin. The infection is carried by direct absorption, or by means of the small veins and lymphatics. The diagnosis is made usually on account of the following symptoms: a child, for example, has an acute otitis media following, as a rule, streptococcus infection. After the drum membrane has been incised, the temperature, instead of declining as it should under ordinary conditions, suddenly rises to 104° or 105° F., only to fall to normal or below normal. The temperature soon rises again as before, and the child may at times complain of chilly sensations or may only have cold hands or feet before the onset of fever. The pulse is usually rapid. The child generally feels bright and happy when the temperature is low, but when the fever is high the little patient is apt to be dull and listless. According to Jansen papillitis is of common occurrence in cases of primary phlebitis of the jugular bulb. McKernon[1] has reported six cases of primary jugular bulb thrombosis in children with five recoveries. The ages varied from six months to two years, and in each instance the mastoid bone was found normal. The thrombus in each case was removed from above by opening the sigmoid sinus. This is the usual procedure with or without ligation or excision of the internal jugular vein according to the conditions found. Grunert[2] has written a monograph in which he gives the histories of twelve personally observed cases from the Schwartze clinic and five of other authors where the jugular bulb was exposed and opened. The operation, according to this writer, is

[1] Transactions of the American Otological Society, 1905.

[2] The Operative Eradication of the Bulb of the Jugular Vein, Archives of Otology, April, 1905.

indicated when, with continuation of severe pyæmic symptoms, the disintegrated thrombus of the jugular bulb cannot be satisfactorily treated by manipulation from the sinus above or the vein below or by irrigation, and in certain forms of primary bulb thrombosis. The operative purpose is to convert the sinus, bulb and jugular vein into a continuous groove with a common external opening.

After an operation on the brain the patient is likely to suffer considerably from shock, so that the pulse becomes weak and rapid. Beside the use of strychnine, digitalis, nitroglycerin, the inhalation of oxygen, and the administration of other remedies, the injection into the rectum of the *normal saline solution* is particularly to be recommended in just such conditions. In extreme cases intravenous injections of the saline solution are necessary.

The Intravenous Injection of the Normal Saline Solution. An excellent solution for intravenous injection is the following one, which is recommended by Hare :

Calcium chloride	0.25
Potassium chloride	0.1
Sodium chloride	9.0
Sterilized water	1000 c. c.

The vein into which the salt solution usually is injected is the median cephalic vein. The patient's arm is uncovered and the skin scrubbed with soap and water, and then washed with a solution of bichloride of mercury (1 : 3000). After a bandage has been applied tightly around the upper arm so as to bring into prominence the median cephalic vein, the skin is dissected away from the vein, and the latter laid bare for the

space of an inch or an inch and a half. The surgeon should apply a small bull-dog clip to the proximal end of the exposed vein, in order to keep the lower part of the vein distended with blood. A ligature should then be applied to the distal extremity of the exposed portion and tied. A second ligature should be passed under the vein, but not tied. The surgeon should then divide with a pair of scissors the distended vein between the clip and the ligature which has been tied, and should insert a glass canula into the opening in the vein. The ligature around the vein which was not tied should then be bound around the canula to hold it in position. It is well, before inserting the canula, to immerse it in the saline solution in order to prevent the blood from coagulating on its tip. A piece of rubber tubing, three or four inches long is attached to the exposed end of the canula. By means of a pipette the rubber tubing and glass canula are filled with the saline solution until all the air has been displaced. In order to prevent the introduction of air into the vein, the surgeon should allow the solution to flow first from the irrigator through the rubber tubing connected with it. Then a glass tip, attached to this rubber tubing, is inserted into the other tube. The solution should be allowed to enter the vein very slowly, and the irrigator should not be held higher than three feet above the patient's arm. The amount injected is usually about a quart. A few moments after the solution enters the vein the patient's pulse usually becomes slower and has more volume, while the respirations are less hurried. The fever, if present, is apt to be less. Shortly afterwards, however, the patient may have a severe chill, with a full and rapid pulse. An hour or two later there should be

marked improvement in the patient's general condition. Great care should be taken to have the irrigator warm, as well as the saline solution. The temperature of the latter should be 101° F.

In many instances the life of a patient is saved by the timely injection of the normal saline solution, particularly after the operation for sinus thrombosis.

LEPTOMENINGITIS.

There are two varieties of leptomeningitis, the *serous* and the *purulent*, the latter being observed more frequently as a complication of ear disease than the former. Leptomeningitis is a disease which, starting in the arachnoid, may spread very rapidly until the entire subdural and intraventricular spaces are involved, as well as the pia, and at times the superficial portion of the brain. The cerebro-spinal fluid becomes increased in quantity, and the sheaths of the spinal and cerebral nerves often are implicated. This form of inflammation is more serious and fatal than either thrombosis of the sigmoid sinus or cerebral abscess, and is in consequence much more dreaded by the surgeon.

In serous leptomeningitis there is at first a hyperæmia, which is followed by exudation of serum, in which often are found flakes of fibrin and at times a few leucocytes. There may be such a rapid accumulation of serum as to give rise to serious pressure-symptoms, the ventricles becoming greatly distended. At times acute serous leptomeningitis is localized. If a disk of bone is removed from the skull in a patient suffering from this disease, the arteries and veins will be found large and full, and the pia swollen and infected, while clear fluid will be contained within the arachnoid mesh-

work. After making a trephine opening and reflecting the dura, there may be so much fluid that the meshes of the arachnoid protrude and form a hernia of the membranes, which are filled with fluid.

Purulent Leptomeningitis. In this disease, which may be due to embolism, there is at first hyperæmia with serous exudation, followed by marked dilatation of the veins. The latter at times contain granular coagula. The perivascular spaces also are filled with an exudation of leucocytes. At first the exudation is limited to the arachnoid, as well as the spaces and clefts in the pia mater; but it is liable to extend to the underlying brain substance, which becomes soft and moist, while the ventricles are filled with sero-pus. When the base of the brain becomes involved, the disease is likely to affect the spinal cord.

Symptoms. Headache is usually the first symptom, which may be localized at first, but which eventually extends over the entire head. Other symptoms are restlessness, insomnia, photophobia, hyperæsthesia of the cutaneous nerves, pupils contracted and responding but little to light, abdomen retracted, pulse full and bounding and accelerated, high temperature, and optic neuritis. In the later stages the patient is dull and stupid, and convulsions, at first confined to the muscles of the face and extremities, become general. The pupils are dilated and often unequal, while the pulse becomes slower, owing to the cerebral pressure. In the last stages paralyses occur, the patient passes feces and urine involuntarily, the respirations are increased in frequency, the pulse is very rapid, the patient becomes comatose, with symptoms of general paralysis, and death follows.

The course of this disease may be extremely rapid, or death may not occur for weeks.

Diagnosis. Symptoms suggestive of meningitis are observed occasionally, more especially in children, which often are due to retention of pus in the middle ear or external auditory canal, so that a diagnosis of this disease is extremely difficult in the early stage. It is only when paralyses occur, and there are optic neuritis and other characteristic symptoms, that the surgeon can be sure of the diagnosis. This disease is associated occasionally with thrombosis of the sigmoid sinus (see Fig. 129) or cerebral abscess, or with both, so that when these complications occur a diagnosis is extremely difficult.

Lumbar Puncture. Quincke in 1891 was the first to make use of lumbar puncture in a number of cases of hydrocephalus for the purpose of diminishing the intracranial pressure. The procedure consists in inserting a hollow needle into the subarachnoid space and drawing off some of the cerebro-spinal fluid. It is of great value, according to some authorities, in making a diagnosis of meningitis caused by suppurative otitis media. After lumbar puncture has been made, one may feel reasonably sure of a diagnosis if pus can be demonstrated in the cerebro-spinal fluid by microscopic and clinical examinations. According to Henle[1] "the spaces between the 3rd and 4th, or 4th and 5th, lumbar vertebræ are best adapted for puncture," although some surgeons insert the needle between the last lumbar vertebra and the sacrum. If possible the patient should sit upright and bend forward. The skin should be scrubbed first with green soap and water, and washed with ether. Then a very small incision should be made in the skin. Quincke's

[1] Von Bergmann's System of Practical Surgery.

method is to introduce the needle a few millimetres to one side of the median line, while Henle enters between the spinous processes and directs the needle upward and inward. The latter uses a canula about 0.6 to 1.2 mm. in diameter and about 5 cm. long (2 inches). There is an advantage in the hollow needle over the trocar in that one can determine immediately whether the dura has been punctured or not. In adults it is safe to introduce the needle to a depth of about 4-6 cm. ($1\frac{1}{2}$-$2\frac{3}{8}$ inches), and in children to a depth of 2-3 cm. ($\frac{3}{4}$-$1\frac{1}{4}$ inches), according to the thickness of the fat and muscles beneath the skin. In order to determine the pressure, a small glass tube should be attached to the needle by means of rubber tubing and then held in a vertical position. In a normal case the pressure varies between 40 and 125 mm. of water. For purposes of diagnosis a few cubic centimetres will suffice, while in cases of meningitis for therapeutic reasons it may be necessary to draw off as much as 100 cc. although it is better to remove much smaller quantities at each time. The fluid should be drawn off very slowly as otherwise the patient may complain of weakness. In infants the fontanelle will be depressed if much fluid is removed so that the surgeon will have that point as a guide in such cases.

Prognosis. Although this disease is generally fatal, the prognosis is not necessarily hopeless, for cases have recovered, especially after operation.

Treatment. As infective meningitis occurring in this connection is usually due to an extension of the disease from the middle ear or mastoid cells to the meninges, it is important first of all to open the antrum and explore the tympanic roof when there are symptoms of

infective meningitis of the middle cranial fossa, and to remove all purulent secretion between the bone and the dura or pia mater. If the symptoms point to involve-



FIG. 129.

Temperature-chart of a fatal case of chronic suppurative otitis media followed by sinus thrombosis and suppurative leptomeningitis.

ment of the base of the brain and spinal cord, the thin plate of bone covering the sigmoid sinus should be cut

away and the posterior cranial fossa investigated. In cases of serous meningitis, after exposing the dura the surgeon should make an incision in it so as to cut into the subarachnoid space, in order to drain as thoroughly as possible the copious serous fluid and thus relieve the pressure symptoms. Lumbar puncture should also be done.

Some difference of opinion exists among surgeons as to the propriety of opening the dura in cases of purulent meningitis. Some are opposed to any operation while others feel that the only hope of saving the patient lies in making a free incision through the dura and in resorting to lumbar puncture.

If a diffuse suppurative meningitis exists, the patient is almost sure to die so that we are justified in operating in such cases, for in a number of instances recovery has followed operative interference. Macewen has reported several cases of purulent lepto-meningitis followed by recovery, in which he found the tympanic roof eroded, the dura congested and covered with pus. A free incision was made in the inflamed membranes, which permitted the purulent intradural exudation to escape. In these cases the area of the infective process was undoubtedly limited, for when there is general implication of the membranes the case is usually, but not always fatal.

It is always well in doubtful cases in which the symptoms are grave and point to intracranial involvement, to make a thorough investigation of the antrum and tympanic roof, as well as of the sigmoid sinus. Then, if there are well-marked clinical symptoms and we find bacteria in the fluid drawn off by lumbar puncture, the surgeon should remove thoroughly all diseased

bone in the mastoid cells and expose the dura. A crucial incision of the size of a half-dollar should then be made in the latter so as to penetrate beneath the subarachnoid space. Sometimes it is necessary to tap the lateral ventricle. In this, as in all other operations, the most thorough antiseptic precautions should be followed, and it is wise to do but little syringing for fear of spreading the infection. The normal salt solution should be used. Lumbar puncture should be practised systematically and repeated as a means of diagnosis and treatment. About 15 cubic centimetres should be withdrawn at a time.

B. A. Randall has reported good results in cases of commencing meningitis by washing out the subdural space with an antiseptic solution.

SUPPURATIVE INFLAMMATION OF THE LABYRINTH.

According to Jansen sixty-two per cent., and according to Whitehead thirty-six per cent. of cases of labyrinth suppuration succumb to meningitis. For some years operations have been performed in cases of suppuration and necrosis of the labyrinth. Jansen was the first to open the vestibule and semicircular canals, followed by Stacke, Kœrner, Lannois, Lermoyez, Ballance and others, and according to these authors operations on the labyrinth, although dangerous, are justifiable. In extensive necrosis of the labyrinth there is apt to be paralysis of the facial nerve. Dundas Grant[1] says: "after the mastoid operation, the persistence or the reappearance of cephalalgia, fever, vertigo, vomiting and nystagmus should

[1] Annales des mal. de l'oreille, Oct. 1904.

make us consider the possibility of suppuration of the labyrinth, symptoms similar to cerebellar abscess but much more frequently they are labyrinthine." He further believes that in addition to the radical operation, cases of serous meningitis should be treated by lumbar puncture and opening of the labyrinth, and in

FIG. 130.



Removal of mastoid cells showing sinus exposed (Gerber). *B.* Horizontal semi-circular canal wall. *Si.* Sigmoid sinus. *Pt.* Promontory. *E.* Point sometimes selected for exploring the cerebellum for abscess.

cases of cerebellar abscess without sinus phlebitis, the labyrinth should be opened and carefully inspected. Schmiegelow [1] has reported a case of chronic otorrhœa of the right side in which there was a cholesteatoma of the temporal bone, followed by necrosis of the labyrinth.

[1] Monatsch. f. Ohrenheil. July, 1904.

Healing was brought about by complete chiseling and resection of the labyrinth. The lateral semicircular canal had been opened through disease. After removal of a sequestrum the cerebro-spinal fluid escaped. The whole under surface of the temporal bone was separated and the cochlea was found carious. Very little of the temporal bone was left and the flow of cerebro-spinal fluid lasted for eight days. The wound eventually healed through Thiersch grafts.

FIG. 131.



Opening of the labyrinth. Posterior and horizontal semicircular canals opened (Gerber). *S.* Sound introduced into the horizontal canal and brought out in the foramen ovale. *Ve.* Posterior semicircular canal. *Co.* Opening into cochlea after removal of the promontory.

Ballance[1] has removed the semicircular canal in a suppurative case, in which the principal symptom was extreme vertigo, with deafness and persistent noises so that the patient could not cross the room. After the

[1] Rev. Heb. de Laryngal., 8 Avril, 1905.

operation the noises and vertigo disappeared, and the hearing was slightly improved.

The route of the infection in suppuration in the labyrinth is usually through the foramen ovale or foramen rotundum, or through an artificial opening. In a suspected case one should look for a carious opening into the horizontal semicircular canal or into the promontory.

S. von Stein [1] has observed four hundred and twenty trepanations with ten cases of necrosis and suppuration of the labyrinth. Of this number one was fatal, one was not improved, but the eight others were cured—that is the suppuration ceased after removal of the necrosed bone in the labyrinth and escape of pus from the labyrinthine cavity but without restoration of the function of hearing. The operation had no favorable effect on the coördination of movement. When operating the surgeon should open first the horizontal semicircular canal as the disease generally starts there. The stapes then should be removed and the foramen ovale enlarged so as to get a good opening into the vestibule, if there is suppuration in the labyrinth. After opening the vestibule, if the surgeon suspects more or less involvement of the cochlea, he should then open the latter through the promontory, being careful not to injure the facial nerve or the carotid artery. After opening the semicircular canal the surgeon should note as to whether the membranous canal is intact and should avoid injuring the latter for fear of setting up a purulent endolabyrinthitis. Von Stein introduces a small syringe, in cases of perilabyrinthitis, into the cavities of the labyrinth and injects peroxide of hydrogen, afterward drying with cotton and filling the cavities with iodoform. The

[1] Annales des Mal. de l'oreille., Jan., 1905.

author would advise against the use of peroxide in such cases for fear of disseminating the infection.

Neumann[1] operates in a different way by liberating first the dura of the posterior cerebral fossa to obtain a good view of the pyramid. Then he separates completely the dura from the posterior covering or wall of the pyramid. After having resected the posterior wall of the pyramid, he uncovers the vestibule and semicircular canals. The advantages of this method, according to Neumann, are that one obtains a good view of the entire pyramid while the approaches or communications of the labyrinth toward the cranial cavity are easily seen and made use of for drainage purposes.

Brieger[2] believes that lumbar puncture should not be neglected in cases of suppuration of the labyrinth. He further says "that it is not unusual for the operation to cause a labyrinth suppuration to flare up and thus develop a meningitis.

In the light of our present knowledge it would seem wise to confine the operation for the most part to those cases in which, together with clinical labyrinthine symptoms, a carious spot is found in the region of the promontory or horizontal semicircular canal, since the opening up of the labyrinth is always a dangerous procedure.

PYÆMIA.

Septicæmia and pyæmia are complications that are liable to occur in connection with suppurative otitis media whenever septic material is absorbed, whether in

[1] Annales des Mal. de l'oreille., Mars, 1905.

[2] Transactions of the Seventh International Otological Congress, 1904.

disease of the middle ear, antrum, or mastoid cells, or in thrombosis of one of the sinuses or of the internal jugular vein. In cases of perforation of the mastoid tip, in which the pus makes its way beneath the deep fascia, unless evacuated at once there is great danger that absorption will take place and give rise to blood-poisoning. If septic particles are carried into the circulation from a disorganized clot, infarctions occur in the lungs as well as other organs, and fatal pyæmia is usually the result. This fact demonstrates the importance of always removing every source of infection whenever the mastoid antrum or cells are involved. Some writers, notably Koerner, believe that metastatic pyæmia may develop from osteophlebitis alone without involvement of the sinus. On the other hand, Jansen[1] states that in sixty-three cases of pyæmia he failed only four times to prove the presence of sinus thrombosis. He further believes that in all cases of pyæmia in which the sinus appears healthy, particularly when following an acute otitis, one should think of a phlebitis of the bulb. Some authorities advocate the internal administration of creosote and bichloride of mercury, but such treatment is usually of little avail. Pyæmia is not always fatal, for in a case[2] of acute purulent otitis media following mastoid disease which came under my care, not only were the lungs affected, but collections of pus were also found in different parts of the body. The case was that of a man, aged twenty-three years, who had an acute attack of earache followed by a profuse discharge. He developed mastoid disease, but disappeared from obser-

[1] Archives of Otology, 1901, vol. xxx. Nos. 4 and 5.

[2] Bacon: Transactions of the American Otological Society, 1885, vol. iii. part iv.

vation for several days. He returned, and complained of nausea, vertigo, and chilly sensations. The mastoid cells were opened immediately, and found filled with pus. He subsequently had chills, high temperature with sudden remissions, sweating, and became much emaciated. He eventually recovered. The case was one in which the pus, after breaking through the mastoid tip (Bezold perforation), burrowed beneath the deep fascia in the neck.

CHAPTER XIII.

DISEASES OF THE SOUND-PERCEIVING APPARATUS.

Diseases of the sound-perceiving apparatus are classified usually as primary or secondary affections, the former being met with much less frequently than the latter. According to the earlier writers, many cases were called "nerve deafness" which should have been classified as affections of the middle ear, but in the light of our present information there are comparatively few cases of primary disease of the internal ear. Affections of the internal ear may develop idiopathically or be due to severe cold, blows, falls, sunstroke, or to malarial fever, typhus and typhoid, scarlatina, syphilis, measles, diphtheria, influenza, leucocythæmia, Bright's disease, diabetes, mumps, and locomotor ataxia. Chemical changes in the blood of patients suffering from general diseases produce disturbances in the auditory centres or in the expansion of the auditory nerve in the labyrinth. Secondary changes take place in the labyrinth from inflammation of the middle ear, but it should not be forgotten that the same cause may produce affections of the middle ear as well as of the labyrinth and the auditory nerve, or that the sound-perceiving and the sound-conducting apparatus may simultaneously be involved. This is especially the case in those insidious forms of adhesive catarrh which lead to anchylosis of the stapes. In cases of acute inflammation of the middle ear, owing

to the anastomosis between the vessels of the tympanum and labyrinth, the latter may become congested, and, as a result, ecchymoses and serous infiltration of the membranous labyrinth may follow, with increase of the intralabyrinthine pressure. When this occurs, the patient may complain of deafness, giddiness, vertigo, nausea, vomiting, tinnitus; and the ability to hear the tuning-fork by bone-conduction may be impaired or even lost. In some cases there is hyperæsthesia of the auditory nerve. In diphtheritic affections the labyrinth may become involved from the entrance of micro-organisms, and in chronic purulent otitis media followed by caries and necrosis the internal ear is liable to be involved. Pathogenic micro-organisms, the streptococcus, the staphylococcus, and the diplococcus of pneumonia, may invade the labyrinth from the cranial cavity by means of the aquæductus vestibuli and aquæductus cochleæ. In consequence of such invasion there is liable to be a rapid destruction of the connective tissue, or there may be a new growth of connective tissue and bloodvessels leading to bone formation, or the micro-organisms may cause thrombo-arteritis and phlebitis. The auditory nerve is affected more frequently by general disease than the other nerves of special sense; and certain drugs, such as quinine and salicylic acid, as well as the excessive use of tobacco, may give rise to subjective noises, and to deafness if used in large doses or continuously.

In certain diseases of the brain and spinal cord, especially epidemic cerebro-spinal meningitis, and in consequence of cerebral tumors, disturbances of hearing are due to pressure on the internal auditory nerve or to the extension of inflammatory products from the cranial

cavity to the labyrinth and auditory nerve trunk. It is the opinion of many writers that in cases of cerebro-spinal meningitis the poison enters the nose, and that the ears may be the tract through which the microbe travels to the brain. According to G. Royet,[1] the pneumococcus is found frequently in the cephalo-rachidian fluid and less frequently the meningococcus of Weichselbaum and the other bacteria. The cerebro-spinal fluid is always opalescent at first, then purulent and greenish. Increased leucocytosis is well-marked with an increase also of the polynuclear elements. In affections of the brain and medulla the nuclei and roots of the auditory nerve may be involved, and as a result deafness follows. Disturbances of hearing are observed at times in diseases of the liver, kidney, uterus, stomach, and also in anæmia and marasmus. Other causes of internal-ear disease are aneurism of proximate blood-vessels and malignant growths beginning in the temporal bone. Fracture at the base of the skull may produce temporary or permanent deafness, with or without hemorrhage from the ear, followed by a serous discharge. Artillerymen and boilermakers are especially liable to nerve deafness, on account of the loud sounds to which they are constantly exposed.

Progressive disease affecting the auditory nerve trunk is seldom confined permanently to one ear, for sooner or later the other ear is likely to be involved. As the root-fibres of the auditory nerve on one side extend into the external auditory nucleus on the other, degenerative processes of one internal auditory nerve may attack the centre and stem of the nerve on the opposite side.

In some affections of the internal ear the deafness

[1] Meningite cérébro-spinale, d'origine otitique, Thèse de Paris, 1905.

comes on very suddenly. Especially is this the case in cerebro-spinal meningitis, syphilis, Ménière's disease, as well as in consequence of falls and blows. In addition to giddiness, deafness, vomiting, and loss of hearing by bone-conduction, the patient often complains of disturbances of equilibration, which are at times so marked, as in Ménière's disease, that the patient is unable to stand without support. Beside the causes already mentioned, it is undoubtedly a fact that hereditary influences play an important rôle, and also that a sudden mental disturbance, such as fright, may give rise to an affection of the internal ear through a sudden disturbance of the circulation of the labyrinth.

Children are more frequently the subjects of disease of the internal ear than adults, for the reasons probably (1) that they are more prone to the exanthematous diseases and cerebro-spinal meningitis, and (2) in children the communications between the cranial cavity and the middle and inner ears, and also between the tympanum and labyrinth, are more direct than in adults, so that inflammatory products are more liable to be carried from the cranial cavity and the middle ear to the labyrinth. Very young children, or the aged, are more frequently the subjects of labyrinthine disease than those of middle age.

In diseases of the auditory nerve the hearing is not so subject to marked fluctuations as in middle-ear disease. If the deafness due to a nerve lesion does not become more marked, it is likely to remain about the same from day to day, although impaired health, bodily fatigue, and worry, as well as over-indulgence in the use of tobacco and alcohol, tend to aggravate the disease.

Prognosis. The prognosis is, as a rule, unfavorable,

although very much depends on the causation, the intensity, and the duration of the disease. In recent syphilitic disease much can be accomplished by general constitutional treatment and by the hypodermic injection of pilocarpine, and undoubtedly in many cases in which there is serous and possibly small-celled infiltration, permanent restoration of hearing may follow. On the other hand, it is proved by clinical observation that deafness often remains after extensive effusion of blood has taken place in the labyrinth, with the invasion of the various cocci, and especially where there is extensive caries or necrosis of the temporal bone. In order to make a differential diagnosis between middle-ear and internal-ear affections, it may be stated, as a general rule, that—

1. In diseases of the inner ear, low-toned tuning-forks are heard better than the higher ones by aërial conduction, if any fork is heard at all; while in disease of the middle ear, if the nerve is not affected, the reverse is true—*i. e.*, the higher forks are heard better.

2. In disease affecting the labyrinth or internal auditory nerve, the bone-conduction becomes impaired or lost, as shown by Rinné's, Weber's, or Schwabach's test (see chapter on Examination of the Ear). In the case of a person with normal hearing, a vibrating tuning-fork should be placed on the mastoid process, and the surgeon should be notified by a signal from the patient when he ceases to hear the sound. If the fork is then held close to the meatus, it will be heard for several seconds longer. In middle-ear or external-ear diseases the tuning-fork will be heard much longer when placed on the mastoid than when held close to the meatus. In a case of deafness in which the labyrinth is involved,

the bone-conduction becomes much impaired or lost altogether. Notes of a low pitch are, as a rule, heard better than high ones. In consequence, the watch is not heard as well as the voice.

Other important instruments for testing the hearing are the acoumeter, the Galton whistle, and König's rods. When the acoumeter is not heard by bone-conduction, there is a reasonable certainty that the internal ear is affected. It should not be forgotten, however, that frequently diminished bone-conduction is observed in cases of adhesive catarrh of the middle ear as a result of secondary implication of the internal ear or of intra-labyrinthine pressure, and to physiological conditions in persons over sixty years of age.

If a patient presents himself suffering from deafness of recent origin, and if an examination shows that the drumhead is normal and the Eustachian tube pervious, it is probable that the case is one of disease of the auditory nerve. It is often difficult for the surgeon to differentiate between a middle-ear affection, in which there are symptoms of vertigo, dizziness, and tinnitus, and an affection of the brain itself. The same symptoms observed in labyrinthine disease may occur in disease of the central nervous system. In the latter there will be such other symptoms as choked disk, paralyses, loss of sensation of one or both sides of the body, and diminished or increased tendon reflexes.

HYPERÆMIA OF THE LABYRINTH.

Hyperæmia of the labyrinth is likely to occur in a severe case of acute purulent otitis media, especially if due to scarlatina, diphtheria, or other acute exanthematous diseases. It is observed also in cases of tumor

of the brain in which there is an obstruction to the escape of blood from the inner ear, and in patients suffering from other general diseases, such as typhoid or typhus fever, cerebro-spinal meningitis, and in cerebral congestion accompanying diseases of the lungs or heart. It follows the administration of certain drugs, such as nitrite of amyl, quinine, and salicylic acid. Hence great caution should be observed in administering this class of drugs to patients suffering from deafness from any cause. Unless the hyperæmia is of a temporary character, there may be pigment exudations in the labyrinth or the membranous labyrinth may become thickened. The symptoms in the acute stage of hyperæmia are vertigo, giddiness, vomiting, inabilty to walk steadily, tinnitus, and a sensation of fulness about the ears. Very often it will be noticed, on examination of the ear, that there is a slight congestion of the vessels along the handle of the malleus and of the bony canal adjoining the drumhead.

It is very difficult to make a differential diagnosis between hyperæmia of the labyrinth and of the brain centre, owing to similarity of the symptoms. The treatment depends altogether on the cause. The patient should be kept quiet and put on a simple and wholesome diet. If occurring in connection with congestion of the brain, cold applications should be applied to the head. Small doses of calomel, to produce watery evacuations, are also useful. When the drumhead is injected simultaneously, the local abstraction of blood from the ante-auricular and mastoid regions is indicated, as well as the application of the Leiter coil.

ANÆMIA OF THE LABYRINTH.

This affection is observed in cases of general anæmia and chlorosis after great loss of blood, or there may be some obstruction to the blood-supply of the labyrinth, due to the pressure of a tumor, an aneurism, or there may be atheromatous changes, etc. In a case of loss of blood, followed by anæmia of the labyrinth, the symptoms are deafness, distressing tinnitus, faintness, vertigo, nausea, and vomiting.

As to treatment, this should be directed to the cause. When the disease is chronic, iron should be given, while a change of air is often advisable.

HEMORRHAGE INTO THE LABYRINTH.

Slight extravasations of blood occur in the labyrinth as a result of disease or of fracture at the base of the skull, or such extravasations may be due to concussion of the skull or to extensive necrosis of the temporal bone. When due to disease, the walls of the vessels may be weakened by atheromatous changes or by intense hyperæmia during the course of scarlet fever, diphtheria, mumps, smallpox, pernicious anæmia, diabetes, and other diseases.

INCREASED PRESSURE OF THE LABYRINTHINE FLUID FROM DISEASE OF THE TYMPANUM.

As already mentioned, vertigo and disturbances of equilibration are prominent symptoms of labyrinthine disease. In cases, however, of middle-ear disease, or those in which a plug of cerumen presses against the drumhead so that the foot-plate of the stapes is forced inward and the pressure of the labyrinthine fluid is

increased, the patient often complains of more or less vertigo. The membrane of the foramen rotundum should, under normal conditions, by bulging outward compensate for the pressure exerted upon the foot-plate of the stapes. It is possible to explain the occurrence of the vertigo as being due to unusual pressure from the labyrinthine fluid on account of congestion of the blood-vessels or to the fact that the membrane covering the foramen rotundum does not bulge outward, owing to thickening having occurred from catarrhal changes.

A patient came under my care a short time since who complained of severe vertigo. He was unable to leave his room for several days on account of the severity of the attack. The history was that he had had a plug of inspissated cerumen removed from the ear by syringing a few days previously, and that since then he had been troubled with severe dizziness. He was a man of fine appearance and in excellent health. An examination was made of the urine, heart, lungs, etc., as well as of the fundus of the eyes. Nothing was found to explain the vertigo except the condition of the ear, there being slight retraction of the drumhead and catarrhal changes. Inflation by Politzer's method daily for several days effected a cure. In syringing the ear at times the patient will complain of sudden vertigo, and occasionally fall from the chair, owing to faintness. Such a phenomenon is undoubtedly due to sudden pressure exerted on the foot-plate of the stapes.

Aural vertigo is induced occasionally by the presence of a polypus which is attached in the neighborhood of the stapes. When this occurs, the patient often complains of a sense of fulness in the ear, with a distinct subjective circulatory tinnitus. The vertigo may be so

severe that the patient is unable to maintain an erect position. The removal of such a polypoid growth should be performed most carefully, as the necessary manipulation is likely to cause increased vertigo and accompanying nausea.

OTITIS INTERNA, OR INFLAMMATION OF THE LABYRINTH.

Inflammation of the lining membrane of the bony labyrinth, as well as of the membranous labyrinth may be either *primary* or *secondary*, the former being extremely infrequent. The secondary form, as a rule, follows inflammation of the middle ear, and is liable to occur as a result of diphtheria, scarlatina, tuberculosis, or extensive caries and necrosis of the temporal bone. In such cases the labyrinth may be filled with pus, while the walls of the membranous labyrinth become thickened and ecchymosed.

Chronic inflammation of the labyrinth, though extremely infrequent, does occur, and, as a result, there is an increase of connective tissue with hyperostosis and calcareous deposits. The saccule and ampullæ become thickened, and finally there may follow atrophy of the membranous labyrinth and disappearance of the organ of Corti.

It is extremely difficult to make a diagnosis of acute inflammation of the labyrinth, for the symptoms, which generally are noticed in children are similar to those observed in meningitis followed by recovery, except that in the former the disease comes on suddenly, and the more serious symptoms last for about a week. The child is apt to have fever, more or less congestion of the

face, vomiting, delirium, deafness, vertigo, convulsions, and coma.

MÉNIÈRE'S DISEASE.

Ménière's disease is an affection of the semicircular canals, even extending into the vestibule. It was first described by Ménière, of Paris, as being due to an effusion of blood or an acute exudation into the semicircular canals. Many cases of aural vertigo, which are due to the presence of a plug of cerumen in the external meatus or to disease of the middle ear, give rise to symptoms of this disease and are classified erroneously as Ménière's disease. The name of Ménière's disease, however, should be limited to those cases in which there is a sudden effusion of blood into the semicircular canal, followed by deafness, giddiness, distressing tinnitus, vertigo, a staggering gait, nausea, and vomiting. The patient may have premonitory symptoms, such as those already mentioned, or may fall suddenly to the ground from loss of consciousness, which is followed in a short time by excessive pallor of the face, cold perspiration, extreme deafness and tinnitus, with an inclination to vomit and more or less vertigo on trying to walk. The disease is not confined to those who are in poor health, but it may attack strong and healthy individuals who previously had not suffered from ear disease. Occasionally, a patient may have premonitory symptoms, such as slight dizziness, tinnitus, and some deafness, before the more serious symptoms manifest themselves.

The cause of the disease is obscure, some observers attributing it to excessive heat of the sun. The deafness is usually very marked, so that the patient can hear only very loud sounds close to the ear. The watch

an acoumeter cannot be heard by contact. Bone-conduction, as shown by tests with the tuning-fork, is very much impaired or lost altogether. If an examination of the ear is made at the time, the drumhead will usually be found normal, as well as the Eustachian tube. Even after the patient has recovered from the vomiting and the unconsciousness, the giddiness and staggering gait may persist and be slow in disappearing, so that it may be some time before the patient can walk without a cane, and he is apt to incline his body to the affected side in walking. The disturbance of hearing generally remains, although there may be some improvement, while the tinnitus is often most distressing. The milder attacks of this disease are mistaken sometimes for biliousness, but a careful consideration of the symptoms already mentioned should make the differential diagnosis comparatively easy. Ménière's disease is sometimes suggestive of disease of the brain. In the latter, however, there are generally symptoms denoting paralysis of spinal or cerebral nerves, but it should not be forgotten that giddiness, vomiting, nausea, and tinnitus are symptoms characteristic of brain affections. I have referred to a case of chronic purulent otitis media (in the chapter on Intracranial Affections), in which the patient had all the symptoms of a cerebellar abscess—giddiness, vomiting, a staggering gait and choked disk—but who finally recovered after proper drainage of the middle ears had been established by scraping away all granulations from the tympanic cavity. A positive diagnosis can only be made of Ménière's disease if the surgeon knows that the patient, before the attack, had normal hearing, and, further, that an examination of the ear made a short time afterward

shows the Eustachian tube to be previous and the drum-head normal.

Prognosis. In cases of long standing the prognosis as to the deafness is unfavorable, and it is only in recent cases that one can hope for even a little improvement.

Treatment. The treatment should consist in the use of cold compresses to the head and the application of the Leiter coil about the ear. Calomel in small doses should be given to produce slight catharsis. Sinapisms should be applied to the back of the neck and the calves of the legs. The patient should be kept absolutely quiet in bed and given a very light diet. Some observers recommend iodide of potash, while others, particularly Charcot, have found much benefit from the administration of quinine in rather large doses. Bromide of potash, and dilute hydrobromic acid in fifteen-drop doses, are of service in relieving the giddiness.

The best results seem to be attained from the injection of five or six drops of a 2 per cent. solution of pilocarpine through the catheter into the tympanum, or from the hypodermic injection of the same remedy and in the same strength. This treatment should be continued for a period of at least four weeks, and injections should be made hypodermically or the solution should be used through the catheter every other day during this time. The surgeon should be very cautious with this drug, and should follow the directions already given (see Chapter VIII.).

Electricity is of but little service in this disease, and should never be used in the early stage for fear of increasing the tinnitus and giddiness. Especial attention should be paid to the patient's general health, particularly if there is a gouty or rheumatic diathesis.

Tonics are indicated in cases where the general condition demands them.

Babinski[1] has reported eight cases of disease of the labyrinth with Ménierè's symptoms which were treated with benefit by lumbar puncture and the tinnitus was improved. He believes that lumbar puncture has a remarkable influence on aural vertigo, and that it also has a favorable influence on tinnitus and deafness but to a less noticeable extent. It is more efficacious generally in pure labyrinthine lesions, according to Babinski, than in mixed lesions of the ear and more so in cicatricial cases of chronic catarrhal otitis media than in the sclerosed cases. He does not think there is any danger in its employment.

Milligan[2] has tried operative measures in three cases of so-called Ménierè's disease where the patients suffered from vertigo for years, incessant noises and progressive deafness. He first performed the radical operation, followed by exposure of the horizontal semicircular canal to the origin of the superior one. The same was done to the posterior canal. Then the horizontal canal was removed with curettage of the cavity and removal of all the soft parts. In two cases the vertigo disappeared and the patients resumed their occupations. There was complete destruction of hearing and slight improvement in the tinnitus.

Richard Lake[3] has reported a case in which he removed a portion of the cochlea, the external wall of the vestibule and the external semicircular canal for the relief of tinnitus and vertigo. The patient was a girl, aged

[1] Deut. Med. Wochenschr., No. 23, 1903.
[2] Rev. Heb. de Laryngol., April 1, 1905.
[3] Rev. Heb. de Laryngol., August 12, 1905.

twenty-six years, who was troubled with intolerable noises and vertigo with a tendency to fall forward. There was complete deafness. Five days after the operation the patient could, with closed eyes, stand up or turn around, and even pick up an object on the floor.

Before resorting to such radical measures, the surgeon should consider the possibility of some intracranial complication arising from such an operation, and should explain thoroughly to the patient the dangers attending such procedures.

MUMPS.

Although disease of the middle ear may occur as a complication of mumps, as shown by the usual appearances of inflammation of the drum membrane, there are a certain number of cases in which the labyrinth is involved either primarily or secondarily. As a rule, only one ear is implicated when the disease affects the labyrinth. The diagnosis is made from the occurrence of sudden deafness with impairment or loss of bone-conduction, as shown by tests made with tuning-forks. Tinnitus is also a troublesome symptom at times. Much can be accomplished by the use of hypodermic injections of pilocarpine if the case is seen at an early date. As a rule, however, these patients do not come to the aural surgeon for treatment until the deafness has existed for a long time. The application of a blister behind the ear is of benefit in some cases.

SYPHILIS.

Syphilis affecting the labyrinth does not generally develop during the secondary stage when the skin is covered by an eruption or there is a throat affection, but

is more likely to be observed during the third stage or even several months or years after the first inoculation, when no signs of syphilis can be discovered on the body. There may be a middle-ear catarrh or purulent inflammation at the same time. In many cases of syphilitic ear disease great thickening of the drumhead takes place, and fibrous bands are found within the tympanic cavity. In recent cases it is possible that there is a plastic exudation in the labyrinth, but this is not certain. In a later stage the following abnormal appearances have been found, viz., condensation of the periosteum of the vestibule, fixation of the foot-plate of the stapes, and great infiltration of the different portions of the membranous labyrinth. The deafness, as a rule, comes on very suddenly, and is attended with considerable tinnitus. Pain is not usually complained of, but there are frequently more or less vertigo and disturbances of equilibration. An examination may show that the Eustachian tube is involved, particularly if there are throat symptoms. Frequently the drumhead is found very much retracted, opaque, and thickened. The deafness is in most cases profound, and is more likely to affect both ears than one ear alone. The disease being confined mostly to the labyrinth, the watch and acoumeter are frequently not heard at all. While tests with tuning-forks show a very much diminished or even absent bone-conduction, low tones are heard much better by aërial conduction than high ones.

Diagnosis. A diagnosis is difficult in many cases unless there is a syphilitic history or there are evidences of the disease in other parts of the body, particularly the eyes or the teeth. The suddenness of the attack is a suspicious symptom ; but it should not be forgotten that the deafness may come on gradually.

Hereditary syphilis is, I believe, much more frequent in children than is generally supposed. There may be middle-ear disease at the same time that the labyrinth is involved. Other symptoms characteristic of syphilis are opacity of the cornea from interstitial keratitis, and peculiar changes in the teeth, Hutchinson's "pegged teeth," so called. Autopsies made in the case of new-born children who were syphilitic have revealed more or less suppurative otitis media, with thickening of the drumhead and adhesions. There was more or less pus in the vestibule, the organ of Corti was destroyed, while the labyrinth was hyperæmic and contained effusions of blood.

Prognosis. The prognosis is, as a general rule, unfavorable, especially in the case of congenital syphilis and when the disease has existed for quite a period of time. It is only in the recent cases that one can hope for much improvement; and even in such, very often, there is but little change for the better. This does not mean that there should not be carried out a prompt and energetic treatment which should be kept up for some time.

Treatment. The treatment depends very much as to whether there is any manifestation of the disease in other parts of the body. In some cases it is advisable to use mercury in addition to the iodide of potash. In prescribing the former, I generally advise inunctions behind the ear over the mastoid process. The best results will generally be obtained from hypodermic injections of a 2 per cent. pilocarpine solution, gradually increasing the dose from four or six drops until the patient experiences the physiological effect of the drug. Great care should be taken in its administration on account of its depressing action.

BOILERMAKERS' DEAFNESS.

The deafness observed in the case of boilermakers, who are subject to the incessant hammering inside of boilers, is due to an affection of the labyrinth. It has been suggested that there may be some loosening of the articulating joints of the ossicula from the excessive vibrations, but from all the cases that have come under my observation, tests made with tuning-forks have demonstrated the fact that there was impaired bone-conduction, showing that the disturbance of hearing and tinnitus were not due primarily to middle-ear inflammation, although in these patients one frequently finds evidence of catarrhal changes in the membrana tympani.

There is very little that can be done in the way of treatment. If the disease is not far advanced, a change of occupation may occasion considerable improvement in the hearing.

ACTION OF CERTAIN DRUGS.

The use of certain drugs, such as quinine and salicylic acid, continued for a long time and in large doses, undoubtedly causes disturbances of hearing and tinnitus, that may in some cases be permanent. It is supposed by some that the action of quinine upon the ear is similar to the effect produced on the retina, viz., anæmia. The common practice of "aborting a cold" by the administration of large doses of quinine is, in my opinion, an exceedingly bad one, and much injury to the hearing has been brought about by such treatment, especially in those who inherit a tendency to deafness and who are sufferers from more or less middle-ear inflammation.

Wittmaack[1] in his experiments on animals claims to

[1] Zeitschr. f. Ohrenh., Bd. xlvi. Heft 1 u 2.

demonstrate that the only action of quinine is on the nerve cells of the ganglion of Corti and that it does not cause any vascular disturbance in the labyrinth. The action of salicylate of soda is very similar to that of quinine. Haike,[1] following experiments by Wittmaack, has made investigations as to salicylic acid and aspirin. Agreeing with Wittmaack he found anatomic changes in the ganglion cells of the vestibular ganglion as well as in the spinal ganglion, which affected principally the Nissl bodies. In the auditory tract a degeneration of the medullated sheaths could be demonstrated.

Arsenic has an effect on the mucous membrane of the middle ear, for cases of serous otitis media have been reported which did not yield to treatment until the patients had ceased taking arsenic.

Tobacco, according to Delie[2] has a direct action on the acoustic nerve. It attacks both ears at a time, and according to this author it is especially advisable that those suffering from deafness due to sclerosis should abstain as far as possible from its use.

PRIMARY DISEASE OF THE BONY LABYRINTHINE CAPSULE.

Politzer,[3] under this heading, has described a disease which, according to earlier writers, was supposed to be a form of chronic adhesive otitis media, but which, from numerous autopsies, he has proved to be due to a "circumscribed disease of the bony labyrinthine capsule," and, as a result, there is formed a neoplastic bony tissue which gradually takes the place of the normal bone, and, finally, closing the foramen ovale, produces anchy-

[1] Arch. f. Ohrenh. Vol. lxiii, p, 78.

[2] Rev. Heb. de Laryngol, d'Otol., Jan. 28, 1905.

[3] Archives of Otology, 1894, vol. xxiii. p. 255.

losis of the stapes. At times the disease attacks the greater portion of the capsule, so that this same neoplastic tissue involves the foramen rotundum and extends toward the cavity of the labyrinth. Most of the patients observed by Politzer were totally deaf, and in only one case was bone-conduction for a loud-ticking watch retained. The age varied from fifty-five to fifty-eight years, and the appearances of the drum membrane were such as are observed generally in old people, viz., thickening of the drumhead with opacities, and, in some instances, atrophic changes. Beside the deafness, other symptoms noted were persistent tinnitus, which in some cases ceased when the deafness was profound, while in others it persisted; occasional sensations of pain of a tearing nature; dulness and heaviness of the head, and vertigo. The sensations in the head and the vertigo might have been due to the advanced ages of the patients. The disease could not be attributed to any particular cause, although gout, syphilis, and heredity probably played an important rôle.

Politzer believes that this disease exists much more frequently than one is justified in assuming from the cases already reported, and, further, that it may produce serious deafness in young subjects. The treatment is unsatisfactory, but Politzer is of the opinion that if a diagnosis can be made at an early period before the deafness is marked, the disease may be prevented from progressing by the administration of large doses of iodide of potash, which should be kept up for a year or more, with stated intervals, and then the drug should be discontinued. An operation for excision of the stapes should not be considered for a moment.

DEAFNESS DUE TO OLD AGE.

After middle life the ability to hear the notes of a Galton whistle or of the König rods becomes lessened, so that in testing the hearing-power of persons of advancing years one must make allowance for the age of the patient. The otologist is frequently consulted by very old patients who complain of inability to hear conversation, although otherwise they enjoy excellent health. Upon making an examination of the drumhead it will very likely be found thickened, retracted, and opaque, and in some cases atrophic, with evidences of implication of the labyrinth and auditory nerve. Very little encouragement should be given to such patients, for the prognosis is usually unfavorable.

There may, however, be another cause for the impaired hearing, namely, *collapse of the cartilaginous meatus*, so that the anterior and posterior walls touch each other and leave but little space for the entrance of sound-waves. I have improved the hearing of such patients by introducing small silver tubes especially constructed for this purpose.

DISEASES AFFECTING THE AUDITORY NERVE.

Diseases affecting this nerve are, as a rule, due to secondary causes, such as meningitis, cerebro-spinal meningitis, aneurism of the basilar artery, when there is likely to be more or less congestion of the neurilemma of the nerve. In purulent meningitis, and when there are extensive caries and necrosis of the temporal bone, the nerve is likely to be inflamed and infiltrated.

Atrophy of the auditory nerve results from diseases

of the brain and cerebellum, and occurs in certain cases in which tumors of the brain press upon the nerve, the latter disappearing altogether in some instances; also from affections of the spinal cord, especially locomotor ataxia.

Bruhl[1] has demonstrated microscopic specimens of the cochlea of a patient, aged forty-four years, who lost his hearing in the course of tabes. Corti's organ and the termination of the vestibular nerve did not show any marked changes. The spiral ganglion and the trunk of the auditory nerve presented distinct changes, which could be followed into the medulla oblongata. According to the grade of the anatomic changes, it can be assumed that the process of degeneration started from the spiral ganglion.

NEW-GROWTHS AFFECTING THE INNER EAR.

New-growths affecting the labyrinth and auditory nerve generally have their origin in the cranial cavity or the middle ear. Cases have, however, been reported of primary sarcoma and of neuroma of the auditory nerve. The new-growths that are observed most frequently in the external meatus and tympanum which involve the petrous bone and inner ear are epitheliomata and sarcomata.

HYPERÆSTHESIA ACUSTICA.

Occasionally in neurasthenic or highly nervous individuals, or those recovering from a long illness, certain sounds give rise to painful sensations in the ear. This same phenomenon is observed also in certain forms of sclerosis of the middle ear and in acute and chronic diseases of the middle ear and labyrinth. It is a curious fact that such patients are often extremely deaf.

[1] German Otological Society, Archives of Otology, vol. xxxiv., No. 6.

CHAPTER XIV.

DEAF-MUTISM.

DEAF-MUTISM may be congenital or acquired—*e. g.*, due to some disease affecting the ear soon after birth. Heredity is one of the most frequent causes, as well as consanguineous marriages. In many instances of congenital deaf-mutism the cause is traceable to some forefather, although occasionally children are born deaf who present no hereditary history whatever.

Acquired deaf-mutism is usually due to some intracranial or general disease, such as meningitis, hydrocephalus, cerebro-spinal meningitis, injuries of the auditory nerve from falls or blows, scarlatina, measles, typhus and typhoid fevers, diphtheria, mumps, inflammation of the labyrinth, and syphilis. Scarlet fever in children is the most frequent cause of permanent deafness and deaf-mutism. According to the statistics of May, in five thousand six hundred and thirteen cases of deaf-mutism, five hundred and seventy-two were due to scarlet fever. It is occasionally impossible to tell whether a case of deaf-mutism in an infant is congenital or acquired, for until the child is a year old the defect in hearing is not noticed, so that the parents often believe that the child had good hearing up to a certain age, but that the deafness came on suddenly.

Anatomical investigations show that congenital deafness is due to some arrest or defect in development

of the temporal bone, such as bilaterial atresia of the external auditory canal, imperfect development or absence of the tympanic cavity, defects in or bony closure of the fenestra rotunda, anchylosis of the stapes, and arrest of development of the labyrinth or auditory nerve. Alexander[1] of Vienna has reported on the microscopic examination of the ear of a deaf-mute. In addition to other changes there were bilateral atrophy (hypoplasia) of the eighth nerve in the trunk and branches, and of all of the ganglia, atrophy of the static nerve, of the terminals of Corti's organs, and bilateral arrest of development of the capsules of the cochlea.

In acquired deaf-mutism, the principal anatomical changes are constriction of the external auditory canals of both sides occurring in early life, which usually is due to a severe attack of diphtheria or scarlatina in which there has been a purulent inflammation of the middle ear, followed by extensive caries and necrosis of the temporal bone. Other causes are the formation of adhesions with anchylosis of the ossicula and certain changes in the labyrinth and auditory nerve from the general diseases already mentioned. A baby brought to my observation a short time ago, although he had learned to say a few words, had gradually lost all power of speech and hearing. On closely questioning the mother, it was ascertained that the child had had a fall from a baby carriage some months previously. In a case of acquired deaf-mutism reported by Schwabach,[2] following an attack of cerebro-spinal meningitis in the seventh year, there was found complete obliteration of

[1] Archives of Otology, Dec., 1904.
[2] Archives of Otology, Dec., 1904.

the cochlea, with newly-formed bone tissue on both sides and a peculiar new formation in the vestibule, also bilateral. This consisted principally in a convolution of medullated nerve fibres, presenting the picture which Virchow has described as an amputation-neuroma.

Children are, as a rule, apt to be dumb if the deafness comes on during the first three or four years of life, and even later. If the child is bright and intelligent, and has already learned to read, dumbness may be prevented if the parent or guardian shows very watchful attention.

Children who lose their hearing after seven years of age are scarcely ever dumb. In testing the hearing of a child that is deaf and dumb the surgeon should stand behind the child, where he cannot be observed, and then either clap his hands or blow a shrill whistle or trumpet. If the child hears any sound whatever, he will naturally turn his head in the direction from which it comes. By some expression of the face the child will often demonstrate to the surgeon whether he hears low or high tuning-forks when they are placed on the mastoid process and portions of the skull. If it seems certain that some perception of sound exists, the surgeon should test each ear separately to see if the patient hears vowel sounds and speech.

Deaf-mutes without hearing are not affected by rotation. The best-hearing deaf-mutes show generally rotating vertigo.

Prognosis. As to the prognosis, it is more favorable in congenital deaf-mutism than in the acquired form. Occasionally, but very infrequently, improvement occurs in the case of congenital deaf-mutism, while in profound deafness which has followed an attack of scarlet fever,

diphtheria, or intracranial disease, a change for the better is seldom observed.

Treatment. If, in a case of deaf-mutism, the drum-heads are retracted, thickened, and present other changes of a catarrhal character, or the Eustachian tubes are constricted so that air does not readily enter the tympanic cavities, the surgeon should inflate the ears by Politzer's method or by catheterization, while the same general treatment should be carried out as has been indicated in the chapter on Chronic Catarrhal Otitis Media. If the external auditory canals have become very much contracted from suppurative inflammation of the middle ear, which has been due to scarlet fever or diphtheria, an attempt should be made to dilate the canal and to treat the case on general surgical principles according to the indications.

When the labyrinth has become involved, owing to cerebro-spinal meningitis or some other disease, the patient should be given strychnine or other nerve tonics, and strict attention should be paid to the patient's general health. Iodide of potash is indicated in some cases, especially of a syphilitic character. Pilocarpine is a remedy that should be tried if the patient's general health is good, and the deafness has been acquired and is due to labyrinthine disease. It is most important to administer this drug very cautiously, on account of its depressing action on the heart; and when given, it should be continued if possible for a period of at least six weeks. In a case[1] of extreme deafness in an adult, caused by typhoid fever, in which the patient was unable to hear the voice through a speaking-trumpet with one ear, it was necessary to give pilocarpine hypodermically

[1] Bacon: New York Medical Journal, July 24, 1897.

for a period of six weeks before any marked improvement took place. It is wiser to discontinue the treatment for a few days should the patient become very weak.

INSTRUCTION AND EDUCATION OF DEAF-MUTES.

There are two methods or systems by which deaf-mutes are educated, one the so-called French system of instruction, consisting in an artificial finger-alphabet, with a series of signs and gestures, and the other the pure oral or German method, which consists in the employment of articulate speech and lip-reading. At the International Congress held at Milan in 1880 it was the unanimous opinion that the German method was the better one, and at the present time lip-reading or speech-reading is generally taught not only in Europe, but also in the United States, to the exclusion of the sign language.

The German method necessitates great skill and patience on the part of both teacher and pupil. In order to become proficient in lip-reading, it is first of all necessary that the child possess ordinary intelligence and normal vocal organs. A child who is a congenital deaf-mute should be at least seven years of age before such instruction is begun, and, in order to be able to understand and communicate perfectly with others, this teaching should be continued for seven or eight years.

Some deaf-mutes are taught to speak very distinctly, while with others it is almost impossible to understand what they say. As it is difficult for deaf-mutes to modulate the voice, there is often an unpleasant harshness of speech.

The French system is employed occasionally in the

case of deaf-mutes who are deficient in intelligence and who cannot master the oral method. It is often impossible to teach lip-reading to those who have been deaf for many years, and who are accustomed to using signs.

When a child has suddenly become deaf from some general disease, such as scarlet fever or diphtheria, it is important for the patient himself to read aloud for an hour or two each day in order not to lose the art of speech. For all who have recently become deaf, and who cannot be cured by treatment, whether children or adults, an attempt should be made at once to teach lip-reading, provided, of course, the child is old enough. When speech has once been acquired, it is much easier to instruct the patient than when the deafness is congenital.

The German method, or the science of lip-, speech-, or expression-reading, consists in teaching deaf-mutes to watch the lips and tongue, as well as to produce first of all the simple articulate sounds, and eventually the more complex ones.

After being instructed in all the sounds necessary to the language, the pupil is then taught how to form words, and the meaning of such words is conveyed to the child by means of pictures, models, etc. The use of signs should be prohibited, for otherwise the pupil will not be apt to master the German method, since the former is much easier to learn than the latter and requires much less patience and perseverance on the part of the patient. The general education of the child should be carried on at the same time. An intelligent and observing pupil soon learns to notice not only the movement of the lips, but also the position of the tongue, the general shape of the mouth, and the facial

expression. He also watches the brow expression, the twinkle in the eye, and the involuntary movement of the head.

Prof. Denker[1] has reported that "in the Bavarian Senate, it was voted that, owing to the favorable results obtained for a number of years in Munich by separate instruction, deaf-mutes are to be collected in separate classes and instructed by a method in which the eye and ear are equally employed; and it would seem desirable that, in the various institutions for deaf-mute children, separate divisions should be formed for those with auditory remnants."

Many persons of a neurasthenic temperament, and who are quite deaf, owing to chronic catarrhal otitis media, and who consequently are unable to take part in general conversation, occasionally become morose and despondent. These unfortunate patients, after receiving instruction in lip-reading or speech-reading, are often enabled to resume business and to become again useful members of society, while the feeling of despondency and isolation gives place to renewed interests and ambitions.

EAR-TRUMPETS AND OTHER AIDS TO HEARING.

Ear-trumpets are of service in certain forms of deafness, more especially when due to chronic catarrh of the middle ear, in which the drumhead and ossicular chain are present. In those cases in which the labyrinth is affected and the function of the auditory nerve is destroyed completely, an ear-trumpet is generally useless. As the object of the hearing-instrument is to collect the sound-waves and conduct them to the auditory apparatus, it stands to reason that the efficiency of the

[1] Archives of Otology, December, 1904.

instrument depends on the surface by which the sound-waves are collected and reflected to the drumhead.

When two persons desire to carry on a conversation, the best instrument for this purpose is the "conversation-tube" (Fig. 132), consisting of a tube about four feet in length, with a bone tip at one end for insertion into the external meatus, while at the other end there is a mouth-piece, shaped like a cup. In talking into the mouth-piece the person speaking should hold the

FIG. 132.



Conversation-tube.

mouth close to it, and should at first speak in an ordinary tone of voice, for loud sounds spoken through the tube are sometimes distressing to the patient and cause giddiness.

For the purpose of hearing at great distances various trumpet-shaped instruments have been devised (Fig. 133). As the external meatus occasionally becomes irritated from inserting the conical end in the canal, an otophone has been constructed (Fig. 134) whereby this annoyance is prevented. The sound-waves in this instrument are transmitted through a rubber disk which

is stretched tightly. Very small trumpets are useless, as well as the various tubes which are introduced into the canal. There is an exception to this rule, and that is in the case of a collapsed meatus, a subject already

Fig. 133.

Ear-trumpet.

referred to. In aged persons the walls of the meatus sometimes collapse, so that it is difficult for the sound-waves to be transmitted to the drumhead. In such cases a silver tube is useful in holding the walls of the

Fig. 134.

Otophone.

meatus apart. Rhodes's audiphone consists of a flexible fan-shaped plate of vulcanite which has been lacquered, and which has a handle attached. By pressing the upper border of the instrument against the upper incisor and canine teeth, a curved surface is presented, with the

convexity forward. The sound-waves are collected by the audiphone and carried through the teeth and bones of the skull to the internal ear. This instrument has a very limited use, and is never recommended unless the patient has a good set of teeth.

Politzer has devised an instrument the "principle of which depends upon the vibration of the cartilaginous plate of the auricle being transmitted by an elastic sound-conductor to the membrana tympani." It consists of a narrow drainage-tube, 4 to 5 mm. long and 2 mm. thick; one end, which is clipped, being brought into contact with the drumhead, while the other end, which lies in the concha, is in contact with the cartilage of the auricle. The effectiveness is increased by having a round rubber plate near its outer end. The instrument is of most benefit in cases of middle-ear catarrh when the membrana tympani is flaccid, but is useless in cases of rigidity and anchylosis of the ossicula.

APPENDIX.

METHOD OF PREPARING SMEARS.

All preparations of pus for the purpose of bacteriological examination should be as thin as possible. The best method is to cleanse thoroughly two medium thick new cover glasses, obtain and place in the centre of one a drop of pus, place the other cover over it, compress the two between the thumb and forefinger of the left hand sufficiently to secure a uniformly thin layer of the secretion, remove the surplus around the edges with a small piece of filter paper, gauze or other material which can be destroyed, slide one cover over the other until the edge of one projects beyond the edge of its companion, then grasp the upper cover between the thumb and forefinger of the right hand, grasp the under cover in a similar manner with the left hand, and continue the sliding movement until the covers are completely separated. They should then be laid down, prepared surface up, and allowed to dry in the air. When thoroughly dry they are ready for the bacteriologist.

If a drop of pus cannot be obtained, the next best method is to secure the specimen desired by means of cotton wound on the end of a cotton carrier or tooth pick, and then smear it over the surface of the covers or clean new slides as thinly and evenly as possible and allow them to dry as above.

CULTIVATIONS.

Almost any antiseptic, including boric acid, will inhibit the growth of germs; consequently considerable care must be used in making cultivations from the canal or tympanum. If the canal has been irrigated with an antiseptic solution, and a cultivation is desired, the canal should be irrigated carefully with warm sterile water, the meatus then being closed with cotton, and allowed to remain undisturbed for several hours, The cotton is then to be removed, the canal thoroughly cleansed with dry sterile cotton, and the pus for the cultivation removed from the tympanum, or blown through the perforation by the patient, who has been directed to blow his nose. If this fails on account of the tube being closed, the pus can be drawn through by exhausting the air in the canal with the Delstanche masseur. The next step is to make the inoculation. The surgeon should have two culture tubes, one of solid media such as blood serum or agar, and the other of bouillon, and two sterile cotton-tipped wire rods such as are supplied by the Boards of Health for diphtheria cultivations. A separate rod should be used for each tube.

If the case "has not perforated" the method of procedure will differ slightly provided it is desirable to make the cultivation at the time of the paracentesis. In this event, the canal should be sterilized, then irrigated with warm sterile water, and thoroughly dried with sterile dry cotton. An assistant should hold the cotton-tipped sterile rod in a position to enable him to hand it to the operator quickly without contamination, so that the moment paracentesis has been done the operator can obtain a specimen of pus escaping from the tympanum

before hemorrhage has obscured the field of operation. The utmost care is necessary to insure success, as Dixon states that failure has frequently resulted from mopping out the canal with alcoholic bichloride previous to paracentesis, and making cultivations as much as an hour after irrigations with so mild an antiseptic as boric acid.

It is seldom possible to demonstrate the presence of the Klebs-Löffler, or the tubercle bacillus in secretions from the ear without cultivation in the case of the former, and the physiological experiment in the case of the latter.

PHYSIOLOGICAL INOCULATION EXPERIMENT.

The inoculation experiment will frequently fail though we are morally certain that tuberculosis exists.

Failing to demonstrate the presence of the tubercle bacillus by means of cover-glass preparations of the pus, the physiological experiment is to be undertaken as follows: Prepare the canal as above, have ready a healthy rabbit with a cocainized eye, and instruments as follows, *sterilized by boiling;* one pair of smooth flat fine-pointed forceps, a pair of fine-pointed ear scissors, and a set of ear specula, for the purpose of removing suspicious tissue, and a fine flat-pointed, rubber-tipped pipette or dropper for the purpose of removing pus for the inoculation, and for the rabbit a pair of fixation forceps and a keratome. Ocular specula are generally too large to be used on the eye of a rabbit; therefore the lids will have to be separated by an assistant. The rabbit can be held by a second assistant who will grasp the ears with one hand, the hind legs with the other, and stretch the animal which will then be helpless, or its body and legs can be confined in a large towel when one

assistant can hold the animal by the ears with one hand and separate the lids with the other. The operator having obtained the tissue or pus required will make as small an incision as will admit the material through the cornea into the anterior chamber. This incision is best made at the upper limbus and the tissue or pus should be deposited in the anterior chamber as low down as possible while firmly holding the globe with the fixation forceps, otherwise the animal will subsequently be able to remove the offending material through the corneal wound by rubbing his eye, or the iris may force it out by retraction, if it be left near the wound. The animal is then to be penned without dressing. During the next twenty-four to thirty-six hours there is generally violent reaction. This will soon subside, and the eye may become almost normal, but about the beginning of the third week, if the material was tubercular, either a tubercular iritis or panophthalmitis will become apparent. The eye should not be removed for examination before the end of the third week, and a negative report should not be given before the end of the fourth or fifth week.

INDEX.

ABSCESS of brain, 370, 376, 377, 378
  cerebellar, 382
  cerebral, 373, 376
  extradural, 373
  intradural or cerebral, 376
  subdural, 373
  temporo-sphenoidal, 376, 378
Acute circumscribed inflammation of the external auditory canal, 123
  diffuse inflammation of the external auditory canal, 120
Adenoid growths, 186
  causation, 188
  diagnosis of, 189
  symptoms of, 187
  treatment of, 189
Adenoids, excision of, causing hemorrhage, 193, 194
  followed by mastoid disease, 194, 339
  improvement in hearing after, 195
  removal of, steel finger-nail for, 190, 193
Alcohol treatment in chronic purulent otitis media, 270
Allen's foreign-body forceps, 142
Ampulla, 57
Anæmia of labyrinth, 443
  causes of, 443
  symptoms of, 443
  treatment of, 443
Anatomy of ear, 30
Angiomata, 292
Annulus tympanicus, 35
Antihelix, 32
Antitragus, 32
Antrum, disease of, excision of ossicles in, 358
Antrum, guide for opening the, 349
  mastoid, 44
Aphasia, 380, 382
Apparatus for generating ozone, 235
Aquæductus cochleæ, 59
  Fallopii, 44
  vestibuli, 57
Artery forceps, 346 (Plate IX.)
Artificial aids to hearing, 464
  leech, 177
Asch's compressing forceps, 215
  operation, 214
  scissors, 214
  separators, 214
  tubes, 215
Atheroma of auricle, 115
Attic, 44, 47, 167
  acute inflammation of, 167
Attollens aurem muscle, 32
Attrahens aurem muscle, 32
Auditory canal, external, 119
    at birth, 36
    diseases of, 119
    eczema of, 132
    foreign bodies in, 138
    furunculosis of, 123
    herpes zoster of, 132
    inflammation of, acute circumscribed, 123
      diffuse, 120
    chronic diffuse, 121
    croupous, 127
    diphtheritic, 128
    malformation of, 119
      operation for, 119
    normal appearance of, 77
    polypi of, 132
    sebaceous tumors of, 133
    seborrhœa of, 132

Auditory canal, syphilis of, 129
nerve, 64
diseases affecting the, 456
causes of, 456
Aural syringe, 136
vertigo, 444
Auricle, 31
abscess of, 115
atheroma of, 115
blood-supply of, 37
congenital deformities of, 100, 101, 102, 103
cysts of, 108
deformity of, from chronic eczema, 112
deposition of urates in, 115
dermatitis of, 109
dermoid cysts of, 115
description of, 31
eczema of, 112
erysipelas of, 110
erythema of, 115
fibroma of, 116
frost-bite of, 111
function of, 64
herpes zoster of, 115
hyperæmia of, 109
injuries of, 104
keloid of, 115
lupus of, 115
malformations of, 100
operation for, 102
malignant growths of, 115
molluscum fibrosum of, 115
myxofibroma of, 115
nervous distribution of, 37
ossification of, 115
porrigo contagiosa of, 115
sebaceous cyst of, 115
syphilitic diseases of, 118
tinea tonsurans of, 115
tumors of, 115

BISTOURY for incising drum-head, 180
Blake's Eustachian catheter, 87
middle-ear syringe, 274
operating otoscope, 75
paper disks, 283
polypus snare, 298
Blake's rubber spring, 242
tuning-fork, 95
Blood-clot in mastoid disease, 353
-supply of auricle, 37
of external meatus, 37
of tympanum, 49
Boilermaker's deafness, 453
causes of, 453
Boucheron's speculum, 74
Brain, abscess of, 370, 376
causes of, 370
course of, 388
diagnosis of, 389
intermediate path, 398
leucocytosis of, 387
pathology of, 370
prognosis of, 388, 391
symptoms of, 378
temperature chart in, 379
treatment of, 391
venous supply of, 400
Brunton's otoscope, 72

CANAL, external auditory, 34
of Huguier, 46
Canalis centralis modioli, 59
reuniens, 63
spiralis modioli, 59
Carcinoma of temporal bone, 173
Caries of posterior bony wall of external meatus, 361
Carotid artery, 312
Catarrh, 203
chronic, 203
atrophic, 206
danger of forcible inflation in, 207
treatment of, 207
hypertrophic, 204
causes of, 204
symptoms of, 205
treatment of, 206
nasopharyngeal, 208
causes of, 208
symptoms of, 208
treatment of, 208
Cat's ear, 102
Cerebellar abscess, 382
facial paralysis, 383
pathology of, 372
prognosis of, 391

Cerebellar abscess, symptoms of, 383, 384, 385
treatment of, 396
Cerebral abscess due to acute otitis media, 370
facial paralysis due to, 382
percussion of skull in, 382
with chills and high temperature, 380
fossa, middle, large carious opening in, 303
veins and sinuses, 400
Cerebro-spinal meningitis, source of infection in, 437
Cerumen inspissatum, 133
diagnosis of, 134
prognosis of, 135
symptoms of, 134
treatment of, 135
Cholesteatoma, 285, 355
Chorda tympani nerve, 46
Chronic diffuse inflammation of external auditory canal, 121
Clicking tinnitus, 256
Cochlea, 58
cupola of, 59
human, Plate IV, Fig. 1
rabbit, Plate II
Colin's nasal speculum, 81
Concha, 32
Conversation-tube, 465
Corti, membrane of, 60
organ of, 60, 61
Cotton-carrier, aluminium, 76
Crista basilaris, 60
helicis, 32
spiralis, 59
vestibuli, 57
Croupous inflammation of external auditory canal, 127
Cultivations, 470
Cupping-glass, 178
Curette forceps, 75

DEAF-MUTISM, 458
causes of, 458
instruction and education. 462
pathology of, 459
prognosis of, 460
treatment of, 461
Deafness due to collapse of cartilaginous meatus, 456
due to old age, 456
prognosis of, 456
from quinine, 160
Decalcified chicken bone drainage-tube, 396
Deiter's cells, 61
Dermatitis of auricle, 109
treatment of, 109
Dermoid cysts of auricle, 115
Diagnostic tube, 83
Differential diagnosis between perichondritis auriculæ and hæmatoma, 107
Diphtheritic cast from the middle ear, 331
inflammation of the external auditory canal, 128
Diplacusis, 156
Disease of middle and internal ears from mumps, 450
diagnosis of, 450
treatment of, 450
Drumhead, bistoury for incising, 180
danger of rupturing, from forcible inflation, 91
perforation of, 283
treatment of, 283
by nitrate of silver, 283
by paper disks, 283
by scarification, 283
rupture of, from diving, 149
Dry method of treatment in chronic purulent otitis media, 270
Ductus cochleæ, 59

EAR, anatomy of, 30
-cough, 143
external, 31
foreign bodies in, 138
persistent cough due to, 139
prognosis of, 139
symptoms of, 139
treatment of, 139
internal, 56
affections of, 436
diseases of, 436

Ear, internal, diseases of, causes of, 436, 439
diagnosis of, 440, 442
due to injuries, 438
due to microörganisms, 437
prognosis of, 439
symptoms of, 439
middle, danger of removal of polypi from, 371
dry catarrh of, 219
examination of, 82
granulations of, 289
treatment of, 295
by sharp curette, 295
inflammation of, chronic purulent, 257
inflation of, Gruber's method, 86
Politzer's method, 83
Valsalvian method, 83
polypi of, 289
diagnosis of, 294
prognosis of, 295
symptoms of, 293
treatment of, 298
by avulsion, 299
by galvano-cautery, 300
with polypus snare, 298
varieties of, 290
angiomato, 290, 292
fibromata, 292
mucous, 290
myxomata, 292
physiology of, 30, 64
syringe, 135, 136
syringing of, basins for, 137
trumpets, 464
Eczema, 112
artificial, from use of improper ear-drops, 113
intertrigo in children, 112
of auricle, 112
symptoms of, 112
treatment of, 113
varieties of, 112
of external auditory canal, 132
Electric light and portable battery, 70
motor and rheostat, 71
Electricity in Ménière's disease, 415
Encephaloscope, Whiting's, 395
Endolymph, 62
Erysipelas of auricle, 110
causation of, 110
treatment of, 111
a complication of acute purulent otitis media, 171
Ether instead of chloroform in mastoid operations, 345
Eustachian catheter, 87
Blake's, 87
obstacles to introduction of, 91
tube, 50
blood-supply of, 53
catheterization of, 88
examination of, 82
orifices of, 203
functions of, 65
narrowing of, 243
diagnosis of, 243
treatment of, 244
by bougies, 244
osseous, (Plate I.), 51
stricture of, 243
Exostoses of external meatus, 143, 145
diagnosis of, 146
treatment of, 146
External auditory canal, 34, 119
isthmus of, 39
ear, 31
meatus, 34
Extradural abscess, 373
facial paralysis in, 374
pathology of, 373
symptoms of, 374
treatment of, 375

FACIAL nerve, 55
disease of, 315
paralysis of central origin, 316
due to extradural abscess, 374
due to malignant disease, 174
operation for cure of, 308
Fenestra ovalis, 43, 56
Fibroma of auricle, 116
Fibromata, 211, 292

Fibromata, diagnosis of, 212
symptoms of, 211
treatment of, 212
Fibro-myxomatous polypi, 210
symptoms of, 210
treatment of, 211
Fissures of Santorini, 34
Fistula auris congenita, 104
Floor cells, 61
Foramen ovale, 44, 56
rotundum, 44
stylo-mastoid, 44
Forceps, foreign-body, 76, 142
Fossa of antihelix, 32
subarcuata, 319
Fovea hemispherica, 56
Free incision in drumhead, 179
Frog face, due to a fibroma in nasopharynx, 211
Frost-bite of auricle, 111
treatment of, 111
Furunculosis of external auditory canal, 123

GALTON whistle, 96, 456
Ganglion spirale, 59 (Plate II.), 64 (Plate IV., fig. 2)
Gottstein curette, 191
Gradle forceps, 195
Gruber's method of inflation of middle ear, 86
speculum, 73
tenotome, 248

HÆMATOMA, appearances of 106
auris, or othæmatoma, 106
causes of, 106
deformity resulting from, 106
treatment of, 106
Hair cells, outer and inner, 61
Hammer, percussion, 95
Head-mirror, 68
Hearing distance, 93
tuning-fork for testing the, 94
watch for testing the, 92
Helicotrema of Breschetius, 59
Helix, 32
Hemorrhage after excision of tonsils, 200
after removal of adenoids, 193
Hemorrhage, from ear, 143
into labyrinth, 443
Hensen's cells, 62
Herpes zoster of auricle, 115
of external auditory canal, 132
Hewitt's apparatus for washing out the attic, 275
incus-hook, 279
Hooper's forceps, 191
method of removing adenoids, 190
Horseley's instrument for exploring brain, 394
Hydrops ex vacuo, 161
Hyperæmia of auricle, 109
treatment of, 109
of labyrinth, 441
causes of, 441, 442
diagnosis of, 441
symptoms of, 442
treatment of, 442
Hyperæsthesia acustica, 229, 457
Hyperostoses of external meatus, 144

INCISURA intertragica, 32
Incus, 45
Infarctions due to sinus thrombosis, 404
Inflation, forcible, danger of rupture of drumhead from, 91
Influenza a cause of mastoid disease, 324, 325, 327
Ingrowing hairs in external meatus, 143
Inoculation experiment, physiological, 471
Inspissated cerumen, 133
Instruments, mastoid, (Plates IX and X.)
Internal ear, 56
jugular vein, excision of, 417, 418
ligation of, 410, 411
thrombosis, 417
Intracranial complications, 370
leucocytosis in, 387
Intravenous injection of normal saline solution (Hare), 421
Iodoform gauze as a dressing in mastoid disease, 352

JACOBSON'S nerve, 49
Jugular bulb, accidental incision of, 182
primary thrombosis of, 419
fossa, 43, 312

KELOID of auricle, 115
Knife for incising membrana tympani and for dividing adhesions, 278
Knight forceps, 190, 191, 192
König rods, 96, 441, 456
Küster's method, 361

LABIUM tympanicum, 59
vestibulare, 59
Labyrinth, 56
anæmia of, 443
blood-supply of, 63
concussion of, due to injury, 151
disease of, due to action of, drugs, 453
hemorrhage into, 443
hyperæmia of, 441
inflammation of, 445
suppurative, 429
membranous, 62
operation on, 449, 450
pressure, increased, 443
syphilis of, 450
tabes of, 457
Labyrinthine disease due to action of drugs, 453
Lamina spiralis, 58
membranous, 59
osseous, 58
Laminated epithelial plugs, or keratosis obturans, 134
Leiter coil, 334
Leiter's electric lamp, 71
Leptomeningitis, 423
acute, due to cerebral abscess, 388
purulent, 423, 424
diagnosis of, 425
lumbar puncture in, 425
prognosis of, 426
symptoms of, 424
temperature chart, 427
treatment of, 426
serous, 423
Leucocytosis, 386
Levator palati muscle, 52
Ligamentum spirale, 59
Ligation of internal jugular vein, 410, 411, 417, 418
Lobule, 31
cleft, due to wearing brass earrings, 104
Löffler bacillus in middle-ear disease, 331
Lucae's pressure probe, 240
Lumbar puncture, 425, 449
Lupus of auricle, 115
Luschka's tonsil, 187
Lymphatics of scalp, 399

MACEWEN'S burrs, 349
Mackenzie's bull's-eye lens and reflector, 69
tonsillotome, 199
Macula acustica, 62
Malignant disease a cause of acute purulent otitis media, 173
Malleus, 44
excision of handle of, 249
fracture of handle of, 153
short process of, 40, 78
Massage, vibratory, 239
Hommel's, 240
Masseur of Delstanche, 79
Mastoid abscess in children, 329
antrum, 44, 312, 319
cells, inflammation of, acute, due to acute suppurative otitis media, 323
causes of, 323
course of, 327
microörganisms in, 330
prognosis of, 331
symptoms of, 324
treatment of, 332
primary acute inflammation of, 321
causes of, 321
differential diagnosis of, 322

Mastoid cells, primary acute inflammation of, prognosis of, 322
symptoms of, 322
treatment of, 323
disease, blood-clot in, 353
due to chronic purulent otitis media, 355
causes of, 356
diagnosis of, 358
prognosis of, 358
symptoms of, 356
treatment of, 358
danger from use of blisters in, 333
diagnostic symptoms of, 325
due to diabetes, 360
due to exanthematous diseases, 335, 336
due to removal of adenoid growths, 339
due to teething, 339
early operation in, 328
facial paralysis in, 328
following excision of adenoids, 194
free incision of drumhead in, 329, 332, 334
iodoform gauze dressing in, 352
microörganisms in, 330
secondary operation in 353
instruments, Plates IX. and X.
operation, 345
indications for, 342
lines of incision in, 347
mastoid exposed, 348
preparation of patient for, 345
operations, injury to dura in, 350
to external semicircular canal, 350
to facial nerve, 350
to sigmoid sinus in, 350
lines of incision in, 347
periosteum, inflammation of, primary, 319
causes of, 320
symptoms of, 320
treatment of, 321
by artificial leech, 321
Mastoid periosteum, inflammation of, primary, treatment, by Leiter coil, 321
process, 54, 310
acute inflammation of, 323
anatomy of, 311
in children, 311
danger of delay in operations on, 343
diploëtic variety of, 54, 317
diseases of, 310
pneumatic diploëtic variety of, 317
varieties of, 54, 317
primary acute inflammation of, 321
sclerosed, 360
treatment of, 360
sequestra from, 306, 307
tip, tenderness on pressure over, 343
vein, 315, 399
Mastoiditis externa, 319
interna, 321
Masto-squamosal suture, 329
Mathieu tonsillotome, 199
Meatus, external, adhesions of, 143
blood-supply of, 37
contractions of, 143
exostoses in, 143, 145
hyperostoses in, 144
ingrowing hairs in, 143
isthmus of, 39
nervous distribution of, 37, 38
or external auditory canal, 34
sarcoma attached to, 290
Membrana basilaris, 60 (Plate III.)
reticularis, 61
tectoria, 60
tympani, 39, 77
anterior folds of, 78
artificial, 280
opening in, 246
flaccida, or Shrapnell's membrane, 42, 78
injuries of, 148, 149
diagnosis of, 150
prognosis of, 151
treatment of, 152

Membrana tympani, normal, 40
posterior fold of, 78
progressive growth of dermoid layer of, 253
section of posterior fold of, 247
vibrans, 78
Membrane of Corti, or membrana tectoria, 60
of Kölliker, 61
of Reissner, 60
Membranous labyrinth, 62
Ménière's disease, 446
causes of, 446
prognosis of, 448
symptoms of, 446
treatment of, 448
Method of incising the drumhead when bulging, 180
Meyer's annular knife for excision of adenoid growths, 189
Microörganisms in acute purulent otitis media, 179
Microtia, 102
Middle ear, 39
granulations and polypi of, 289
paths of infection of, 313
tuberculosis of, 266
Molluscum fibrosum of auricle, 115
Mouth-breathing due to adenoids, 188
Mucous polypi of middle ear, 290
Mumps, 450
Muscle, attollens aurem, 32
attrahens aurem, 32
of auricle, 32
levator palati, 52
retrahens aurem, 33
stapedius, 48
tensor palati, 52
tympani, 45, 48
Myringitis, acute, 148
symptoms of, 148
treatment of, 149
chronic, 149
symptoms of, 149
treatment of, 149
Myxofibroma of auricle, 115
Myxomata, 292

NASAL passages, appearance of normal, 80
diseases of, 185
polypi, 209
treatment of, 210
septum, cartilaginous, deviation of, 212
diagnosis of, 213
symptoms of, 213
treatment of, 214
Nasopharynx, 158
appearance of normal, 80
digital examination of, 158
Nerve, auditory, 64
chorda tympani, 46
-deafness, 436
facial, 55
Jacobson's, 49
supply of auricle, 38
of external meatus, 38
of tympanum, 49
New growths affecting the inner ear, 457
Normal saline solution, intravenous injection of, 421
Nose, examination of, 200

OSSEOUS erosion of tympanic roof, 370
meatus, 35
Ossicles, or ossicula, 45
excision of, 245, 249, 284, 358
Ossification of auricle, 115
Osteomyelitis of the temporal bone, 340
glycosuria as a complication of, 341
Otalgia, 255
Othæmatoma, 106
Otitis externa circumscripta acuta, 123
causes of, 123
diagnosis of, 125
symptoms of, 123
treatment of, 125
crouposa, 127
diffusa acuta, 120
causes of, 120
symptoms of, 120

Otitis externa circumscripta, diffusa acuta, treatment of, 122
chronica, 121
diphtheritica, 128
hæmorrhagica, 126
interna or inflammation of labyrinth, 445
causes of, 445
diagnosis of, 445
media acuta, or acute inflammation of middle ear, 153
acute catarrhal, 153
purulent, 164
catarrhalis acuta, 153
appearance of membrana tympani in, 153, 157
causation of, 154
symptoms of, 155
treatment of, 158
chronica, 219
exudative variety of, 219
sclerotic variety of, 219
adhesions in, 221
anchylosis of ossicles in, 222
appearances of membrana tympani in, 225
atrophic changes of drumhead in, 227
causes of, 224
diagnosis of, 230
heredity in, 224
obliteration of attic in, 221
prognosis of, 231
symptoms of, 227
treatment of, 233
subacuta, 160
treatment of, 162
chronic catarrhal, 219
purulent, 257
purulenta acuta, 164
appearance of membrana tympani in, 166
complications of, 169
course of, 169
diagnosis of, 172
etiology of 165
free incision in, 179
Otitis media, purulenta acuta, microörganism in, 179
pathology of, 164
permanent disturbances of hearing in, 171
prognosis of, 174
symptoms of, 168
treatment of, 176
use of powders in, 182
chronica, 257
anatomical appearances of, 258
appearances of external auditory canal in, 261
of membrana tympani in, 261
character of discharge in, 260
diagnosis of, 265
perforation in Shrapnell's membrane in, 263
prognosis of, 265
symptoms of, 259
treatment of, 266
by Blake's middle-ear syringe, 273
by caustic applications, 271
by excision of carious ossicles, 277
by Hewitt's apparatus, 275
by instillation of alcohol, 270
of astringent solutions, 269
by insufflation of powders, 270
by sharp spoons, 273
by syringing through Eustachian tube, 268
by wick method, 272
subacute catarrhal, 160
Otoliths, 62
Otomycosis, 129
aspergillus flavus, 129
fumigatus, 129
niger or nigricans, 129

Otomycosis, etiology of, 129
symptoms of, 130
treatment of, 131
Otophone, 466
Ozæna, 207
treatment of, 207

PARACUSIS Willisiana, 229
Parietal clot, 402
Parotiditis due to inflammation of external meatus, 311
Percussion hammer, 95
Perichondritis auriculæ, 107
appearances of, 107
deformity resulting from, 108
treatment of, 108
Perilymph, 62
Petroso-squamosal suture, 42, 354, 371
Pharynx, appearance of normal, 80
Physiology of hearing, 64
Physiological inoculation experiment, 471
Plexus, tympanic, 49
Pneumococci a cause of mastoid disease, 414
Pneumonia a complication of mastoid disease, 337
Politzer's acoumeter, 95
artificial membrana tympani, 281
bag, 85
eyelet, 247
forceps, 278
head-mirror, 68
method of inflation of middle ear, 83
of removing fluid from tympanum, 162
ring-knife, 296
sharp spoons, 273
speculum, 73
Polyotia, 100
operation for, 102
Polypi, fibro-myxomatous, 210
mucous, 209
nasal, 209
of external auditory canal, 132
of naso pharynx, 210
Porrigo contagiosa of auricle, 115
Primary disease of bony labyrinthine capsule, 454
symptoms of, 455
treatment of, 455
jugular bulb thrombosis, 421
papillitis in, 420
Processus cochleariformis, 47
Prominentia spiralis, 60
Promontory, 43
Prussak's space, 46, 47
Pyæmia, 433
causes of, 434
symptoms of, 434
Pyramid, 44

RAREFACTEUR, or masseur of Delstanche, 79
Recumbent method of removing adenoids, 192
Reid's base line, 384
Reissner, membrane of, 60
Report of case of sinus thrombosis in child, 408
Retrahens aurem muscle, 32
Rhinitis, 204
atrophic, 206
chronic atrophic, 206
nasopharyngeal, 208
hypertrophic, 204
Rhinoscopy, anterior, 200
posterior, 202
mirror for, 81
Rhodes' audiphone, 466
Rinné's test, 97
Rivinian segment, 35
Rongeur forceps, 346 (Plate X., Fig. 1) 397

SACCULE, 62
Saline solution, normal, 421
Scala media, 59 (Plate II.)
tympani, 59
vestibuli, 59
Scarificator, 177
Schwabach's test, 97
Schwartze's chisels and gouges, Plate IX.
Schwartze-Stacke operation, 366
Thiersch grafts in, 369

Sclerotic otitis media, 224
Sebaceous cyst of auricle, 115
Seborrhœa of external auditory canal, 132
Semicircular canals, 57
functions of, 65
Septicæmia, 433
causes of, 434
symptoms of, 435
Sero-mucous catarrh, 161
prognosis of, 162
symptoms of, 162
treatment of, 162
Sharp curettes, 296
spoons, 346 (Plate X., Fig. 1)
Schrapnell's membrane, 42, 78
perforation in, 263, 273
Siegle's pneumatic speculum, 79
Sigmoid groove, defects in, 319
sinus, 360
inadvertent exposure of, 351
knee or anterior convexity of, 399
Sinus thrombosis due to acute otitis media, 413
re-establishment of blood current in, 418
time to operate in, 412
Sinuses, cerebral veins, 400
Smears, method of preparing, 469
Sound-conducting apparatus, 30
-perceiving apparatus, 30
diseases of, 436
epidemic cerebro-spinal meningitis in, 437
Specula, aural, 73, 74
Spina suprameatum, 55
Stacke operation, 362
modifications of 368
Stapedectomy, 249
alarming symptoms following, 252
Stapedius muscle, 48
Stapes, 45, 222
excision of, 249
Steel finger-nail for removing adenoids, 192, 193
Sterilizer, 345
Stoker's ozone apparatus, 235
Streptococci in brain abscess, 396
in sinus thrombosis, 409
Stria vascularis, 60
Stylo-mastoid foramen, 44
Sulcus spiralis externus, 60
internus, 59
tympanicus, 40
Suppurative inflammation of labyrinth, 434
leptomeningitis due to osseous erosion, 371
Suture, petroso-squamosal, 42
Symptoms of Ménière's disease due to middle-ear disease, 446
Syphilis of external auditory canal, 129
of labyrinth, 450
diagnosis of, 451
prognosis of, 452
symptoms of, 451
treatment of, 452
Syphilitic diseases of auricle, 118

TEETHING a cause of acute purulent otitis media, 171
Tegmen tympani, defects in, 313
Temperature-chart, brain abscess, 379, 381
mastoiditis, acute, and pneumonia, 337
sinus thrombosis, leptomeningitis, 427
thrombosis sigmoid sinus, 407
Temporal bone at birth, 37
caries of, 300
cholesteatoma of, 285
diagnosis of, 287
pathology of, 286
symptoms of, 286
treatment of, 287
malignant growth of, 173
necrosis of, 300
causes of, 301
frequency of involvement of different portions, 329
prognosis of, 305
symptoms of, 304
treatment of, 305
Temporo-sphenoidal abscess, 376, 378
causes of, 376
temperature-chart of, 379
prognosis of, 388

Temporo-sphenoidal abscess, symptoms of, 378
treatment of, 391
Tenotomy of tensor tympani muscle, 248
Tensor palati muscle, 52
Tests for hearing, 91
Thiersch grafts, 369
Thrombosis of sigmoid and other sinuses, 398
course of, 402
diagnosis of, 412
different types of, 411
method of operation, 415
pathology of, 372, 402
prognosis of, 415
symptoms of, 404
time to operate, 412
treatmeni of, 415
Thrombus, 398
internal jugular vein, 417
primary or marasmic, 401
causation of, 401
secondary or infective, 401
causes of, 401
microörganisms in, 401
Tinea tonsurans of auricle, 115
Tongue depressor, 202
Tonsils enlarged, 195
causes of, 196
symptoms of, 196
tonsillotomy for, 198
treatment of, 196
excision of, hemorrhage from, 200
Toxic effects of iodoform gauze, 396
Toynbee's artificial membrane, 281
Tragus, 32
Treatment of acute circumscribed inflammation of external auditory canal, 125
diffuse inflammation of external auditory canal, 122
inflammation of mastoid cells occurring in connection with acute purulent otitis media, 323
of adenoid growths, 189
of adhesions and contractions of external meatus, 145
Treatment of anæmia of labyrinth, 443
of brain abscess, 391
of caries and necrosis of temporal bone, 305
of cerebellar abscess, 396
of cerumen inspissatum, or inspissated cerumen, 135
of cholesteatoma of temporal bone, 287
of chronic atrophic catarrh, 207
hypertrophic catarrh, 206
nasopharyngeal catarrh, 209
of deaf-mutism, 461
of dermatitis of auricle, 109
of deviation of cartilaginous nasal septum, 214
of disease of middle and internal ear due to mumps, 450
of eczema of auricle, 113
of enlarged tonsils, 196
of erysipelas of auricle, 111
of exostoses of external meatus, 146
of extradural abscess, 375
of facial paralysis due to caries of temporal bone, 308
of fibroma of auricle, 116
of fibromata of nasopharynx, 212
of fibro-myxomatous polypi of nasopharynx, 211
of foreign bodies in ear, 139
of frost-bite of auricle, 111
of furunculosis of external auditory canal, 125
of granulations of middle ear, 295
of hæmatoma auris or othæmatoma, 106
of hyperæmia of auricle, 109
of labyrinth, 441
of increased pressure of labyrinthine fluid, 444
of injuries of membrana tympani, 152
of leptomeningitis, 426
of mastoid disease in connection with chronic purulent otitis media, 358
of Ménière's disease, 448

Treatment of myrinigitis, acute, 149
chronic, 149
of narrowing and stricture of Eustachian tube, 244
of nasal polypi, 210
of otitis externa circumscripta acuta, 125
diffusa acuta, 122
media, catarrhalis acuta, 158
chronica, 233
subacuta, 162
purulenta acuta, 176
chronica, 266
of otomycosis, 131
of ozæna 207
of perforation of drumhead, 283
in Schrapnell's membrane 264, 273
of perichondritis auriculæ, 108,
of polypi of middle ear, 298
of primary acute inflammation of mastoid cells, 321
of disease of bony labyrinthine capsule, 455
of primary inflammation of mastoid periosteum, 319
of sclerosed mastoid process, 360
of sebaceous cyst of auricle, 116
of sero-mucous catarrh, 162
of syphilis of labyrinth, 452
of temporo-sphenoidal abscess, 391
of thrombosis of sigmoid and other sinuses, 415
of tumors of auricle, 115
Trephining for abscess in temporo-sphenoidal lobe, 393
for cerebellar abscess, 397
Tubal catarrh, 161
Tuberculosis of mastoid, 321
of middle ear, 266
Tumors, of auricle, 115
sebaceous, of external auditory canal, 133
Tuning-fork, Blake's, 96
-forks, for testing hearing distance, 94
Hartmann's series, 94, 95
Politzer's low fork, 95
Tympanic cavity, 40
epithelial masses in, 285
plexus, 49
Tympanotomy, exploratory, 253
Tympanum, blood supply of, 49
description of, 40
nervous distribution of, 49

UMBO, 41, 77
Urates, deposition of, in auricle, 115
Utricle, 62

VALSALVIAN method of inflation of middle ear, 83
Veins and sinuses, cerebral, 400
Vestibule, 56
Vicarious menstruation, 143
Villate's solution, 306
Viola tricolor in treatment of acute eczema, 113
Volkmann's spoon, 373

WATCH for testing hearing distance, 92
Weber's test, 97
White erysipelas of scalp a complication of mastoid disease, 341, 342
Whiting's encephaloscope, 395

ZONA arcuata, 6, (Plate III.)

MEar
B

Bacon, Gorham
A manual of otology. 4th.ed. rev.and enl.

497368

DATE.

NAME OF BORROWER.

University of Toronto
Library

DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET